

A categorification of the Casimir of quantum $\mathfrak{sl}(2)$

Anna Beliakova, Mikhail Khovanov, and Aaron D. Lauda

January 10, 2011

Abstract

We categorify the Casimir element of the idempotented form of quantum $\mathfrak{sl}(2)$.

Contents

1	Introduction	2
2	Casimir element and idempotented form of quantum \mathfrak{sl}_2	7
2.1	Quantum \mathfrak{sl}_2 and the Casimir element	7
2.2	Idempotented rings and their centers	9
2.3	BLM $\dot{\mathcal{U}}$	9
3	Brief review of $\mathfrak{sl}(2)$-calculus	11
3.1	The 2-category \mathcal{U}	11
3.2	The 2-categories $\dot{\mathcal{U}}$, $Kom(\dot{\mathcal{U}})$, and $Com(\dot{\mathcal{U}})$	18
3.3	Symmetry 2-functors	22
4	The Casimir complex	25
4.1	The Casimir complex and its symmetries	25
4.2	Commutativity of the Casimir complex	32
4.3	Indecomposability	36
4.4	Symmetries of the commutativity chain maps	41
5	Naturality of the Casimir complex	47
5.1	Natural transformations κ and $\widehat{\kappa}$	47
5.2	Naturality with respect to 2-morphisms	49
	References	61

1 Introduction

The Witten-Reshetikhin-Turaev invariant [20, 21] of a 3-manifold presented by surgery along a framed link is given by summing over labellings of the components of the link by irreducible representations of the quantum group $U_q(\mathfrak{sl}_2)$, then evaluating the corresponding coloured Jones polynomial at a root of unity. Crane and Frenkel [6] conjectured that quantum 3-manifold invariants could be categorified using the categorified representation theory of $U_q(\mathfrak{sl}_2)$. While the Jones polynomial and coloured Jones polynomial have been categorified [3, 11], the problem of categorification at a root of unity has seen little progress.

A universal invariant of knots taking values in (an appropriate completion of) the center of the quantum $U_q(\mathfrak{sl}_2)$ was constructed by Lawrence [16, 17]. This universal invariant dominates all coloured Jones polynomials. The center of $U_q(\mathfrak{sl}_2)$ as a $\mathbb{Z}[q, q^{-1}]$ -algebra is freely generated by the Casimir element C . For $p \geq 0$, let us introduce

$$\sigma_p = \prod_{i=1}^p (C^2 - (q^{2i} + 2 + q^{-2i}))$$

which are monic polynomials of degree p in C^2 . The universal invariant J_K of a knot K can be written [8, Theorem 4.5] as

$$J_K = \sum_{p \geq 0} a_p(K) \sigma_p.$$

The coefficients $a_p(K) \in \mathbb{Z}[q, q^{-1}]$, $p \in \mathbb{N}$, determine the Witten-Reshetikhin-Turaev invariant of any integral homology 3-sphere obtained by surgery on the knot K . Therefore it is natural to seek a categorification of this universal invariant within the context of categorified representation theory of $U_q(\mathfrak{sl}_2)$. Here we take a first step in this ambitious program by categorifying the quantum Casimir element for $U_q(\mathfrak{sl}_2)$.

Igor Frenkel conjectured [7] the existence of a categorification of the integral idempotent version ${}_{\mathcal{A}}\dot{\mathbf{U}}$ of the quantum enveloping algebra of \mathfrak{sl}_2 at generic q . The algebra ${}_{\mathcal{A}}\dot{\mathbf{U}}$ is the $\mathbb{Z}[q, q^{-1}]$ -subalgebra of the algebra $\dot{\mathbf{U}}$ defined by Beilinson, Lusztig and MacPherson [2] and generalized to arbitrary types by Lusztig [18]. In $\dot{\mathbf{U}}$ the identity element $1 \in \mathbf{U}_q(\mathfrak{sl}_2)$ is replaced by a collection of orthogonal idempotents 1_n indexed by the weight lattice for \mathfrak{sl}_2 . We recall the definition of $\mathbf{U}_q(\mathfrak{sl}_2)$ in Section 2.1 and $\dot{\mathbf{U}}$ in Section 2.3.

In [15] the third author introduced a categorification $\dot{\mathcal{U}}$ of ${}_{\mathcal{A}}\dot{\mathbf{U}}$ given by the idempotent completion of an additive 2-category \mathcal{U} whose objects $n \in \mathbb{Z}$ are parameterized by the integral weight lattice of \mathfrak{sl}_2 . The 1-morphisms from n to m are given by direct sums of 1-morphisms $\mathcal{E}_{\underline{\varepsilon}} \mathbf{1}_n \{t\} = \mathcal{E}_{\varepsilon_1} \dots \mathcal{E}_{\varepsilon_k} \mathbf{1}_n \{t\}$ where $\underline{\varepsilon} = \varepsilon_1 \dots \varepsilon_k$ with $\varepsilon_1, \dots, \varepsilon_k \in \{+, -\}$, $m - n = 2 \sum_{i=1}^k \varepsilon_i$, $\mathcal{E}_+ = \mathcal{E}$, $\mathcal{E}_- = \mathcal{F}$ and $t \in \mathbb{Z}$. The 1-morphisms \mathcal{E} and \mathcal{F} can be thought of as categorifications of the generators E and F of quantum \mathfrak{sl}_2 . The 2-morphisms are given by \mathbb{k} -linear combinations of certain planar diagrams modulo local relations. In [15] it was shown that the split Grothendieck ring of $\dot{\mathcal{U}}$ is isomorphic to ${}_{\mathcal{A}}\dot{\mathbf{U}}$,

$$K_0(\dot{\mathcal{U}}) \cong {}_{\mathcal{A}}\dot{\mathbf{U}}, \tag{1.1}$$

when the ground ring is a field \mathbb{k} . In [13] this result was proven with \mathbb{k} replaced by the integers.

The quantum Casimir element for $U_q(\mathfrak{sl}_2)$ has the form

$$c := EF + \frac{q^{-1}K + qK^{-1}}{(q - q^{-1})^2} = FE + \frac{qK + q^{-1}K^{-1}}{(q - q^{-1})^2}. \quad (1.2)$$

This element is central and is preserved by various (anti)linear (anti)involutions defined on $U_q(\mathfrak{sl}_2)$. In this paper we categorify the integral idempotent version of the Casimir element obtained from (1.2) by clearing denominators by multiplying by $(q - q^{-1})^2$ and projecting via 1_n . We also multiply by -1 for convenience and obtain the integral idempotent Casimir element \dot{C} for $\dot{\mathbf{U}}$:

$$\dot{C} = \prod_{n \in \mathbb{Z}} C1_n, \quad (1.3)$$

$$C1_n = 1_n C := (-q^2 + 2 - q^{-2})EF1_n - (q^{n-1} + q^{1-n})1_n, \quad (1.4)$$

$$= (-q^2 + 2 - q^{-2})FE1_n - (q^{n+1} + q^{-1-n})1_n. \quad (1.5)$$

This element belongs to the center, defined in Section 2.2, of the idempotent ring $\dot{\mathbf{U}}$.

To categorify the component $C1_n$ of the idempotent Casimir element of $\dot{\mathbf{U}}$ in the form given in (1.4) we must lift elements $q^a EF1_n$ and $q^b 1_n$ to 1-morphisms $\mathcal{EF}1_n\{a\}$ and $1_n\{b\}$. We follow the now standard procedure of lifting powers of q to grading shifts and using complexes whenever minus signs are present. This requires us to work with the 2-category $Kom(\dot{\mathcal{U}})$ of bounded complexes over the 2-category $\dot{\mathcal{U}}$ whose objects are integers $n \in \mathbb{Z}$, 1-morphisms are bounded complexes of 1-morphisms in $\dot{\mathcal{U}}$, and 2-morphisms are chain maps constructed from the 2-morphisms in $\dot{\mathcal{U}}$. We also consider the 2-category $Com(\dot{\mathcal{U}})$ which has the same objects and 1-morphisms as $Kom(\dot{\mathcal{U}})$, and whose 2-morphisms are chain maps up to homotopy.

We are looking for a complex with four copies of $\mathcal{EF}1_n$ and two copies of 1_n with grading shifts:

$$\mathcal{EF}1_n\{-2\}, \quad \mathcal{EF}1_n, \quad \mathcal{EF}1_n, \quad \mathcal{EF}1_n\{2\}, \quad 1_n\{1-n\}, \quad 1_n\{n-1\}.$$

The minus signs in (1.4) indicate that the terms $\mathcal{EF}1_n\{-2\}$, $\mathcal{EF}1_n\{2\}$, $1_n\{1-n\}$, $1_n\{n-1\}$ should live in odd homological degrees, and the remaining two copies of $\mathcal{EF}1_n$ in even degrees.

The positioning of these terms in the complex is naturally dictated by the q -degrees of the possible maps between them. Negative degree endomorphisms of $\mathcal{EF}1_n$ exist only for $n > 1$, while there are obvious degree two endomorphisms given by placing a dot on one of two vertical lines in the diagram of the identity map: $\uparrow \downarrow$ and $\downarrow \uparrow$ (see Section 3.1 for a review of the 2-category \mathcal{U}). We can arrange the above four copies of $\mathcal{EF}1_n$ with the

appropriate shifts and cohomological degrees into a complex just using these maps

$$\begin{array}{ccccc}
 & & \mathcal{E}\mathcal{F}\mathbf{1}_n & & \\
 & \uparrow \downarrow & \nearrow & \searrow & \\
 \mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} & & & & \mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \\
 & \downarrow \uparrow & \searrow & \nearrow & \\
 & & \mathcal{E}\mathcal{F}\mathbf{1}_n & &
 \end{array}$$

where the exact position of the minus sign is unimportant. To find the room for the two shifted copies of $\mathbf{1}_n$ we observe that clockwise cup and cap 2-morphisms have degree $1 - n$, perfectly matching the difference in degrees of these $\mathbf{1}_n$ and those of the middle $\mathcal{E}\mathcal{F}\mathbf{1}_n$ in the complex, leading to a commutative square

$$\begin{array}{ccccc}
 & & \mathcal{E}\mathcal{F}\mathbf{1}_n & & \\
 & \curvearrowright & \nearrow & \searrow & \\
 \mathbf{1}_n\{1-n\} & & & & \mathbf{1}_n\{n-1\} \\
 & \curvearrowleft & \searrow & \nearrow & \\
 & & \mathcal{E}\mathcal{F}\mathbf{1}_n & &
 \end{array}$$

These two commutative squares can be glued into a single complex $\mathcal{C}\mathbf{1}_n$ centered in homological degree zero:

$$\begin{array}{ccccccc}
 & & \mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} & \xrightarrow{\uparrow \downarrow} & \mathcal{E}\mathcal{F}\mathbf{1}_n & \xrightarrow{-\uparrow \downarrow} & \mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \\
 & & \searrow & \nearrow & \curvearrowright & \searrow & \nearrow \\
 \mathcal{C}\mathbf{1}_n := \oplus & & & & \oplus & & \oplus \\
 & & \nearrow & \searrow & \curvearrowleft & \nearrow & \searrow \\
 & & \mathbf{1}_n\{1-n\} & \xrightarrow{\curvearrowleft} & \mathcal{E}\mathcal{F}\mathbf{1}_n & \xrightarrow{-\curvearrowright} & \mathbf{1}_n\{n-1\}
 \end{array} \tag{1.6}$$

We call the above complex the Casimir complex. The image of this complex in the Grothendieck ring of $Com(\mathcal{U})$ is $\mathcal{C}\mathbf{1}_n$ expressed in the form given by the right hand side of equation (1.4).

Starting with the form of the idempotented Casimir element given in (1.5) we obtain a

different complex:

$$\begin{array}{ccccc}
\mathcal{FE}\mathbf{1}_n\{2\} & \xrightarrow{\downarrow \uparrow} & \mathcal{FE}\mathbf{1}_n & \xrightarrow{-\downarrow \uparrow} & \mathcal{FE}\mathbf{1}_n\{-2\} \\
\downarrow \uparrow & \searrow & \downarrow \uparrow & \searrow & \downarrow \uparrow \\
\mathcal{C}'\mathbf{1}_n := \bigoplus & & \bigoplus & & \bigoplus \\
\mathbf{1}_n\{1+n\} & \xrightarrow{\curvearrowright} & \mathcal{FE}\mathbf{1}_n & \xrightarrow{-\curvearrowright} & \mathbf{1}_n\{-n-1\}
\end{array} \quad (1.7)$$

However, we will show that the complex $\mathcal{C}'\mathbf{1}_n$ is homotopy equivalent to $\mathcal{C}\mathbf{1}_n$. These two complexes behave well under certain symmetry 2-functors ψ, ω, σ defined for the 2-category \mathcal{U} in [15], and extended here in Section 3.3 to the 2-categories $Kom(\dot{\mathcal{U}})$ and $Com(\dot{\mathcal{U}})$. In particular, $\mathcal{C}'\mathbf{1}_n = \sigma(\mathcal{C}\mathbf{1}_{-n})$. These symmetry 2-functors categorify certain (anti)linear (anti)involutions on the algebras $\dot{\mathcal{U}}$ with the various (anti)linearity and (anti)involution properties being reflected in the (contravariant)covariant behaviour of the 2-functors. Just as one can go between the two forms of the Casimir in (1.4) and (1.5) using these (anti)involutions on the algebras $\dot{\mathcal{U}}$, we relate the complexes above together with their alternative versions obtained by moving the minus signs and reordering the dot 2-morphisms via these categorified symmetries of $\dot{\mathcal{U}}$.

Our results can be summarized as follows:

Theorem 1.1.

a) There are canonical mutually-inverse isomorphisms

$$\varrho^\sigma : \mathcal{C}\mathbf{1}_n \rightarrow \mathcal{C}'\mathbf{1}_n, \quad \widehat{\varrho}^\sigma : \mathcal{C}'\mathbf{1}_n \rightarrow \mathcal{C}\mathbf{1}_n \quad (1.8)$$

in $Com(\dot{\mathcal{U}})$. If $n \leq 0$ the complex $\mathcal{C}\mathbf{1}_n$ is indecomposable in $Kom(\dot{\mathcal{U}})$, and complex $\mathcal{C}'\mathbf{1}_n$ is isomorphic to a direct sum of $\mathcal{C}\mathbf{1}_n$ and a contractible complex. If $n \geq 0$ the complex $\mathcal{C}'\mathbf{1}_n$ is indecomposable in $Kom(\dot{\mathcal{U}})$, and complex $\mathcal{C}\mathbf{1}_n$ is isomorphic to a direct sum of $\mathcal{C}'\mathbf{1}_n$ and a contractible complex. If $n = 0$ complexes $\mathcal{C}\mathbf{1}_n$ and $\mathcal{C}'\mathbf{1}_n$ are isomorphic in $Kom(\dot{\mathcal{U}})$.

b) Under the isomorphism $K_0(\dot{\mathcal{U}}) \cong K_0(Com(\dot{\mathcal{U}})) \cong {}_{\mathcal{A}}\dot{\mathcal{U}}$ we have $[\mathcal{C}\mathbf{1}_n] = \mathcal{C}\mathbf{1}_n$, so that the complex $\mathcal{C}\mathbf{1}_n$ in $Com(\dot{\mathcal{U}})$ descends to the component $\mathcal{C}\mathbf{1}_n$ of the Casimir element \dot{C} of $\dot{\mathcal{U}}$ after passing to the Grothendieck ring.

c) The complex $\mathcal{C}\mathbf{1}_n$ is invariant under the symmetries ψ and $\omega\sigma$ of $Kom(\mathcal{U})$. Symmetry σ takes $\mathcal{C}\mathbf{1}_n = (1.6)$ to the complex $\mathcal{C}'\mathbf{1}_{-n}$ given by (1.7) for $-n$.

d) *Commutativity*: There exists a collection of invertible 2-morphisms of complexes

$$\kappa_X : XC \rightarrow CX, \quad (1.9)$$

with inverses

$$\widehat{\kappa}_X: \mathcal{C}X \rightarrow XC, \quad (1.10)$$

for all X in $Com(\dot{\mathcal{U}})$.

Naturality: The collection of invertible chain maps κ_X is natural in the sense that for any 2-morphism $f: X \rightarrow Y$ the squares

$$\begin{array}{ccc} Y\mathcal{C} & \xrightarrow{\kappa_Y} & \mathcal{C}Y \\ \uparrow fc & & \uparrow cf \\ X\mathcal{C} & \xrightarrow{\kappa_X} & \mathcal{C}X \end{array} \quad \begin{array}{ccc} \mathcal{C}Y & \xrightarrow{\widehat{\kappa}_Y} & Y\mathcal{C} \\ \uparrow cf & & \uparrow fc \\ \mathcal{C}X & \xrightarrow{\widehat{\kappa}_X} & X\mathcal{C} \end{array} \quad (1.11)$$

commute in $Com(\dot{\mathcal{U}})$. By construction these invertible chain maps are compatible with composition in $Com(\dot{\mathcal{U}})$ given by the tensor product of complexes and juxtaposition of diagrams. That is, for complexes $Y = \mathbf{1}_{n'}Y\mathbf{1}_n$ and $X = \mathbf{1}_{n''}X\mathbf{1}_{n'}$, with $\mathcal{C}XY\mathbf{1}_n = \mathbf{1}_{n''}\mathcal{C}\mathbf{1}_{n''}X\mathbf{1}_{n'}Y\mathbf{1}_n$, we have a commutative diagram

$$\begin{array}{ccc} \mathcal{C}XY\mathbf{1}_n & \xrightarrow{\kappa_{XY}} & XY\mathcal{C}\mathbf{1}_n \\ & \searrow \kappa_{XY} & \nearrow X\kappa_Y \\ & & X\mathcal{C}Y\mathbf{1}_n \end{array} \quad (1.12)$$

in $Com(\dot{\mathcal{U}})$.

Parts a) and d) are difficult, while parts b) and c) are obvious. The indecomposability of the Casimir and the resulting simplifications are discussed in section 4.3. The rest of part a) and part c) of the Theorem can be found in Proposition 4.1. The construction of the commutativity chain isomorphisms is given in Section 4.2, while the naturality of these maps is proven in Section 5.

As explained in [12, Section 3.7] an additive 2-category can be viewed as an idempotent monoidal category by regarding 1-morphisms as objects of the monoidal category. The 2-morphisms in the 2-category become 1-morphisms in the monoidal category. The composition operation for 1-morphisms and the horizontal composition for 2-morphisms in the original 2-category gives rise to the monoidal structure, allowing objects and morphisms to be tensored together. It is sometimes convenient to view $Kom(\dot{\mathcal{U}})$ and $Com(\dot{\mathcal{U}})$ as idempotent additive monoidal categories in this way.

The commutativity and naturality statements in the third property above imply that the complex $\mathcal{C}\mathbf{1}_n$ is in the (Drinfeld) center of the additive monoidal category $Com(\dot{\mathcal{U}})$ [9, 19, 5]. The collection of chain maps κ_X define an invertible natural transformation of functors $\kappa: - \otimes \mathcal{C} \Rightarrow \mathcal{C} \otimes -$, where $- \otimes \mathcal{C}$ and $\mathcal{C} \otimes -$ are the endofunctors of $Com(\dot{\mathcal{U}})$ given by tensoring on the right, respectively left, with the complex $\mathcal{C}\mathbf{1}_n$ for appropriate n .

The categorification of the Casimir element for quantum \mathfrak{sl}_2 presented here demonstrates the increase in combinatorial complexity that arises when lifting structures to the

categorical level: the Casimir complex only commutes with other complexes up to chain homotopies, which are rather involved. By appealing extensively to the graphical calculus for categorified $\dot{\mathbf{U}}$ and its symmetries, we are able to study the Casimir complex and construct explicit chain maps giving commutativity of the Casimir up to chain homotopy. This paper presents new identities that are used for simplifying 2-morphisms in categorified \mathfrak{sl}_2 . We hope that the calculations in this paper will serve to further illustrate how complex computations can be performed in the graphical calculus for \mathcal{U} .

Acknowledgments: A.B. would like to acknowledge the Swiss National Science Foundation for support via grant PP002-119088. M.K. is grateful to the NSF for partially supporting him via grants DMS-0706924 and DMS-0739392. A.L. was partially supported by the NSF grants DMS-0739392 and DMS-0855713 and would like to thank the MSRI for support in Spring 2010 when this work was almost completed.

2 Casimir element and idempotent form of quantum \mathfrak{sl}_2

2.1 Quantum \mathfrak{sl}_2 and the Casimir element

The quantum group $\mathbf{U}_q(\mathfrak{sl}_2)$ is the associative algebra (with unit) over $\mathbb{Q}(q)$ with generators E, F, K, K^{-1} and relations

$$KK^{-1} = 1 = K^{-1}K, \quad (2.1)$$

$$KE = q^2EK, \quad (2.2)$$

$$KF = q^{-2}FK, \quad (2.3)$$

$$EF - FE = \frac{K - K^{-1}}{q - q^{-1}}. \quad (2.4)$$

For simplicity the algebra $\mathbf{U}_q(\mathfrak{sl}_2)$ is written \mathbf{U} . For more details on quantum groups see [10].

For $a \geq 0$ we put $[a] = \frac{q^a - q^{-a}}{q - q^{-1}}$, $[a]! = [a][a-1] \dots [1]$ and $E^{(a)} = \frac{E^a}{[a]!}$, $F^{(a)} = \frac{F^a}{[a]!}$. We further define the integral form $\mathcal{A}\mathbf{U}$ to be the $\mathbb{Z}[q, q^{-1}]$ -subalgebra of \mathbf{U} generated by

$$\{E^{(a)}, F^{(a)}, K^{\pm 1} \mid a \in \mathbb{Z}_+\}. \quad (2.5)$$

There are several $\mathbb{Z}[q, q^{-1}]$ -(anti)linear (anti)automorphisms that will be used in this paper. Let $\bar{}$ be the \mathbb{Q} -linear involution of $\mathbb{Q}(q)$ which maps q to q^{-1} .

- The $\mathbb{Q}(q)$ -antilinear algebra involution $\underline{\psi}: \mathbf{U} \rightarrow \mathbf{U}$ is given by

$$\underline{\psi}(E) = E, \quad \underline{\psi}(F) = F, \quad \underline{\psi}(K) = K^{-1}, \quad \underline{\psi}(fx) = \bar{f}\underline{\psi}(x) \quad \text{for } f \in \mathbb{Q}(q) \text{ and } x \in \mathbf{U}.$$

- The $\mathbb{Q}(q)$ -linear algebra involution $\underline{\omega}: \mathbf{U} \rightarrow \mathbf{U}$ is given by

$$\begin{aligned}\underline{\omega}(E) &= F, & \underline{\omega}(F) &= E, & \underline{\omega}(K) &= K^{-1}, \\ \underline{\omega}(fx) &= f\underline{\omega}(x), & & \text{for } f \in \mathbb{Q}(q) \text{ and } x \in \mathbf{U}, \\ \underline{\omega}(xy) &= \underline{\omega}(x)\underline{\omega}(y), & & \text{for } x, y \in \mathbf{U}.\end{aligned}$$

- The $\mathbb{Q}(q)$ -linear algebra antiinvolution $\underline{\sigma}: \mathbf{U} \rightarrow \mathbf{U}$ is given by

$$\begin{aligned}\underline{\sigma}(E) &= E, & \underline{\sigma}(F) &= F, & \underline{\sigma}(K) &= K^{-1}, \\ \underline{\sigma}(fx) &= f\underline{\sigma}(x), & & \text{for } f \in \mathbb{Q}(q) \text{ and } x \in \mathbf{U}, \\ \underline{\sigma}(xy) &= \underline{\sigma}(y)\underline{\sigma}(x), & & \text{for } x, y \in \mathbf{U}.\end{aligned}$$

The (anti)linear (anti)involutions pairwise commute and generate the group $G = (\mathbb{Z}_2)^3$ of (anti)linear (anti)automorphisms acting on \mathbf{U} . Throughout the paper we will also use the index two subgroup $G_1 = \{1, \underline{\psi}, \underline{\omega}\underline{\sigma}, \underline{\psi}\underline{\omega}\underline{\sigma}\}$ of G and the coset $G \setminus G_1 = \{\underline{\omega}, \underline{\sigma}, \underline{\psi}\underline{\omega}, \underline{\psi}\underline{\sigma}\}$.

The Casimir element for \mathbf{U} is given by

$$c := EF + \frac{q^{-1}K + qK^{-1}}{(q - q^{-1})^2} = FE + \frac{qK + q^{-1}K^{-1}}{(q - q^{-1})^2}. \quad (2.6)$$

It is easy to verify that $Ec = cE$, $Fc = cF$ and $Kc = cK$. Moreover, c generates the center of \mathbf{U} , and

$$Z(\mathbf{U}) = \mathbb{Q}(q)[c]. \quad (2.7)$$

We will be most interested in the element

$$C := -(q - q^{-1})^2 c = (-q^2 + 2 - q^{-2})EF - q^{-1}K - qK^{-1}, \quad (2.8)$$

$$= (-q^2 + 2 - q^{-2})FE - qK - q^{-1}K^{-1}. \quad (2.9)$$

Of course,

$$Z(\mathbf{U}) = \mathbb{Q}(q)[C]. \quad (2.10)$$

The element C belongs to the integral form ${}_{\mathcal{A}}\mathbf{U}$ of \mathbf{U} , and we call C the Casimir element.

The symmetries in G preserve the Casimir element:

$$\underline{\psi}(C) = \underline{\omega}(C) = \underline{\sigma}(C) = C. \quad (2.11)$$

Notice that the involutions $\underline{\psi}$, $\underline{\omega}\underline{\sigma}$, $\underline{\psi}\underline{\omega}\underline{\sigma}$ in G_1 preserve the form of the Casimir in (2.8) and (2.9), while the involutions $\underline{\omega}$, $\underline{\sigma}$, $\underline{\psi}\underline{\omega}$, $\underline{\psi}\underline{\sigma}$ in $G \setminus G_1$ map one form of the Casimir element in (2.8) and (2.9) to the other.

All of these symmetries preserve the integral form ${}_{\mathcal{A}}\mathbf{U}$ of \mathbf{U} .

2.2 Idempotent rings and their centers

An idempotent ring A is a not necessarily unital associative ring equipped with a family of mutually-orthogonal idempotents 1_i , indexed by elements i of a set I , such that

$$A = \bigoplus_{i,j \in I} 1_i A 1_j. \quad (2.12)$$

The center $Z(A)$ of A is a subspace of $\prod_{i \in I} 1_i A 1_i$ consisting of elements $\prod_{i \in I} z_i$ such that

$$z_i x = x z_j \quad (2.13)$$

for any $i, j \in I$ and $x \in 1_i A 1_j$. $Z(A)$ is a commutative ring isomorphic to the center of the category of idempotent A -modules, i.e. A -modules M such that

$$M = \bigoplus_{i \in I} 1_i M. \quad (2.14)$$

An idempotent ring A has a unit element if and only if the set I is finite, in which case

$$1 = \sum_{i \in I} 1_i. \quad (2.15)$$

For unital A , the center of A defined as above coincides with the usual center of A .

2.3 BLM $\dot{\mathbf{U}}$

The Belinson-Lusztig-MacPherson [2] algebra $\dot{\mathbf{U}}$ is the $\mathbb{Q}(q)$ -algebra obtained by modifying \mathbf{U} by replacing the unit element with a collection of orthogonal idempotents 1_n for $n \in \mathbb{Z}$,

$$1_n 1_m = \delta_{n,m} 1_n, \quad (2.16)$$

indexed by the weight lattice of \mathfrak{sl}_2 , such that

$$K 1_n = 1_n K = q^n 1_n, \quad E 1_n = 1_{n+2} E = 1_{n+2} E 1_n, \quad F 1_n = 1_{n-2} F = 1_{n-2} F 1_n. \quad (2.17)$$

Similarly, the $\mathbb{Z}[q, q^{-1}]$ -subalgebra ${}_{\mathcal{A}}\dot{\mathbf{U}}$ of $\dot{\mathbf{U}}$ is obtained from ${}_{\mathcal{A}}\mathbf{U}$ by replacing the unit element by a collection of orthogonal idempotents (2.16), such that

$$\begin{aligned} K 1_n &= 1_n K = 1_n K 1_n = q^n 1_n, \\ E^{(a)} 1_n &= 1_{n+2a} E^{(a)} = 1_{n+2a} E^{(a)} 1_n, \\ F^{(a)} 1_n &= 1_{n-2a} F^{(a)} = 1_{n-2a} F^{(a)}. \end{aligned} \quad (2.18)$$

The diagram below illustrates the various algebras considered so far

$$\begin{array}{ccc} {}_{\mathcal{A}}\mathbf{U} & \longrightarrow & \mathbf{U} \\ \downarrow \wr & & \downarrow \wr \\ {}_{\mathcal{A}}\dot{\mathbf{U}} & \longrightarrow & \dot{\mathbf{U}}, \end{array} \quad (2.19)$$

where the rightward pointing arrows are the inclusions of subalgebras, and the squiggly arrows denote passing to the idempotent form of the algebra. See [18] and the references therein for more details on the algebra $\dot{\mathbf{U}}$.

There are direct sum decompositions

$$\dot{\mathbf{U}} = \bigoplus_{n,m \in \mathbb{Z}} 1_m \dot{\mathbf{U}} 1_n \quad \mathcal{A}\dot{\mathbf{U}} = \bigoplus_{n,m \in \mathbb{Z}} 1_m (\mathcal{A}\dot{\mathbf{U}}) 1_n$$

with $1_m (\mathcal{A}\dot{\mathbf{U}}) 1_n$ the $\mathbb{Z}[q, q^{-1}]$ -submodule spanned by $1_m E^{(a)} F^{(b)} 1_n$ and $1_m F^{(b)} E^{(a)} 1_n$ for $a, b \in \mathbb{Z}_+$ (these elements are zero unless $m = n + 2a - 2b$).

The algebra $\dot{\mathbf{U}}$ does not have the unit since the infinite sum $\sum_{n \in \mathbb{Z}} 1_n$ does not belong to $\dot{\mathbf{U}}$; instead, the system of idempotents $\{1_n | n \in \mathbb{Z}\}$ serves as a substitute for 1. Lusztig's basis \mathbb{B} of $\dot{\mathbf{U}}$ consists of the following elements of $\dot{\mathbf{U}}$:

- (i) $E^{(a)} F^{(b)} 1_n$ for $a, b \in \mathbb{Z}_+, n \in \mathbb{Z}, n \leq b - a$,
- (ii) $F^{(b)} E^{(a)} 1_n$ for $a, b \in \mathbb{Z}_+, n \in \mathbb{Z}, n \geq b - a$,

where $E^{(a)} F^{(b)} 1_{b-a} = F^{(b)} E^{(a)} 1_{b-a}$.

The (anti)involutions $\underline{\psi}$, $\underline{\omega}$, and $\underline{\sigma}$ all naturally extend to $\dot{\mathbf{U}}$ if we set

$$\underline{\psi}(1_n) = 1_n, \quad \underline{\omega}(1_n) = 1_{-n}, \quad \underline{\sigma}(1_n) = 1_{-n}.$$

Taking direct sums of the induced maps on each summand $1_m \dot{\mathbf{U}} 1_n$ allows these maps to be extended to $\dot{\mathbf{U}}$ and $\mathcal{A}\dot{\mathbf{U}}$. These $\mathbb{Z}[q, q^{-1}]$ -(anti)linear (anti)algebra homomorphisms are recorded below on some elements of $\mathcal{A}\dot{\mathbf{U}}$:

$$\underline{\omega} : q^s 1_m E^{(a)} F^{(b)} 1_n \mapsto q^s 1_{-m} F^{(a)} E^{(b)} 1_{-n}, \quad (2.20)$$

$$\underline{\sigma} : q^s 1_n E^{(a)} F^{(b)} 1_n \mapsto q^s 1_{-n} F^{(b)} E^{(a)} 1_{-m}, \quad (2.21)$$

$$\underline{\psi} : q^s 1_m E^{(a)} F^{(b)} 1_n \mapsto q^{-s} 1_m E^{(a)} F^{(b)} 1_n, \quad (2.22)$$

where $m = n + 2a - 2b$. The group G acts on both $\dot{\mathbf{U}}$ and $\mathcal{A}\dot{\mathbf{U}}$.

There is a natural homomorphism from the center of \mathbf{U} to the center of its idempotent form $\dot{\mathbf{U}}$ that sends $x \in Z(\mathbf{U})$ to $\prod_{n \in \mathbb{Z}} 1_n x 1_n$. It is not hard to check that this homomorphism is, in fact, an isomorphism. Denote by

$$\dot{C} = \prod_{n \in \mathbb{Z}} 1_n C \quad (2.23)$$

the image of C in $\dot{\mathbf{U}}$ under this homomorphism. Let $\dot{C}_{ev} = \prod_{n \in 2\mathbb{Z}} 1_n C$ and $\dot{C}_{od} = \prod_{n \in 2\mathbb{Z}+1} 1_n C$. Then $\dot{C} = \dot{C}_{ev} + \dot{C}_{od}$ and

$$Z(\dot{\mathbf{U}}) \cong \mathbb{Q}(q)[\dot{C}_{ev}] \times \mathbb{Q}(q)[\dot{C}_{od}]. \quad (2.24)$$

We call $C1_n$ components or terms of the Casimir element \dot{C} . They are given by

$$1_n C = C1_n = (-q^2 + 2 - q^{-2})EF1_n - (q^{n-1} + q^{1-n})1_n, \quad (2.25)$$

$$= (-q^2 + 2 - q^{-2})FE1_n - (q^{n+1} + q^{-1-n})1_n. \quad (2.26)$$

Components are preserved by the symmetries in $G_1 = \{1, \underline{\psi}, \underline{\omega\sigma}, \underline{\psi\sigma\omega}\}$:

$$g(C1_n) = C1_n \quad (2.27)$$

for any $g \in G_1$. They are interchanged by the elements in the coset $G \setminus G_1 = \{\underline{\sigma}, \underline{\omega}, \underline{\psi\sigma}, \underline{\psi\omega}\}$:

$$g'(C1_n) = C1_{-n} \quad (2.28)$$

for any $g' \in G \setminus G_1$. In addition to sending 1_n to 1_{-n} , the involutions in $G \setminus G_1$ map one form of the Casimir in (2.25) and (2.26) to the other form.

3 Brief review of $\mathfrak{sl}(2)$ -calculus

3.1 The 2-category \mathcal{U}

3.1.1 Definition of \mathcal{U}

Fix a field \mathbb{k} . Here we recall the definition of the \mathbb{k} -linear 2-category $\mathcal{U} = \mathcal{U}(\mathfrak{sl}_2)$ introduced in [15], see also [14, 12].

Definition 3.1. The 2-category \mathcal{U} is the additive 2-category consisting of

- objects: n for $n \in \mathbb{Z}$.

The hom $\mathcal{U}(n, n')$ between two objects n, n' is an additive \mathbb{k} -linear category:

- objects of $\mathcal{U}(n, n')$: for a signed sequence $\underline{\varepsilon} = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_m)$, where $\varepsilon_1, \dots, \varepsilon_m \in \{+, -\}$, define

$$\mathcal{E}_{\underline{\varepsilon}} := \mathcal{E}_{\varepsilon_1} \mathcal{E}_{\varepsilon_2} \dots \mathcal{E}_{\varepsilon_m}$$


where $\mathcal{E}_+ := \mathcal{E}$ and $\mathcal{E}_- := \mathcal{F}$. An object of $\mathcal{U}(n, n')$, called a 1-morphism in \mathcal{U} , is a formal finite direct sum of 1-morphisms

$$\mathcal{E}_{\underline{\varepsilon}} \mathbf{1}_n \{t\} = \mathbf{1}_{n'} \mathcal{E}_{\underline{\varepsilon}} \mathbf{1}_n \{t\}$$


for any $t \in \mathbb{Z}$ and signed sequence $\underline{\varepsilon}$ such that $n' = n + \sum_{j=1}^m \varepsilon_j 2$.

- morphisms of $\mathcal{U}(n, n')$: given objects $\mathcal{E}_{\underline{\varepsilon}} \mathbf{1}_n \{t\}, \mathcal{E}_{\underline{\varepsilon}'} \mathbf{1}_n \{t'\} \in \mathcal{U}(n, n')$, the hom sets $\mathcal{U}(\mathcal{E}_{\underline{\varepsilon}} \mathbf{1}_n \{t\}, \mathcal{E}_{\underline{\varepsilon}'} \mathbf{1}_n \{t'\})$ of $\mathcal{U}(n, n')$ are \mathbb{k} -vector spaces given by linear combinations of diagrams with degree $t - t'$, modulo certain relations, built from composites of:

- i) Degree zero identity 2-morphisms 1_x for each 1-morphism x in \mathcal{U} ; the identity 2-morphisms $1_{\mathcal{E}1_n\{t\}}$ and $1_{\mathcal{F}1_n\{t\}}$ are represented graphically by











and more generally, for a signed sequence $\underline{\varepsilon} = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_m)$, the identity $1_{\mathcal{E}_{\underline{\varepsilon}}1_n\{t\}}$ 2-morphism is represented as



where the strand labelled $\mathcal{E}_{\varepsilon_\alpha}$ is oriented up if $\varepsilon_\alpha = +$ and oriented down if $\varepsilon_\alpha = -$. We will often omit labels from the strands since the labels can be deduced from the orientation of a strand.

- ii) For each $n \in X$ the 2-morphisms

2-morphism:				
Degree:	2	2	-2	-2

2-morphism:				
Degree:	1 + n	1 - n	1 + n	1 - n

such that the following identities hold:

- cups and caps are biadjointness morphisms up to grading shifts:

$$\begin{array}{ccc}
 \begin{array}{c} n+2 \\ \uparrow \\ \downarrow \\ n \end{array} & = & \begin{array}{c} n+2 \\ \uparrow \\ n \end{array} \\
 \begin{array}{c} n \\ \uparrow \\ \downarrow \\ n+2 \end{array} & = & \begin{array}{c} n \\ \uparrow \\ n+2 \end{array}
 \end{array} \tag{3.1}$$

where the second equality in (3.7) and (3.8) follow from (3.6). We also write

$$\begin{array}{c} | \\ | \\ \bullet \\ | \\ \alpha \end{array} \begin{array}{c} n \\ \\ \\ \end{array} := \left(\begin{array}{c} | \\ | \\ \bullet \\ | \\ \alpha \end{array} \begin{array}{c} n \\ \\ \\ \end{array} \right)^\alpha$$

to denote the α -fold vertical composite of a dot with itself.

- All dotted bubbles of negative degree are zero. That is,

$$\begin{array}{c} n \\ \curvearrowright \\ \bullet \\ \alpha \end{array} = 0 \quad \text{if } \alpha < n - 1, \quad \begin{array}{c} n \\ \curvearrowleft \\ \bullet \\ \alpha \end{array} = 0 \quad \text{if } \alpha < -n - 1 \quad (3.9)$$

for all $\alpha \in \mathbb{Z}_+$. A dotted bubble of degree zero equals 1:

$$\begin{array}{c} n \\ \curvearrowright \\ \bullet \\ n-1 \end{array} = 1 \quad \text{for } n \geq 1, \quad \begin{array}{c} n \\ \curvearrowleft \\ \bullet \\ -n-1 \end{array} = 1 \quad \text{for } n \leq -1.$$

It is often convenient to express dotted bubbles using a notation introduced in [13] that emphasizes the degree:

$$\begin{array}{c} n \\ \curvearrowright \\ \bullet \\ \spadesuit + \alpha \end{array} := \begin{array}{c} n \\ \curvearrowright \\ \bullet \\ (n-1) + \alpha \end{array} \quad \begin{array}{c} n \\ \curvearrowleft \\ \bullet \\ \spadesuit + \alpha \end{array} := \begin{array}{c} n \\ \curvearrowleft \\ \bullet \\ (-n-1) + \alpha \end{array}$$

so that

$$\deg \left(\begin{array}{c} n \\ \curvearrowright \\ \bullet \\ \spadesuit + \alpha \end{array} \right) = 2\alpha \quad \deg \left(\begin{array}{c} n \\ \curvearrowleft \\ \bullet \\ \spadesuit + \alpha \end{array} \right) = 2\alpha.$$

The value of \spadesuit depends on the orientation, $\spadesuit = n - 1$ for clockwise oriented bubbles and $\spadesuit = -n - 1$ for counter-clockwise oriented bubbles. Notice that for some values of n it is possible that $\spadesuit + \alpha$ is a negative number even though $\alpha \geq 0$. While vertically composing a generator with itself a negative number of times does not make sense, having these symbols around greatly simplifies the calculus. For each $\spadesuit + \alpha < 0$, where

$$\deg \left(\begin{array}{c} n \\ \curvearrowright \\ \bullet \\ \spadesuit + \alpha \end{array} \right) \geq 0 \quad \deg \left(\begin{array}{c} n \\ \curvearrowleft \\ \bullet \\ \spadesuit + \alpha \end{array} \right) \geq 0,$$

we introduce formal symbols, called *fake bubbles*, inductively defined by the equation

$$\left(\begin{array}{c} n \\ \curvearrowright \\ \bullet \\ \spadesuit + 0 \end{array} + \begin{array}{c} n \\ \curvearrowright \\ \bullet \\ \spadesuit + 1 \end{array} t + \cdots + \begin{array}{c} n \\ \curvearrowright \\ \bullet \\ \spadesuit + \alpha \end{array} t^\alpha + \cdots \right) \left(\begin{array}{c} n \\ \curvearrowleft \\ \bullet \\ \spadesuit + 0 \end{array} + \cdots + \begin{array}{c} n \\ \curvearrowleft \\ \bullet \\ \spadesuit + \alpha \end{array} t^\alpha + \cdots \right) = 1. \quad (3.10)$$

and the additional condition

$$\begin{array}{c} n \\ \circlearrowleft \\ \blacktriangle +0 \end{array} = \begin{array}{c} n \\ \circlearrowright \\ \blacktriangle +0 \end{array} = 1.$$

Equation (3.10) is called the infinite Grassmannian relation. It remains valid even in high degree when most of the bubbles involved are not fake bubbles. See [15] for more details.

- For the following relations we employ the convention that all summations are increasing, so that $\sum_{f=0}^{\alpha}$ is zero if $\alpha < 0$.

$$\begin{array}{c} n \\ \text{X} \\ n \end{array} = - \sum_{f_1+f_2=-n} \begin{array}{c} f_1 \\ | \\ \bullet \end{array} \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangle +f_2 \end{array} \quad \begin{array}{c} n \\ \text{X} \\ n \end{array} = \sum_{g_1+g_2=n} \begin{array}{c} n \\ \circlearrowright \\ \blacktriangle +g_2 \end{array} \begin{array}{c} g_1 \\ | \\ \bullet \end{array} \quad (3.11)$$

$$\begin{array}{c} n \\ | \\ n \end{array} \begin{array}{c} | \\ n \end{array} = - \begin{array}{c} \text{X} \\ n \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangle +f_2 \\ \circlearrowleft \\ f_3 \end{array} \begin{array}{c} f_1 \\ | \\ \bullet \end{array} \\ \begin{array}{c} n \\ | \\ n \end{array} \begin{array}{c} | \\ n \end{array} = - \begin{array}{c} \text{X} \\ n \end{array} + \sum_{\substack{g_1+g_2+g_3 \\ =-n-1}} \begin{array}{c} n \\ \circlearrowright \\ \blacktriangle +g_2 \\ \circlearrowright \\ g_3 \end{array} \begin{array}{c} g_1 \\ | \\ \bullet \end{array} \quad (3.12)$$

for all $n \in \mathbb{Z}$. In equations (3.11) and (3.12) whenever the summations are nonzero they utilize fake bubbles.

- the additive composition functor $\mathcal{U}(n, n') \times \mathcal{U}(n', n'') \rightarrow \mathcal{U}(n, n'')$ is given on 1-morphisms of \mathcal{U} by

$$\mathcal{E}_{\underline{\varepsilon}'} \mathbf{1}_{n'} \{t'\} \times \mathcal{E}_{\underline{\varepsilon}} \mathbf{1}_n \{t\} \mapsto \mathcal{E}_{\underline{\varepsilon}' \underline{\varepsilon}} \mathbf{1}_n \{t+t'\} \quad (3.13)$$

for $n' = n + \sum_{j=1}^m \varepsilon_j 2$, and on 2-morphisms of \mathcal{U} by juxtaposition of diagrams

$$\left(\begin{array}{c} n'' \\ \text{X} \\ n' \end{array} \right) \times \left(\begin{array}{c} n' \\ \text{X} \\ n \end{array} \right) \mapsto \begin{array}{c} n'' \\ \text{X} \\ n \end{array}$$

3.1.2 Relations in \mathcal{U}

In this section we collect some relations that follow from the definition of \mathcal{U} . These relations were proven in [15].

$$\begin{array}{c} \alpha \\ \bullet \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \bullet \\ n \end{array} - \begin{array}{c} \diagup \quad \diagdown \\ \bullet \\ \alpha \\ \bullet \\ \diagdown \quad \diagup \\ n \end{array} = \begin{array}{c} \alpha \\ \bullet \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ n \end{array} - \begin{array}{c} \diagup \quad \diagdown \\ \bullet \\ \alpha \\ \bullet \\ \diagdown \quad \diagup \\ n \end{array} = \sum_{f_1+f_2=\alpha-1} \begin{array}{c} f_1 \\ \uparrow \\ \bullet \\ \uparrow \\ f_2 \\ n \end{array} \quad (3.14)$$

For all $n \in \mathbb{Z}$ the following equation holds

$$\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ n \end{array} - \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ n \end{array} = \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n}} \begin{array}{c} \bullet \\ \curvearrowright \\ f_1 \\ \bullet \\ \curvearrowleft \\ f_3 \\ \bullet \\ \curvearrowright \\ f_2 \\ \bullet \\ \curvearrowleft \\ f_4 \\ \bullet \\ n \end{array} + \sum_{\substack{g_1+g_2+g_3+g_4 \\ =-n-2}} \begin{array}{c} g_2 \\ \bullet \\ \curvearrowright \\ g_1 \\ \bullet \\ \curvearrowleft \\ g_3 \\ \bullet \\ \curvearrowright \\ g_4 \\ \bullet \\ n \end{array} \quad (3.15)$$

where the first sum is over all $f_1, f_2, f_3, f_4 \geq 0$ with $f_1 + f_2 + f_3 + f_4 = n$ and the second sum is over all $g_1, g_2, g_3, g_4 \geq 0$ with $g_1 + g_2 + g_3 + g_4 = -n - 2$. Recall that all summations in this paper are increasing, so that the first summation is zero if $n < 0$ and the second is zero when $-2 < n$. By rotating this equation and shifting n we also have

$$\begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ n \end{array} - \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ n \end{array} = \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n-2}} \begin{array}{c} f_2 \\ \bullet \\ \curvearrowright \\ f_1 \\ \bullet \\ \curvearrowleft \\ f_3 \\ \bullet \\ \curvearrowright \\ f_4 \\ \bullet \\ n \end{array} + \sum_{\substack{g_1+g_2+g_3+g_4 \\ =-n}} \begin{array}{c} g_1 \\ \bullet \\ \curvearrowright \\ g_2 \\ \bullet \\ \curvearrowleft \\ g_3 \\ \bullet \\ \curvearrowright \\ g_4 \\ \bullet \\ n \end{array} \quad (3.16)$$

Dotted curl relations:

$$\begin{array}{c} \bullet \\ \curvearrowright \\ \bullet \\ \bullet \\ x \\ \bullet \\ \curvearrowleft \\ \bullet \\ n \end{array} = - \sum_{f_1+f_2=x-n} \begin{array}{c} f_1 \\ \bullet \\ \uparrow \\ \bullet \\ \uparrow \\ f_2 \\ \bullet \\ n \end{array} \begin{array}{c} \bullet \\ \curvearrowright \\ \bullet \\ \bullet \\ \curvearrowleft \\ \bullet \\ n \end{array} \quad \begin{array}{c} \bullet \\ \curvearrowleft \\ \bullet \\ \bullet \\ x \\ \bullet \\ \curvearrowright \\ \bullet \\ n \end{array} = \sum_{g_1+g_2=x+n} \begin{array}{c} g_1 \\ \bullet \\ \uparrow \\ \bullet \\ \uparrow \\ g_2 \\ \bullet \\ n \end{array} \begin{array}{c} \bullet \\ \curvearrowright \\ \bullet \\ \bullet \\ \curvearrowleft \\ \bullet \\ n \end{array} \quad (3.17)$$

One can also show the relations:

$$\begin{array}{c} n \\ \uparrow \\ \bullet \\ x \\ \uparrow \\ \bullet \\ y \\ n \end{array} = - \begin{array}{c} \bullet \\ \curvearrowright \\ \bullet \\ \bullet \\ y \\ \bullet \\ \curvearrowleft \\ \bullet \\ x \\ n \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =x+y+n-1}} \begin{array}{c} n \\ \curvearrowright \\ \bullet \\ f_1 \\ \bullet \\ \curvearrowleft \\ f_2 \\ \bullet \\ \curvearrowright \\ f_3 \\ \bullet \\ n \end{array} \\ \begin{array}{c} n \\ \uparrow \\ \bullet \\ x \\ \uparrow \\ \bullet \\ y \\ n \end{array} = - \begin{array}{c} \bullet \\ \curvearrowleft \\ \bullet \\ \bullet \\ y \\ \bullet \\ \curvearrowright \\ \bullet \\ x \\ n \end{array} + \sum_{\substack{g_1+g_2+g_3 \\ =x+y+n-1}} \begin{array}{c} n \\ \curvearrowright \\ \bullet \\ g_1 \\ \bullet \\ \curvearrowleft \\ g_2 \\ \bullet \\ \curvearrowright \\ g_3 \\ \bullet \\ n \end{array} \quad (3.18)$$

Bubble slide equations:

$$\begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} \circlearrowleft \\ \blacktriangleright + j \end{array} = \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangleright + (j-2) \end{array} \begin{array}{c} \bullet \\ \downarrow \\ \bullet \\ \downarrow \end{array} - 2 \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangleright + (j-1) \end{array} \begin{array}{c} \bullet \\ \downarrow \end{array} + \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangleright + j \end{array} \begin{array}{c} \downarrow \end{array} \quad (3.19)$$

$$\begin{array}{c} \circlearrowleft \\ \blacktriangleright + j \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} = \begin{array}{c} \bullet \\ \downarrow \\ \bullet \\ \downarrow \end{array} \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangleright + (j-2) \end{array} - 2 \begin{array}{c} \bullet \\ \downarrow \end{array} \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangleright + (j-1) \end{array} + \begin{array}{c} \downarrow \end{array} \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangleright + j \end{array} \quad (3.20)$$

$$\begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangleright + j \end{array} = \sum_{f=0}^j (j+1-f) \begin{array}{c} n+2 \\ \circlearrowleft \\ \blacktriangleright + f \end{array} \begin{array}{c} \bullet \\ \downarrow \\ \bullet \\ \downarrow \end{array} \begin{array}{c} j-f \\ \downarrow \end{array} \quad (3.21)$$

$$\begin{array}{c} \circlearrowleft \\ \blacktriangleright + j \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} = \sum_{f=0}^j (j+1-f) \begin{array}{c} \bullet \\ \downarrow \\ \bullet \\ \downarrow \end{array} \begin{array}{c} j-f \\ \downarrow \end{array} \begin{array}{c} n \\ \circlearrowleft \\ \blacktriangleright + f \end{array} \quad (3.22)$$

Below we collect a few additional identities that have not appeared in the literature previously.

The following relation together with its image under the various symmetries of the 2-category \mathcal{U} will be used extensively in the paper.

Proposition 3.2.

$$\begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} - \begin{array}{c} \bullet \\ \downarrow \\ \bullet \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} + \begin{array}{c} \bullet \\ \downarrow \\ \bullet \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} + \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} = 0. \quad (3.23)$$

The proof utilizes the nilHecke relations to slide dots as well as (3.16).

Proposition 3.3 (Step functions).

$$\begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} = \begin{cases} - \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} & \text{if } n \leq 0 \\ 0 & \text{otherwise.} \end{cases} \quad (3.24)$$

$$\begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} = \begin{cases} - \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n \\ \downarrow \\ \downarrow \\ \downarrow \end{array} & \text{if } n \geq 0 \\ 0 & \text{otherwise.} \end{cases} \quad (3.25)$$

Proof. This proposition follows immediately from the Dotted Curl Relations (3.17). \square

Proposition 3.4.

$$\text{Diagram with } n \text{ crossings} = \begin{cases} \text{Diagram with } 0 \text{ crossings and } \spadesuit_{+1} & \text{if } n = 0, \\ 0 & \text{otherwise.} \end{cases} \quad (3.26)$$

Proof. This proposition follows from the Curl relations in \mathcal{U} together with the positivity of bubbles axiom. \square

3.2 The 2-categories $\dot{\mathcal{U}}$, $Kom(\dot{\mathcal{U}})$, and $Com(\dot{\mathcal{U}})$

3.2.1 Additive categories, homotopy categories, and Karoubi envelopes

For an additive category \mathcal{M} we write $Kom(\mathcal{M})$ for the category of bounded complexes in \mathcal{M} . An object (X, d) of $Kom(\mathcal{M})$ is a collection of objects X^i of \mathcal{M} together with maps d_i

$$\cdots \xrightarrow{d} X^{i-1} \xrightarrow{d_{i-1}} X^i \xrightarrow{d_i} X^{i+1} \xrightarrow{d_{i+1}} \cdots \quad (3.27)$$

such that $d_{i+1}d_i = 0$ and only finitely many objects are nonzero. A morphism $f: (X, d) \rightarrow (Y, d)$ in $Kom(\mathcal{M})$ is a collection of morphisms $f_i: X^i \rightarrow Y^i$ such that

$$\begin{array}{ccccccc} \cdots & \xrightarrow{d} & X^{i-1} & \xrightarrow{d_{i-1}} & X^i & \xrightarrow{d_i} & X^{i+1} \xrightarrow{d_{i+1}} \cdots \\ & & \downarrow f_{i-1} & & \downarrow f_i & & \downarrow f_{i+1} \\ \cdots & \xrightarrow{d} & Y^{i-1} & \xrightarrow{d_{i-1}} & Y^i & \xrightarrow{d_i} & Y^{i+1} \xrightarrow{d_{i+1}} \cdots \end{array} \quad (3.28)$$

commutes.

Given a pair of morphisms $f, g: (X, d) \rightarrow (Y, d)$ in $Kom(\mathcal{M})$, we say that f is *homotopic* to g if there exists morphisms $h^i: X^i \rightarrow Y^{i-1}$ such that $f_i - g_i = h^{i+1}d_i + d_{i-1}h^i$ for all i . A morphism of complexes is said to be *null-homotopic* if it is homotopic to the zero map.

Definition 3.5. The homotopy category $Com(\mathcal{M})$ has the same objects as $Kom(\mathcal{M})$, and morphisms are morphisms in $Kom(\mathcal{M})$ modulo null-homotopic morphisms.

The Karoubi envelope $Kar(\mathcal{M})$ of a category \mathcal{M} is an enlargement of \mathcal{M} in which all idempotents split. An idempotent $e: b \rightarrow b$ in a category \mathcal{M} is said to split if there exist morphisms

$$b \xrightarrow{g} b' \xrightarrow{h} b$$

such that $e = hg$ and $gh = \text{Id}_{b'}$. More precisely, the Karoubi envelope $Kar(\mathcal{M})$ is a category whose objects are pairs (b, e) where $e: b \rightarrow b$ is an idempotent of \mathcal{M} and whose morphisms are triples of the form

$$(e, f, e'): (b, e) \rightarrow (b', e')$$

where $f: b \rightarrow b'$ in \mathcal{M} making the diagram

$$\begin{array}{ccc} b & \xrightarrow{f} & b' \\ e \downarrow & \searrow f & \downarrow e' \\ b & \xrightarrow{f} & b' \end{array} \quad (3.29)$$

commute. Thus, f must satisfy $f = e'fe$, which is equivalent to $f = e'f = fe$. Composition is induced from the composition in \mathcal{M} , and the identity morphism is $(e, e, e): (b, e) \rightarrow (b, e)$. When \mathcal{M} is an additive category, the splitting of idempotents allows us to write $(b, e) \in Kar(\mathcal{M})$ as $\text{im } e$, and we have $b \cong \text{im } e \oplus \text{im } (\text{Id}_b - e)$.

The identity map $\text{Id}_b: b \rightarrow b$ is an idempotent, and this gives rise to a fully faithful functor $\mathcal{M} \rightarrow Kar(\mathcal{M})$. In $Kar(\mathcal{M})$ all idempotents of \mathcal{M} split and this functor is universal with respect to functors which split idempotents in \mathcal{M} . When \mathcal{M} is additive the inclusion $\mathcal{M} \rightarrow Kar(\mathcal{M})$ is an additive functor (see [15, Section 9] and references therein).

Proposition 3.6. For any additive category \mathcal{M} there exists a canonical equivalence

$$Kom(Kar(\mathcal{M})) \cong Kar(Kom(\mathcal{M})). \quad (3.30)$$

Proof. Define the functor

$$\rho_{\mathcal{M}}: Kom(Kar(\mathcal{M})) \rightarrow Kar(Kom(\mathcal{M})) \quad (3.31)$$

as follows. An object of $Kom(Kar(\mathcal{M}))$ has the form

$$(X, e) = \cdots \xrightarrow{d_{i-1}} (X^i, e_i) \xrightarrow{d_i} (X^{i+1}, e_{i+1}) \xrightarrow{d_{i+1}} \cdots \quad (3.32)$$

where $e_i^2 = e_i$, $d_{i+1}d_i = 0$, and $d_i = e_{i+1}d_i e_i$. Here $e_i: X^i \rightarrow X^i$ is an idempotent and $d_i: X^i \rightarrow X^{i+1}$. The functor $\rho_{\mathcal{M}}$ takes this object to the pair in $Kar(Kom(\mathcal{M}))$ consisting of the complex

$$X = \cdots \xrightarrow{d_{i-1}} X^i \xrightarrow{d_i} X^{i+1} \xrightarrow{d_{i+1}} \cdots \quad (3.33)$$

and the idempotent chain map $(\dots, e_i, e_{i+1}, \dots)$. A morphism $f: (X, e) \rightarrow (X', e')$ in $Kom(Kar(\mathcal{M}))$ is a collection of maps $f_i: X^i \rightarrow (X')^i$ such that the squares

$$\begin{array}{ccc} X^i & \xrightarrow{d_i} & X^{i+1} \\ f_i \downarrow & & \downarrow f_{i+1} \\ (X')^i & \xrightarrow{d'_i} & (X')^{i+1} \end{array} \quad (3.34)$$

commute, and $f_i = e'_i f_i e_i$. The functor $\rho_{\mathcal{M}}$ takes the morphism f to the “same” morphism $\{f_i\}$ of complexes equipped with idempotents $(\dots, e_i, e_{i+1}, \dots)$ and $(\dots, e'_i, e'_{i+1}, \dots)$.

It is easy to see that $\rho_{\mathcal{M}}$ is fully-faithful. To show $\rho_{\mathcal{M}}$ is essentially surjective note that any object Y of $Kar(Kom(\mathcal{M}))$ is isomorphic to $\rho_{\mathcal{M}}(\tilde{Y})$ for some object \tilde{Y} of $Kom(Kar(\mathcal{M}))$. The object Y consists of a complex

$$Y = \dots \xrightarrow{d_{i-1}} Y^i \xrightarrow{d_i} Y^{i+1} \xrightarrow{d_{i+1}} \dots \quad (3.35)$$

in $Kom(\mathcal{M})$ together with idempotents $e_i: Y^i \rightarrow Y^i$ such that

$$e_{i+1}d_i = d_ie_i. \quad (3.36)$$

Let \tilde{Y} be the object

$$\dots \xrightarrow{d_{i-1}e_{i-1}} (Y^i, e_i) \xrightarrow{d_ie_i} (Y^{i+1}, e_{i+1}) \xrightarrow{d_{i+1}e_{i+1}} \dots \quad (3.37)$$

in $Kom(Kar(\mathcal{M}))$. We can define morphisms $\varphi_1: Y \rightarrow \rho_{\mathcal{M}}(\tilde{Y})$ and $\varphi_2: \rho_{\mathcal{M}}(\tilde{Y}) \rightarrow Y$ in $Kar(Kom(\mathcal{M}))$ that are both given on Y^i as multiplication by e_i . Then $\varphi_2\varphi_1 = \text{Id}_Y$ and $\varphi_1\varphi_2 = \text{Id}_{\rho_{\mathcal{M}}(\tilde{Y})}$, showing that Y and $\rho_{\mathcal{M}}(\tilde{Y})$ are isomorphic. Together with fully-faithfulness of $\rho_{\mathcal{M}}$ this completes the proof that $\rho_{\mathcal{M}}$ is an equivalence of categories. \square

Proposition 3.7. For any additive \mathbb{k} -linear category \mathcal{M} with finite dimensional hom spaces there exists a canonical equivalence

$$Com(Kar(\mathcal{M})) \cong Kar(Com(\mathcal{M})). \quad (3.38)$$

Proof. The functor $\rho_{\mathcal{M}}$ descends to a functor

$$\rho_{\mathcal{M}}^c: Com(Kar(\mathcal{M})) \rightarrow Kar(Com(\mathcal{M})). \quad (3.39)$$

Given an object X of $Com(Kar(\mathcal{M}))$ we can view it as an object of $Kom(Kar(\mathcal{M}))$. By the idempotent-lifting property for finite-dimensional algebras [4, Chapter 1] an idempotent of $Com(Kar(\mathcal{M}))$ lifts to an idempotent in $Kom(Kar(\mathcal{M}))$, and the latter category is idempotent-complete by the previous proposition. Therefore, $Com(Kar(\mathcal{M}))$ is idempotent-complete as well, allowing us to define a functor

$$\tilde{\rho}_{\mathcal{M}}: Kar(Com(\mathcal{M})) \rightarrow Kar(Com(Kar(\mathcal{M}))) \simeq Com(Kar(\mathcal{M})) \quad (3.40)$$

such that $\rho_{\mathcal{M}}^c\tilde{\rho}_{\mathcal{M}} \cong \text{Id}_{Kar(Com(\mathcal{M}))}$, $\tilde{\rho}_{\mathcal{M}}\rho_{\mathcal{M}}^c \cong \text{Id}_{Com(Kar(\mathcal{M}))}$, showing that $\rho_{\mathcal{M}}^c$ is an equivalence. \square

Alternatively, the result follows from [1, Corollary 2.12].

3.2.2 Karoubian envelope of \mathcal{U}

Definition 3.8. Define the additive 2-category $\dot{\mathcal{U}}$ to have the same objects as \mathcal{U} and hom additive categories given by $\dot{\mathcal{U}}(n, m) = Kar(\mathcal{U}(n, m))$. The fully-faithful additive functors $\mathcal{U}(n, m) \rightarrow \dot{\mathcal{U}}(n, m)$ combine to form an additive 2-functor $\mathcal{U} \rightarrow \dot{\mathcal{U}}$ universal with respect to splitting idempotents in the hom categories $\dot{\mathcal{U}}(n, m)$. The composition functor $\dot{\mathcal{U}}(n, n') \times \dot{\mathcal{U}}(n', n'') \rightarrow \dot{\mathcal{U}}(n, n'')$ is induced by the universal property of the Karoubi envelope from the composition functor for \mathcal{U} . The 2-category $\dot{\mathcal{U}}$ has graded 2-homs given by

$$HOM_{\dot{\mathcal{U}}}(x, y) := \bigoplus_{t \in \mathbb{Z}} Hom_{\dot{\mathcal{U}}}(x\{t\}, y). \quad (3.41)$$

Theorem 3.9 (Theorem 9.1.3 [15]). There is an isomorphism

$$\gamma: \dot{\mathcal{U}}_{\mathcal{A}} \rightarrow K_0(\dot{\mathcal{U}}), \quad (3.42)$$

where $K_0(\dot{\mathcal{U}})$ is the split Grothendieck ring of $\dot{\mathcal{U}}$.

In [13, Corollary 5.14] this result is proven when the ground field \mathbb{k} is replaced by the commutative ring \mathbb{Z} .

3.2.3 Karoubian envelopes of $Kom(\mathcal{U})$ and $Com(\mathcal{U})$

Definition 3.10. Define $Kom(\mathcal{U})$ to be the additive 2-category with objects $n \in \mathbb{Z}$ and additive hom categories $Kom(\mathcal{U})(n, m) := Kom(\mathcal{U}(n, m))$. The additive composition functor $Kom(\mathcal{U}(n', n'')) \times Kom(\mathcal{U}(n, n')) \rightarrow Kom(\mathcal{U}(n, n''))$ is given by the tensor product of complexes using the additive composition functor on \mathcal{U} to tensor 1-morphisms via composition.

Definition 3.11. Define $Com(\mathcal{U})$ to be the additive 2-category with the same objects and 1-morphisms as $Kom(\mathcal{U})$ and 2-morphisms given by identifying homotopy equivalent 2-morphisms in $Kom(\mathcal{U})$.

Recall that $\dot{\mathcal{U}} = Kar(\mathcal{U})$. By Propositions 3.6 and 3.7 there are equivalences

$$Kar(Kom(\mathcal{U})) \cong Kom(\dot{\mathcal{U}}), \quad Kar(Com(\mathcal{U})) \cong Com(\dot{\mathcal{U}}). \quad (3.43)$$

The 2-categories we consider fit into the following table where the horizontal arrows denote passage to the Karoubian envelope and vertical arrows stand for passage to complexes and modding out by null-homotopic maps.

$$\begin{array}{ccc}
 \mathcal{U} & \longrightarrow & \dot{\mathcal{U}} = Kar(\mathcal{U}) \\
 \downarrow & & \downarrow \\
 Kom(\mathcal{U}) & \longrightarrow & Kom(\dot{\mathcal{U}}) \cong Kar(Kom(\mathcal{U})) \\
 \downarrow & & \downarrow \\
 Com(\mathcal{U}) & \longrightarrow & Com(\dot{\mathcal{U}}) \cong Kar(Com(\mathcal{U}))
 \end{array} \quad (3.44)$$

3.3 Symmetry 2-functors

A covariant/contravariant functor $\alpha: \mathcal{M} \rightarrow \mathcal{M}'$ extends canonically to a functor

$$Kar(\alpha): Kar(\mathcal{M}) \rightarrow Kar(\mathcal{M}').$$

An additive covariant/contravariant functor $\alpha: \mathcal{M} \rightarrow \mathcal{M}'$ between additive categories extends canonically to an additive functor

$$Kom(\alpha): Kom(\mathcal{M}) \rightarrow Kom(\mathcal{M}')$$

and an exact functor $Com(\alpha): Com(\mathcal{M}) \rightarrow Com(\mathcal{M}')$ between triangulated categories.

Given an exact endofunctor $\alpha: \mathcal{M} \rightarrow \mathcal{M}$ these extensions respect the equivalence $\rho_{\mathcal{M}}$ in (3.31), in the sense that the diagrams

$$\begin{array}{ccc} Kom(Kar(\mathcal{M})) & \xrightarrow{\rho_{\mathcal{M}}} & Kar(Kom(\mathcal{M})) \\ \downarrow Kom(Kar(\alpha)) & & \downarrow Kar(Kom(\alpha)) \\ Kom(Kar(\mathcal{M})) & \xrightarrow{\rho_{\mathcal{M}}} & Kar(Kom(\mathcal{M})) \end{array} \quad (3.45)$$

$$\begin{array}{ccc} Com(Kar(\mathcal{M})) & \xrightarrow{\rho_{\mathcal{M}}} & Kar(Com(\mathcal{M})) \\ \downarrow Com(Kar(\alpha)) & & \downarrow Kar(Com(\alpha)) \\ Com(Kar(\mathcal{M})) & \xrightarrow{\rho_{\mathcal{M}}} & Kar(Com(\mathcal{M})) \end{array} \quad (3.46)$$

commute.

In this section we recall several 2-functor involutions ω, ψ, σ on the 2-category \mathcal{U} defined in [15] and extend them to 2-functors on all the 2-categories in (3.44). We use the same notation for these extended 2-functors.

Denote by \mathcal{U}^{op} the 2-category with the same objects as \mathcal{U} but the 1-morphisms reversed. The direction of the 2-morphisms remain fixed. The 2-category \mathcal{U}^{co} has the same objects and 1-morphism as \mathcal{U} , but the directions of the 2-morphisms is reversed. That is, $\mathcal{U}^{\text{co}}(x, y) = \mathcal{U}(y, x)$ for 1-morphisms x and y . Finally, $\mathcal{U}^{\text{coop}}$ denotes the 2-category with the same objects as \mathcal{U} , but the directions of the 1-morphisms and 2-morphisms have been reversed.

Using the symmetries of the diagrammatic relations imposed on \mathcal{U} 2-functors were defined in [15] that categorify various $\mathbb{Z}[q, q^{-1}]$ -(anti)linear (anti)automorphisms of the algebra $\dot{\mathbf{U}}$. The various forms of contravariant behaviour for 2-functors on \mathcal{U} translate into properties of the corresponding homomorphism in $\dot{\mathbf{U}}$ as the following table summarizes:

2-functors	Algebra maps
$\mathcal{U} \rightarrow \mathcal{U}$	$\mathbb{Z}[q, q^{-1}]$ -linear homomorphisms
$\mathcal{U} \rightarrow \mathcal{U}^{\text{op}}$	$\mathbb{Z}[q, q^{-1}]$ -linear antihomomorphisms
$\mathcal{U} \rightarrow \mathcal{U}^{\text{co}}$	$\mathbb{Z}[q, q^{-1}]$ -antilinear homomorphisms
$\mathcal{U} \rightarrow \mathcal{U}^{\text{coop}}$	$\mathbb{Z}[q, q^{-1}]$ -antilinear antihomomorphisms

Rescale, invert the orientation, and send $n \mapsto -n$: Consider the operation on the diagrammatic calculus that rescales the crossing $\begin{array}{c} \nearrow \\ \searrow \end{array} \mapsto - \begin{array}{c} \searrow \\ \nearrow \end{array}$, inverts the orientation of each strand and sends $n \mapsto -n$:

This gives a strict invertible 2-functor $\omega: \mathcal{U} \rightarrow \mathcal{U}$

$$\begin{aligned} \omega: \mathcal{U} &\rightarrow \mathcal{U} \\ n &\mapsto -n \\ \mathbf{1}_m \mathcal{E}^{\alpha_1} \mathcal{F}^{\beta_1} \mathcal{E}^{\alpha_2} \dots \mathcal{E}^{\alpha_k} \mathcal{F}^{\beta_k} \mathbf{1}_n \{s\} &\mapsto \mathbf{1}_{-m} \mathcal{F}^{\alpha_1} \mathcal{E}^{\beta_1} \mathcal{F}^{\alpha_2} \dots \mathcal{F}^{\alpha_k} \mathcal{E}^{\beta_k} \mathbf{1}_{-n} \{s\}. \end{aligned} \quad (3.47)$$

This 2-functor extends to a 2-endofunctor

$$\begin{aligned} \omega: \text{Kom}(\mathcal{U}) &\rightarrow \text{Kom}(\mathcal{U}), \\ n &\mapsto -n \\ (X, d) &\mapsto \dots \longrightarrow \omega(X^{i-1}) \xrightarrow{\omega(d_{i-1})} \omega(X^i) \xrightarrow{\omega(d_i)} \omega(X^{i+1}) \longrightarrow \dots \\ f_i: X \rightarrow Y &\mapsto \omega(f_i): \omega(X) \rightarrow \omega(Y), \end{aligned} \quad (3.48)$$

and a 2-endofunctor on all the other 2-categories in (3.44).

Rescale, reflect across the vertical axis, and send $n \mapsto -n$: The operation on diagrams that rescales the crossing $\begin{array}{c} \nearrow \\ \searrow \end{array} \mapsto - \begin{array}{c} \searrow \\ \nearrow \end{array}$, reflects a diagram across the vertical axis, and sends n to $-n$ leaves invariant the relations on the 2-morphisms of \mathcal{U} . This operation

is contravariant for composition of 1-morphisms, covariant for composition of 2-morphisms, and preserves the degree of a diagram. This symmetry gives an invertible 2-functor

$$\begin{aligned} \sigma: \mathcal{U} &\rightarrow \mathcal{U}^{\text{op}}, \\ n &\mapsto -n \\ \mathbf{1}_m \mathcal{E}^{\alpha_1} \mathcal{F}^{\beta_1} \mathcal{E}^{\alpha_2} \dots \mathcal{E}^{\alpha_k} \mathcal{F}^{\beta_k} \mathbf{1}_n \{s\} &\mapsto \mathbf{1}_{-n} \mathcal{F}^{\beta_k} \mathcal{E}^{\alpha_k} \mathcal{F}^{\beta_{k-1}} \dots \mathcal{F}^{\beta_1} \mathcal{E}^{\alpha_1} \mathbf{1}_{-m} \{s\} \end{aligned}$$

that acts on 2-morphisms via the symmetry described above. This 2-functor extends to a 2-functor

$$\begin{aligned} \sigma: \text{Kom}(\mathcal{U}) &\rightarrow \text{Kom}(\mathcal{U}) \\ n &\mapsto -n \\ (X, d) &\mapsto \dots \longrightarrow \sigma(X^{i-1}) \xrightarrow{\sigma(d_{i-1})} \sigma(X^i) \xrightarrow{\sigma(d_i)} \sigma(X^{i+1}) \longrightarrow \dots \\ f_i: X \rightarrow Y &\mapsto \sigma(f_i): \sigma(X) \rightarrow \sigma(Y), \end{aligned} \quad (3.49)$$

and, likewise, a 2-endofunctor on all the other 2-categories from (3.44).

Reflect across the x-axis and invert orientation: Here we are careful to keep track of what happens to the shifts of sources and targets

$$\psi \left(\begin{array}{c} \text{Diagram with strands } m, n, \{t\}, \{t'\} \end{array} \right) = \begin{array}{c} \text{Diagram with strands } m, n, \{-t\}, \{-t'\} \end{array}$$

Shift reversals on the right hand side are required for this transformation to preserve the degree of a diagram. This gives an invertible 2-functor

$$\begin{aligned} \psi: \mathcal{U} &\rightarrow \mathcal{U}^{\text{co}}, \\ n &\mapsto n \\ \mathbf{1}_m \mathcal{E}^{\alpha_1} \mathcal{F}^{\beta_1} \mathcal{E}^{\alpha_2} \dots \mathcal{E}^{\alpha_k} \mathcal{F}^{\beta_k} \mathbf{1}_n \{s\} &\mapsto \mathbf{1}_m \mathcal{E}^{\alpha_1} \mathcal{F}^{\beta_1} \mathcal{E}^{\alpha_2} \dots \mathcal{E}^{\alpha_k} \mathcal{F}^{\beta_k} \mathbf{1}_n \{-s\}, \end{aligned} \quad (3.50)$$

and on 2-morphisms ψ reflects the diagrams across the x -axis and inverts the orientation.

Since ψ is contravariant on 2-morphisms in \mathcal{U} , this 2-functor extends to a 2-functor

$$\begin{aligned} \psi: \text{Kom}(\mathcal{U}) &\rightarrow \text{Kom}(\mathcal{U}), \\ n &\mapsto n \\ (X, d) &\mapsto \dots \longrightarrow \psi(X^{i+1}) \xrightarrow{\psi(d_i)} \psi(X^i) \xrightarrow{\psi(d_{i-1})} \psi(X^{i-1}) \longrightarrow \dots \\ f_i: X \rightarrow Y &\mapsto \psi(f_i): \psi(Y) \rightarrow \psi(X), \end{aligned} \quad (3.51)$$

and, likewise, a 2-endofunctor on all 2-categories in (3.44). Notice that ψ inverts the homological degree so that ψ acts on a complex (X, d) in $Kom(\mathcal{U})$ by $(\psi X)^i = \psi(X^{-i})$.

These 2-functors are involutive and commute with each other ‘on-the-nose’:

$$\omega\sigma = \omega\sigma, \quad \sigma\psi = \psi\sigma, \quad \omega\psi = \psi\omega, \quad (3.52)$$

generating a group $\mathcal{G} = (\mathbb{Z}_2)^3$ of 2-functors acting on all the 2-categories in (3.44). Equivalences in table (3.44) respect this action. On the Grothendieck group of \mathcal{U} the 2-functors ψ, ω, σ descend to (anti)involutions $\underline{\psi}, \underline{\omega},$ and $\underline{\sigma}$ on ${}_{\mathcal{A}}\dot{\mathcal{U}}$.

4 The Casimir complex

4.1 The Casimir complex and its symmetries

We sometimes represent the Casimir complex (1.6) using the notation

$$\mathcal{C}\mathbf{1}_n := \left(\begin{array}{c} \uparrow \downarrow \quad \smile \\ \uparrow \downarrow \quad \smile \end{array} \right) \xrightarrow{\mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} \oplus \mathbf{1}_n\{1-n\}} \mathcal{E}\mathcal{F}\mathbf{1}_n \oplus \mathcal{E}\mathcal{F}\mathbf{1}_n \left(\begin{array}{c} - \uparrow \downarrow \quad \uparrow \downarrow \\ \curvearrowright \quad - \quad \curvearrowright \end{array} \right) \xrightarrow{\mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \oplus \mathbf{1}_n\{n-1\}} \mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \oplus \mathbf{1}_n\{n-1\}.$$

or

$$\mathcal{C}\mathbf{1}_n := \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1-n\} \end{array} \right) \left(\begin{array}{c} \uparrow \downarrow \quad \smile \\ \uparrow \downarrow \quad \smile \end{array} \right) \xrightarrow{\mathcal{E}\mathcal{F}\mathbf{1}_n} \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathbf{1}_n \\ \mathcal{E}\mathcal{F}\mathbf{1}_n \end{array} \right) \left(\begin{array}{c} - \uparrow \downarrow \quad \uparrow \downarrow \\ \curvearrowright \quad - \quad \curvearrowright \end{array} \right) \xrightarrow{\mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\}} \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{n-1\} \end{array} \right) \quad (4.1)$$

We will interchange freely between these methods of depicting the complex $\mathcal{C}\mathbf{1}_n$.

The placement of minus signs in the above complex is arbitrary as long as each square anticommutes. In fact, we get different placements of the minus signs and dots using the symmetry 2-functors defined in Section 3.3:

$$\psi(\mathcal{C}\mathbf{1}_n) = \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1-n\} \end{array} \right) \left(\begin{array}{c} - \uparrow \downarrow \quad \smile \\ \uparrow \downarrow \quad - \smile \end{array} \right) \xrightarrow{\mathcal{E}\mathcal{F}\mathbf{1}_n} \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathbf{1}_n \\ \mathcal{E}\mathcal{F}\mathbf{1}_n \end{array} \right) \left(\begin{array}{c} \uparrow \downarrow \quad \uparrow \downarrow \\ \curvearrowright \quad \curvearrowright \end{array} \right) \xrightarrow{\mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\}} \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{n-1\} \end{array} \right) \quad (4.2)$$

$$\sigma\omega(\mathcal{C}\mathbf{1}_n) = \begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1-n\} \end{pmatrix} \xrightarrow{\begin{pmatrix} \uparrow\downarrow \cdot & \cup \\ \uparrow\uparrow & \cup \end{pmatrix}} \underline{\begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n \\ \mathcal{E}\mathcal{F}\mathbf{1}_n \end{pmatrix}} \xrightarrow{\begin{pmatrix} -\uparrow\downarrow & \uparrow\downarrow \\ \curvearrowright & -\curvearrowright \end{pmatrix}} \begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{n-1\} \end{pmatrix} \quad (4.3)$$

$$\sigma\omega\psi(\mathcal{C}\mathbf{1}_n) = \begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1-n\} \end{pmatrix} \xrightarrow{\begin{pmatrix} -\uparrow\downarrow & \cup \\ \uparrow\downarrow \cdot & -\cup \end{pmatrix}} \underline{\begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n \\ \mathcal{E}\mathcal{F}\mathbf{1}_n \end{pmatrix}} \xrightarrow{\begin{pmatrix} \uparrow\downarrow & \uparrow\downarrow \\ \curvearrowright & \curvearrowright \end{pmatrix}} \begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{n-1\} \end{pmatrix} \quad (4.4)$$

It is trivial to check (see also Proposition 4.1 below) that these complexes are all isomorphic to the Casimir complex $\mathcal{C}\mathbf{1}_n$. We will write $\mathcal{G}_1 = \{\text{Id}, \psi, \omega\sigma, \psi\omega\sigma\}$ for the subgroup of symmetry 2-functors in $\mathcal{G} = (\mathbb{Z}_2)^3$ that preserve the Casimir complex.

Just as the symmetries in $G \setminus G_1 = \{\underline{\omega}, \underline{\sigma}, \underline{\psi\omega}, \underline{\psi\sigma}\}$ interchange between the two forms of the Casimir element

$$\begin{array}{ccc} & G \setminus G_1 & \\ \curvearrowright & & \curvearrowleft \\ (-q^2 + 2 - q^{-2})EF - q^{-1}K - qK^{-1} & & (-q^2 + 2 - q^{-2})FE - qK - q^{-1}K^{-1}, \\ \curvearrowleft & & \curvearrowright \\ & G \setminus G_1 & \end{array}$$

we can write down a categorification of the idempotented Casimir element in the form

$$(-q^2 + 2 - q^{-2})FE\mathbf{1}_n - (q^{-n-1} + q^{1+n})\mathbf{1}_n \quad (4.5)$$

by applying symmetry 2-functors in $\mathcal{G} \setminus \mathcal{G}_1 = \{\omega, \sigma, \psi\omega, \psi\sigma\}$ to the Casimir complex $\mathcal{C}\mathbf{1}_{-n}$. Depending on which 2-functor in $\mathcal{G} \setminus \mathcal{G}_1$ is chosen, we will get a different placement of minus signs and dots:

$$\sigma(\mathcal{C}\mathbf{1}_{-n}) = \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1+n\} \end{pmatrix} \xrightarrow{\begin{pmatrix} \downarrow\uparrow \cdot & \cup \\ \downarrow\uparrow & \cup \end{pmatrix}} \underline{\begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n \\ \mathcal{F}\mathcal{E}\mathbf{1}_n \end{pmatrix}} \xrightarrow{\begin{pmatrix} -\downarrow\uparrow & \downarrow\uparrow \\ \curvearrowleft & -\curvearrowleft \end{pmatrix}} \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{-n-1\} \end{pmatrix} \quad (4.6)$$

$$\omega(\mathcal{C}\mathbf{1}_{-n}) = \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1+n\} \end{pmatrix} \xrightarrow{\begin{pmatrix} \downarrow\uparrow & \cup \\ \downarrow\uparrow & \cup \end{pmatrix}} \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n \\ \mathcal{F}\mathcal{E}\mathbf{1}_n \end{pmatrix} \xrightarrow{\begin{pmatrix} -\downarrow\uparrow & \downarrow\uparrow \\ \curvearrowright & -\curvearrowright \end{pmatrix}} \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{-n-1\} \end{pmatrix} \quad (4.7)$$

$$\sigma\psi(\mathcal{C}\mathbf{1}_{-n}) = \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1+n\} \end{pmatrix} \xrightarrow{\begin{pmatrix} -\downarrow\uparrow & \cup \\ \downarrow\uparrow & -\cup \end{pmatrix}} \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n \\ \mathcal{F}\mathcal{E}\mathbf{1}_n \end{pmatrix} \xrightarrow{\begin{pmatrix} \downarrow\uparrow & \downarrow\uparrow \\ \curvearrowright & \curvearrowright \end{pmatrix}} \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{-n-1\} \end{pmatrix} \quad (4.8)$$

$$\omega\psi(\mathcal{C}\mathbf{1}_{-n}) = \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1+n\} \end{pmatrix} \xrightarrow{\begin{pmatrix} -\downarrow\uparrow & \cup \\ \downarrow\uparrow & -\cup \end{pmatrix}} \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n \\ \mathcal{F}\mathcal{E}\mathbf{1}_n \end{pmatrix} \xrightarrow{\begin{pmatrix} \downarrow\uparrow & \downarrow\uparrow \\ \curvearrowright & \curvearrowright \end{pmatrix}} \begin{pmatrix} \mathcal{F}\mathcal{E}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{-n-1\} \end{pmatrix} \quad (4.9)$$

The first complex is identical to the complex $\mathcal{C}'\mathbf{1}_n$ in (1.7). The four complexes above are isomorphic to one another in $Kom(\dot{\mathcal{U}})$. In Proposition 4.1 below we will show that they are homotopy equivalent to the complex $\mathcal{C}\mathbf{1}_n$.

Proposition 4.1.

- a) For any $g \in \mathcal{G}_1$ and $g' \in \mathcal{G} \setminus \mathcal{G}_1$ there are chain isomorphisms $\mathcal{C}\mathbf{1}_n \simeq g(\mathcal{C}\mathbf{1}_n)$ and homotopy equivalences $\mathcal{C}\mathbf{1}_n \simeq g'(\mathcal{C}\mathbf{1}_{-n})$ given by chain maps

$$\begin{array}{ccc} & \sigma(\mathcal{C}\mathbf{1}_{-n}) & \xrightarrow{\sigma(\varrho^\psi)} & \sigma\psi(\mathcal{C}\mathbf{1}_{-n}) \\ & \nearrow \varrho^\sigma & \downarrow \sigma(\varrho^{\sigma\omega}) & \nearrow \psi(\widehat{\varrho}^\sigma) \\ \mathcal{C}\mathbf{1}_n & \xrightarrow{\varrho^\psi} & \psi(\mathcal{C}\mathbf{1}_n) & \downarrow \psi\sigma(\widehat{\varrho}^{\sigma\omega}) \\ & \searrow \varrho^\omega & \downarrow \psi(\widehat{\varrho}^\omega) & \\ & \omega(\mathcal{C}\mathbf{1}_{-n}) & \xrightarrow{\omega(\varrho^\psi)} & \omega\psi(\mathcal{C}\mathbf{1}_{-n}) \\ & \nearrow \omega(\widehat{\varrho}^\sigma) & \downarrow \psi(\widehat{\varrho}^{\sigma\omega}) & \nearrow \psi\omega(\varrho^\sigma) \\ \sigma\omega(\mathcal{C}\mathbf{1}_n) & \xrightarrow{\sigma\omega(\varrho^\psi)} & \sigma\omega\psi(\mathcal{C}\mathbf{1}_n) & \end{array} \quad (4.10)$$

where the solid lines represent chain isomorphisms and the dotted lines represent chain homotopy equivalences. We denote the inverse of a map ϱ^g by $\widehat{\varrho}^g$ for $g \in \mathcal{G}$.

- b) The triangles on the left and right in (4.10) commute and the four remaining squares anticommute.
- c) Complexes $\mathcal{C}\mathbf{1}_n$ and $\mathcal{C}'\mathbf{1}_n$ are homotopy equivalent.

Proof. We define explicit chain isomorphisms

$$\mathcal{C}\mathbf{1}_n \begin{array}{c} \xrightarrow{\varrho^\psi} \\ \xleftarrow{\widehat{\varrho}^\psi} \end{array} \psi(\mathcal{C}\mathbf{1}_n) \quad \mathcal{C}\mathbf{1}_n \begin{array}{c} \xrightarrow{\varrho^{\sigma\omega}} \\ \xleftarrow{\widehat{\varrho}^{\sigma\omega}} \end{array} \sigma\omega(\mathcal{C}\mathbf{1}_n),$$

as well as a homotopy equivalence

$$\mathcal{C}\mathbf{1}_n \begin{array}{c} \xrightarrow{\varrho^\sigma} \\ \xleftarrow{\widehat{\varrho}^\sigma} \end{array} \sigma(\mathcal{C}\mathbf{1}_n).$$

While the maps ϱ^ψ and $\varrho^{\sigma\omega}$ are rather uninteresting, it is convenient to fix them.

Define $\varrho^\psi: \mathcal{C}\mathbf{1}_n \rightarrow \psi(\mathcal{C}\mathbf{1}_n)$ and its inverse $\widehat{\varrho}^\psi: \psi(\mathcal{C}\mathbf{1}_n) \rightarrow \mathcal{C}\mathbf{1}_n$ by

$$(\varrho^\psi)_{-1} := \begin{pmatrix} -\uparrow\downarrow & 0 \\ 0 & 1 \end{pmatrix} \quad (\varrho^\psi)_0 := \begin{pmatrix} 0 & \uparrow\downarrow \\ -\uparrow\downarrow & 0 \end{pmatrix} \quad (\varrho^\psi)_{+1} := \begin{pmatrix} \uparrow\downarrow & 0 \\ 0 & -1 \end{pmatrix} \quad (4.11)$$

$$(\widehat{\varrho}^\psi)_{-1} := \begin{pmatrix} -\uparrow\downarrow & 0 \\ 0 & 1 \end{pmatrix} \quad (\widehat{\varrho}^\psi)_0 := \begin{pmatrix} 0 & -\uparrow\downarrow \\ \uparrow\downarrow & 0 \end{pmatrix} \quad (\widehat{\varrho}^\psi)_{+1} := \begin{pmatrix} \uparrow\downarrow & 0 \\ 0 & -1 \end{pmatrix} \quad (4.12)$$

where $\mathcal{C}\mathbf{1}_n$ and $\psi(\mathcal{C}\mathbf{1}_n)$ are given in (4.1) and (4.8). The map $\varrho^{\sigma\omega}: \mathcal{C}\mathbf{1}_n \rightarrow \sigma\omega(\mathcal{C}\mathbf{1}_n)$ and its inverse $\widehat{\varrho}^{\sigma\omega}: \sigma\omega(\mathcal{C}\mathbf{1}_n) \rightarrow \mathcal{C}\mathbf{1}_n$ are given by the chain maps

$$(\varrho^{\sigma\omega})_{-1} := \begin{pmatrix} \uparrow\downarrow & 0 \\ 0 & 1 \end{pmatrix} \quad (\varrho^{\sigma\omega})_0 := \begin{pmatrix} 0 & \uparrow\downarrow \\ \uparrow\downarrow & 0 \end{pmatrix} \quad (\varrho^{\sigma\omega})_{+1} := \begin{pmatrix} -\uparrow\downarrow & 0 \\ 0 & -1 \end{pmatrix} \quad (4.13)$$

$$(\widehat{\varrho}^{\sigma\omega})_{-1} := \begin{pmatrix} \uparrow\downarrow & 0 \\ 0 & 1 \end{pmatrix} \quad (\widehat{\varrho}^{\sigma\omega})_0 := \begin{pmatrix} 0 & \uparrow\downarrow \\ \uparrow\downarrow & 0 \end{pmatrix} \quad (\widehat{\varrho}^{\sigma\omega})_{+1} := \begin{pmatrix} -\uparrow\downarrow & 0 \\ 0 & -1 \end{pmatrix} \quad (4.14)$$

with $\sigma\omega(\mathcal{C}\mathbf{1}_n)$ given by (4.3). We sometimes express chain maps using cube-like diagrams. For example, the (rather obvious) chain maps $\varrho^{\sigma\omega}: \mathcal{C}\mathbf{1}_n \rightarrow \sigma\omega(\mathcal{C}\mathbf{1}_n)$ and $\widehat{\varrho}^{\sigma\omega}: \sigma\omega(\mathcal{C}\mathbf{1}_n) \rightarrow$

$\mathcal{C}1_n$ can be depicted as

$$(4.15)$$

where

$$\rho_1^{\sigma\omega} = \rho_3^{\sigma\omega} = \rho_4^{\sigma\omega} = -\rho_5^{\sigma\omega} = \text{Id}_{\mathcal{E}\mathcal{F}1_n} \quad \rho_2^{\sigma\omega} = -\rho_6^{\sigma\omega} = \text{Id}_{1_n}, \quad (4.16)$$

and

$$\widehat{\rho}_1^{\sigma\omega} = \widehat{\rho}_3^{\sigma\omega} = \widehat{\rho}_4^{\sigma\omega} = -\widehat{\rho}_5^{\sigma\omega} = \text{Id}_{\mathcal{F}\mathcal{E}1_n} \quad \widehat{\rho}_2^{\sigma\omega} = -\widehat{\rho}_6^{\sigma\omega} = \text{Id}_{1_n}. \quad (4.17)$$

The interesting maps here are $\rho^\sigma : \mathcal{C}1_n \rightarrow \sigma(\mathcal{C}1_{-n})$ and its homotopy inverse $\widehat{\rho}^\sigma : \sigma(\mathcal{C}1_{-n}) \rightarrow \mathcal{C}1_n$, given by the diagram

$$(4.18)$$

where

$$\begin{aligned}
 \varrho_1^\sigma &= \varrho_5^\sigma = \varrho_6^\sigma = \varrho_7^\sigma = \begin{array}{c} \diagup \\ \diagdown \end{array} & \varrho_2^\sigma &= - \begin{array}{c} \curvearrowright \end{array} & \varrho_3^\sigma &= - \sum_{\substack{f_1+f_2 \\ =-n-1}} \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \\
 \varrho_4^\sigma &= - \begin{array}{c} \curvearrowleft \\ \bullet \\ \curvearrowleft \end{array} & \varrho_8^\sigma &= - \sum_{\substack{f_1+f_2 \\ =n-1}} \begin{array}{c} \curvearrowleft \\ \bullet \\ \curvearrowleft \\ \bullet \\ \curvearrowleft \end{array} & \varrho_9^\sigma &= - \begin{array}{c} \curvearrowleft \end{array} & \varrho_{10}^\sigma &= \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \\
 \widehat{\varrho}_1^\sigma &= \widehat{\varrho}_5^\sigma = \widehat{\varrho}_6^\sigma = \widehat{\varrho}_7^\sigma = - \begin{array}{c} \diagup \\ \diagdown \end{array} & \widehat{\varrho}_2^\sigma &= - \begin{array}{c} \curvearrowright \end{array} & \widehat{\varrho}_3^\sigma &= - \sum_{\substack{f_1+f_2 \\ =n-1}} \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \\
 \widehat{\varrho}_4^\sigma &= - \begin{array}{c} \curvearrowleft \\ \bullet \\ \curvearrowleft \end{array} & \widehat{\varrho}_8^\sigma &= - \begin{array}{c} \curvearrowleft \end{array} & \widehat{\varrho}_9^\sigma &= - \sum_{\substack{f_1+f_2 \\ =-n-1}} \begin{array}{c} \curvearrowleft \\ \bullet \\ \curvearrowleft \\ \bullet \\ \curvearrowleft \end{array} & \widehat{\varrho}_{10}^\sigma &= \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \end{array}
 \end{aligned}$$

The chain homotopies $\varrho^\sigma \widehat{\varrho}^\sigma - \text{Id} \simeq 0$, $\widehat{\varrho}^\sigma \varrho^\sigma - \text{Id} \simeq 0$ are given by

$$\begin{array}{ccccc}
 \mathcal{EF}1_n\{2\} & \xrightarrow{\uparrow\downarrow} & \mathcal{EF}1_n & \xrightarrow{-\uparrow\downarrow} & \mathcal{EF}1_n\{-2\} \\
 \uparrow\downarrow & \searrow & \curvearrowright & \searrow & \uparrow\downarrow \\
 \mathbf{1}_n\{1-n\} & \xrightarrow{\curvearrowright} & \mathcal{EF}1_n & \xrightarrow{\curvearrowright} & \mathbf{1}_n\{n-1\} \\
 & & & & \curvearrowright \\
 & & & & \mathbf{1}_n\{n-1\}
 \end{array}
 \begin{array}{c}
 \xrightarrow{h_1} \\
 \xrightarrow{h_2} \\
 \xrightarrow{h_3} \\
 \xrightarrow{h_4}
 \end{array}$$

$$\begin{array}{ccccc}
 \mathcal{FE}1_n\{2\} & \xrightarrow{\downarrow\uparrow} & \mathcal{FE}1_n & \xrightarrow{-\downarrow\uparrow} & \mathcal{FE}1_n\{-2\} \\
 \downarrow\uparrow & \searrow & \curvearrowright & \searrow & \downarrow\uparrow \\
 \mathbf{1}_n\{1+n\} & \xrightarrow{\curvearrowright} & \mathcal{FE}1_n & \xrightarrow{\curvearrowright} & \mathbf{1}_n\{-n-1\} \\
 & & & & \curvearrowright \\
 & & & & \mathbf{1}_n\{-n-1\}
 \end{array}
 \begin{array}{c}
 \xrightarrow{h'_1} \\
 \xrightarrow{h'_2} \\
 \xrightarrow{h'_3} \\
 \xrightarrow{h'_4}
 \end{array}
 \tag{4.19}$$

where

$$h_1 = h_3 = - \sum_{\substack{f_1+f_2+f_3 \\ =n-2}} \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \quad h_2 = - \sum_{\substack{f_1+f_2 \\ =n-1}} \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \quad h_4 = \sum_{\substack{f_1+f_2 \\ =n-1}} \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \quad (4.20)$$

$$h'_1 = h'_3 = - \sum_{\substack{g_1+g_2+g_3 \\ =x+y-n-2}} \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \quad h'_2 = - \sum_{\substack{f_1+f_2 \\ =-n-1}} \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \quad h'_4 = \sum_{\substack{f_1+f_2 \\ =-n-1}} \begin{array}{c} \curvearrowright \\ \bullet \\ \curvearrowright \\ \bullet \\ \curvearrowright \end{array} \quad (4.21)$$

One can verify the following equations

$$\varrho^\sigma \widehat{\varrho}^\sigma - \text{Id} = hd + dh, \quad (4.22)$$

$$\widehat{\varrho}^\sigma \varrho^\sigma - \text{Id} = h'd + dh'. \quad (4.23)$$

The chain maps just defined satisfy

$$\sigma(\varrho^\sigma) = \widehat{\varrho}^\sigma, \quad \sigma(\widehat{\varrho}^\sigma) = \varrho^\sigma, \quad (4.24)$$

$$\psi(\varrho^\psi) = -\varrho^\psi, \quad \psi(\widehat{\varrho}^\psi) = -\widehat{\varrho}^\psi, \quad (4.25)$$

$$\omega(\varrho^{\sigma\omega}) = \sigma(\widehat{\varrho}^{\sigma\omega}), \quad \omega(\widehat{\varrho}^{\sigma\omega}) = \sigma(\varrho^{\sigma\omega}). \quad (4.26)$$

To prove part b of the proposition one can check by direct computation that the front square in (4.10) anticommutes. The back solid square is just σ applied to the front solid square so it also anticommutes. The leftmost square commutes on the nose and we define maps

$$\varrho^\omega := \sigma(\varrho^{\sigma\omega}) \circ \varrho^\sigma \quad \widehat{\varrho}^\omega := \widehat{\varrho}^\sigma \circ \sigma(\widehat{\varrho}^{\sigma\omega}), \quad (4.27)$$

using these commutative squares. The rightmost square is just ψ applied to the leftmost square so it also commutes on the nose. The top square can be shown to anticommute. After observing that $\psi\omega(\varrho^\sigma) = \psi\omega\sigma(\widehat{\varrho}^\sigma)$ and $\omega(\widehat{\varrho}^\sigma) = \omega\sigma(\varrho^\sigma)$ the anticommutativity of the bottom square follows since it is just $\omega\sigma$ applied to the top square.

Part c follows immediately from part a since $\mathcal{C}'\mathbf{1}_n = \sigma(\mathcal{C}\mathbf{1}_{-n})$. \square

Theorem 3.9 and the results in section 3.2 imply that

$$K_0(Kar(Kom(\mathcal{U}))) \cong K_0(Kom(Kar(\mathcal{U}))) = K_0(Kar(\mathcal{U})) \cong K_0(\dot{\mathcal{U}}) \cong {}_{\mathcal{A}}\dot{\mathcal{U}}. \quad (4.28)$$

Under this isomorphism

$$[\mathcal{C}\mathbf{1}_n] = C\mathbf{1}_n. \quad (4.29)$$

4.2 Commutativity of the Casimir complex

4.2.1 Commutativity chain maps ξ^- and $\widehat{\xi}^-$

Definition 4.2. Define chain maps $\xi^- = \xi^- \mathbf{1}_n: \mathcal{FC}\mathbf{1}_n \rightarrow \mathcal{CF}\mathbf{1}_n$ and $\widehat{\xi}^- = \widehat{\xi}^- \mathbf{1}_n: \mathcal{CF}\mathbf{1}_n \rightarrow \mathcal{FC}\mathbf{1}_n$ as follows:

(4.30)

where

$$\begin{aligned}
 \xi_1^- &= \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n-2 \\ 1 \leq f_3}} \begin{array}{c} f_1 \\ f_2 \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \\
 \xi_2^- &= \begin{array}{c} \uparrow \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_3}} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \\
 \xi_3^- &= \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n-2}} \begin{array}{c} f_1 \\ f_2 \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \\
 \xi_4^- &= \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \\
 \xi_5^- &= \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} f_1 \\ f_3 \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \\
 \xi_6^- &= \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_1}} \begin{array}{c} f_1 \\ f_3 \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \\
 \xi_7^- &= - \sum_{\substack{f_1+f_2=n \\ 1 \leq f_1}} \begin{array}{c} f_1 \\ f_2 \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \\
 \xi_8^- &= \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} + \delta_{n,1} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \\
 \xi_9^- &= \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \delta_{n,1} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array}
 \end{aligned}$$

The map $\widehat{\xi}^- \mathbf{1}_n$ is defined via $\xi^- \mathbf{1}_n$ using symmetry 2-functors:

$$\widehat{\xi}^- \mathbf{1}_n := \mathcal{F} \widehat{\varrho}^\psi \circ \psi(\xi^- \mathbf{1}_n) \circ \varrho^\psi \mathcal{F}. \quad (4.31)$$

More explicitly,

$$\begin{aligned} \widehat{\xi}_1^- &= \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_1}} \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} & \widehat{\xi}_2^- &= \sum_{\substack{f_1+f_2 \\ =n-1 \\ 1 \leq f_1}} \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} \\ \widehat{\xi}_3^- &= - \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} + \delta_{n,1} \begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array} & \widehat{\xi}_4^- &= \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \end{array} + \delta_{n,1} \begin{array}{c} \text{Diagram 13} \\ \text{Diagram 14} \end{array} \\ \widehat{\xi}_5^- &= \begin{array}{c} \text{Diagram 15} \\ \text{Diagram 16} \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \text{Diagram 17} \\ \text{Diagram 18} \end{array} & \widehat{\xi}_6^- &= - \begin{array}{c} \text{Diagram 19} \\ \text{Diagram 20} \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \text{Diagram 21} \\ \text{Diagram 22} \end{array} \\ \widehat{\xi}_7^- &= \begin{array}{c} \text{Diagram 23} \\ \text{Diagram 24} \end{array} - \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n-2}} \begin{array}{c} \text{Diagram 25} \\ \text{Diagram 26} \end{array} & & \\ \widehat{\xi}_8^- &= \begin{array}{c} \text{Diagram 27} \\ \text{Diagram 28} \end{array} - \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n-2 \\ 1 \leq f_3}} \begin{array}{c} \text{Diagram 29} \\ \text{Diagram 30} \end{array} & \widehat{\xi}_9^- &= - \begin{array}{c} \text{Diagram 31} \\ \text{Diagram 32} \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_3}} \begin{array}{c} \text{Diagram 33} \\ \text{Diagram 34} \end{array} \end{aligned}$$

Remark 4.3. Note that when $n \leq 0$ all terms involving bubbles are zero.

Proposition 4.4. The maps ξ^- and $\widehat{\xi}^-$ defined above are mutually-inverse chain homotopy equivalences between $\mathcal{FC}\mathbf{1}_n$ and $\mathcal{CF}\mathbf{1}_n$.

Proof. To check that the maps ξ^- and $\widehat{\xi}^-$ are chain maps it suffices to verify this claim for the map ξ^- since $\widehat{\xi}^-$ is defined from ξ^- via (4.31). This is proven by direct computation.

Below we give explicit chain homotopies $\text{Id} - \xi^- \widehat{\xi}^- = hd + dh$ and $\text{Id} - \widehat{\xi}^- \xi^- = h'd + dh'$ showing that ξ^- and $\widehat{\xi}^-$ are mutually-inverse homotopy equivalences.

To simplify notation we write

$$\Theta(n) := \begin{cases} 1 & \text{if } n \geq 0, \\ 0 & \text{if } n < 0. \end{cases} \quad (4.33)$$

$$h_1 = \uparrow \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} + \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n-3}} \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \quad (4.34)$$

$$h_2 = \sum_{\substack{f_1+f_2+f_3 \\ =n-2}} \begin{array}{c} \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \quad -h_3 = h_4 = \uparrow \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \quad (4.35)$$

$$h_5 = \uparrow \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} + \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n-3}} \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \quad h_6 = - \sum_{\substack{f_1+f_2+f_3 \\ =n-2}} \begin{array}{c} \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \quad (4.36)$$

$$h'_1 = h'_6 = - \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \quad h'_2 = -h'_5 = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-2}} \begin{array}{c} \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \quad (4.37)$$

$$h'_3 = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} - \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \quad n = \Theta(n-1) \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \quad n - \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \quad (4.38)$$

$$h'_4 = - \text{diagram} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_1}} \text{diagram} = \text{diagram} - \sum_{f_1+f_2=n-1} \text{diagram} \quad (4.39)$$

$$h'_7 = - \text{diagram} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_1}} \text{diagram} = \text{diagram} - \sum_{f_1+f_2=n-1} \text{diagram} \quad (4.40)$$

$$h'_8 = - \text{diagram} + \text{diagram} \cdot n = \Theta(n-1) \text{diagram} - \sum_{f_1+f_2+f_3=n-1} \text{diagram} \cdot n \quad (4.41)$$

The rather nontrivial computation that these maps are the required homotopies is omitted from the paper. \square

4.2.2 Commutativity chain maps ξ^+ and $\widehat{\xi}^+$

Definition 4.5. We introduce chain maps

$$\xi^+ = \xi^+ \mathbf{1}_n := \mathcal{E} \widehat{\varrho}^{\sigma\omega} \circ \omega\sigma(\xi^- \mathbf{1}_n) \circ \varrho^{\sigma\omega} \mathcal{E} : \mathcal{CE} \mathbf{1}_n \rightarrow \mathcal{E} \mathcal{C} \mathbf{1}_n \quad (4.42)$$

and

$$\widehat{\xi}^+ = \widehat{\xi}^+ \mathbf{1}_n := \widehat{\varrho}^{\sigma\omega} \mathcal{E} \circ \omega\sigma(\widehat{\xi}^- \mathbf{1}_n) \circ \mathcal{E} \varrho^{\sigma\omega} : \mathcal{E} \mathcal{C} \mathbf{1}_n \rightarrow \mathcal{CE} \mathbf{1}_n. \quad (4.43)$$

The diagrams

$$\begin{array}{ccc} \mathcal{CE} \mathbf{1}_n & \xrightarrow{\xi^+} & \mathcal{E} \mathcal{C} \mathbf{1}_n \\ \varrho^{\sigma\omega} \mathcal{E} \downarrow & & \uparrow \mathcal{E} \widehat{\varrho}^{\sigma\omega} \\ \sigma\omega(\mathcal{C} \mathbf{1}_n) \mathcal{E} \mathbf{1}_n & \xrightarrow{\omega\sigma(\xi^-)} & \mathcal{E} \sigma\omega(\mathcal{C} \mathbf{1}_n) \mathbf{1}_n \end{array} \quad \begin{array}{ccc} \mathcal{E} \mathcal{C} \mathbf{1}_n & \xrightarrow{\widehat{\xi}^+} & \mathcal{CE} \mathbf{1}_n \\ \mathcal{E} \varrho^{\sigma\omega} \downarrow & & \uparrow \widehat{\varrho}^{\sigma\omega} \mathcal{E} \\ \mathcal{E} \sigma\omega(\mathcal{C} \mathbf{1}_n) \mathbf{1}_n & \xrightarrow{\omega\sigma(\widehat{\xi}^-)} & \sigma\omega(\mathcal{C} \mathbf{1}_n) \mathcal{E} \mathbf{1}_n \end{array} \quad (4.44)$$

commute by definition.

Proposition 4.6. The maps ξ^+ and $\widehat{\xi}^+$ defined above are mutually-inverse chain homotopy equivalences between $\mathcal{E} \mathcal{C} \mathbf{1}_n$ and $\mathcal{CE} \mathbf{1}_n$.

Proof. The Proposition follows at once from (4.42), (4.43), and Proposition 4.4. \square

4.3 Indecomposability

4.3.1 Indecomposability of Casimir complexes

Since $\mathbf{1}_n$ is indecomposable for all n , morphisms $\mathbf{1}_n\{1-n\}$, $\mathbf{1}_n\{n-1\}$ are as well. The 1-morphisms $\mathcal{EF}\mathbf{1}_n\{-2\}$, $\mathcal{EF}\mathbf{1}_n$, and $\mathcal{EF}\mathbf{1}_n\{2\}$ appearing in various direct summands of the complex $\mathcal{C}\mathbf{1}_n$ are indecomposable when $n \leq 0$. Assuming $n \leq 0$ each of the eight maps describing the differential in $\mathcal{C}\mathbf{1}_n$ belongs to the graded Jacobson radical of the category $\dot{\mathcal{U}}(n, n)$. This implies that $\mathcal{C}\mathbf{1}_n$ does not contain any contractible summands if $n \leq 0$.

Proposition 4.7. The complex $\mathcal{C}\mathbf{1}_n$ given by (1.6) is indecomposable in $Kom(\dot{\mathcal{U}})$ if $n \leq 0$.

Proof. Assume $n \leq 0$ so that $\mathcal{C}\mathbf{1}_n$ consists of 6 terms, all indecomposable. Those terms belong to the category $\dot{\mathcal{U}}(n, n)$, which is Krull-Schmidt with finite dimensional hom spaces.

There are no homs (of degree zero) $\mathbf{1}_n\{1-n\} \rightarrow \mathcal{EF}\mathbf{1}_n\{2\}$ for any n . The degree zero hom space $\mathcal{EF}\mathbf{1}_n\{2\} \rightarrow \mathbf{1}_n\{1-n\}$ (for $n \leq 0$) is nontrivial only when $n = 0$, and then it is spanned by the diagram \curvearrowright . Consequently, $(\mathcal{C}\mathbf{1}_n)^{-1} = \mathcal{EF}\mathbf{1}_n\{2\} \oplus \mathbf{1}_n\{1-n\}$ has only one possible direct sum decomposition for $n < 0$, and a one-parameter family of direct sum decompositions for $n = 0$

$$(\mathcal{C}\mathbf{1}_0)^{-1} = X \oplus \mathbf{1}_0\{1\}, \quad (4.45)$$

where $X \cong \mathcal{EF}\mathbf{1}_0\{2\}$ is the image of $\mathcal{EF}\mathbf{1}_0\{2\}$ in $(\mathcal{C}\mathbf{1}_0)^{-1}$ under the homomorphism

$$\mathcal{EF}\mathbf{1}_0\{2\} \xrightarrow{\begin{pmatrix} \text{Id} \\ a \curvearrowright \end{pmatrix}} \mathcal{EF}\mathbf{1}_0\{2\} \oplus \mathbf{1}_0\{1\} \quad (4.46)$$

for $a \in \mathbb{k}$. Any direct sum decomposition of $(\mathcal{C}\mathbf{1}_n)^0 = \mathcal{EF}\mathbf{1}_n \oplus \mathcal{EF}\mathbf{1}_n$ is determined by a 2×2 invertible matrix with coefficients in \mathbb{k} . There are no homs (of degree zero) $\mathcal{EF}\mathbf{1}_n\{-2\} \rightarrow \mathbf{1}_n\{n-1\}$, and the degree zero hom space $\mathbf{1}_n\{n-1\} \rightarrow \mathcal{EF}\mathbf{1}_n\{-2\}$ is nontrivial only when $n = 0$, and then it is spanned by \curvearrowleft . Therefore, for $n < 0$ direct sum decomposition of $(\mathcal{C}\mathbf{1}_n)^1$ is unique, and for $n = 0$ any direct sum decomposition in $\dot{\mathcal{U}}(0, 0)$ of $(\mathcal{C}\mathbf{1}_0)^1 = \mathcal{EF}\mathbf{1}_0\{-2\} \oplus \mathbf{1}_0\{-1\}$ has the form

$$(\mathcal{C}\mathbf{1}_0)^1 \simeq \mathcal{EF}\mathbf{1}_0\{-2\} \oplus Y \quad (4.47)$$

where $Y \simeq \mathbf{1}_0\{n-1\}$ is the image of $\mathbf{1}_0\{-1\}$ in $(\mathcal{C}\mathbf{1}_0)^1$ under the homomorphism

$$\mathbf{1}_0\{-1\} \xrightarrow{\begin{pmatrix} b \curvearrowleft \\ \text{Id} \end{pmatrix}} \mathcal{EF}\mathbf{1}_0\{-2\} \oplus \mathbf{1}_0\{-1\} \quad (4.48)$$

for some $b \in \mathbb{k}$.

Suppose that for some $n \leq 0$ there exists a nontrivial direct sum decomposition $\mathcal{C}\mathbf{1}_n \simeq \mathcal{C}_1 \oplus \mathcal{C}_2$ in $Kom(\dot{\mathcal{U}})$. Then, from the above discussion, we know that the summand $\mathbf{1}_n\{1-n\} \subset (\mathcal{C}\mathbf{1}_n)^{-1}$ must be either in \mathcal{C}_1 or \mathcal{C}_2 . We can assume it belongs to \mathcal{C}_1 . Applying the

differential in $\mathcal{C}\mathbf{1}_n$ and the classification of direct sum decompositions of $(\mathcal{C}\mathbf{1}_n)^0$ we see that \mathcal{C}_1 must contain the diagonal summand of $(\mathcal{C}\mathbf{1}_n)^0$, the image of

$$\mathcal{E}\mathcal{F}\mathbf{1}_n \xrightarrow{\begin{pmatrix} \text{Id} \\ \text{Id} \end{pmatrix}} \mathcal{E}\mathcal{F}\mathbf{1}_n \oplus \mathcal{E}\mathcal{F}\mathbf{1}_n. \quad (4.49)$$

Further application of the differential and very few available direct sum decompositions of $(\mathcal{C}\mathbf{1}_n)^1$ tell us that \mathcal{C}_1 must contain the summand $\mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\}$ of $(\mathcal{C}\mathbf{1}_n)^1$.

If $(\mathcal{C}_2)^{-1} \neq 0$, then $(\mathcal{C}_2)^{-1} = X$, described above, for some $a \in \mathbb{k}$. Then dX must lie inside a summand of $(\mathcal{C}\mathbf{1}_n)^0$ isomorphic to $\mathcal{E}\mathcal{F}\mathbf{1}_n$. A simple computation shows that this is impossible. Therefore, $(\mathcal{C}_2)^{-1} = 0$ and $(\mathcal{C}_{-1}) = (\mathcal{C}\mathbf{1}_n)^{-1}$. Applying the differential to $(\mathcal{C}_1)^{-1}$ we quickly conclude that $(\mathcal{C}_1)^0 = (\mathcal{C}\mathbf{1}_n)^0$ and then $(\mathcal{C}_1)^1 = (\mathcal{C}\mathbf{1}_n)^1$. Hence $\mathcal{C}_2 = 0$ supplying a contradiction. \square

Corollary 4.8. The complex $\mathcal{C}'\mathbf{1}_n = \sigma(\mathcal{C}\mathbf{1}_{-n})$ is indecomposable in $\text{Kom}(\dot{\mathcal{U}})$ if $n \geq 0$.

Assume $n \leq 0$. Since $\mathcal{C}\mathbf{1}_n$ and $\mathcal{C}'\mathbf{1}_n$ are isomorphic in $\text{Com}(\dot{\mathcal{U}})$ by Proposition 4.1, and $\mathcal{F}\mathcal{E}\mathbf{1}_n \cong \mathcal{E}\mathcal{F}\mathbf{1}_n \oplus_{[n]} \mathbf{1}_n$, we conclude that $\mathcal{C}'\mathbf{1}_n$ is isomorphic in $\text{Kom}(\dot{\mathcal{U}})$ to the direct sum of $\mathcal{C}\mathbf{1}_n$, contractible complexes

$$0 \longrightarrow \mathbf{1}_n\{n-1-2\ell+2\} \xrightarrow{\text{Id}} \mathbf{1}_n\{n-1-2\ell+2\} \longrightarrow 0, \quad 0 \leq \ell \leq n-1 \quad (4.50)$$

concentrated in cohomological degrees -1 and 0, and contractible complexes

$$0 \longrightarrow \mathbf{1}_n\{n-1-2\ell-2\} \xrightarrow{\text{Id}} \mathbf{1}_n\{n-1-2\ell-2\} \longrightarrow 0, \quad 0 \leq \ell \leq n-1 \quad (4.51)$$

concentrated in cohomological degrees 0 and 1. When $n = 0$ complexes $\mathcal{C}\mathbf{1}_n$ and $\mathcal{C}'\mathbf{1}_n$ are isomorphic in $\text{Kom}(\dot{\mathcal{U}})$ via Proposition 4.1.

When $n \geq 0$ there is a similar isomorphism in $\text{Kom}(\dot{\mathcal{U}})$ between $\mathcal{C}\mathbf{1}_n$ and the direct sum of $\mathcal{C}'\mathbf{1}_n$ and contractible complexes.

In the complex $\mathcal{C}'\mathbf{1}_n = \sigma(\mathcal{C}\mathbf{1}_{-n})$ the 1-morphisms $\mathbf{1}_n\{1+n\}$, $\mathbf{1}_n\{-n-1\}$, $\mathcal{F}\mathcal{E}\mathbf{1}_n\{-2\}$, $\mathcal{F}\mathcal{E}\mathbf{1}_n$, and $\mathcal{F}\mathcal{E}\mathbf{1}_n\{2\}$ in the direct summands are all indecomposable when $n \geq 0$. In this way, the symmetry 2-functor σ plays an important role allowing us to switch between complexes $\mathcal{C}\mathbf{1}_n$ and $\mathcal{C}'\mathbf{1}_n$.

The commutativity chain maps studied above reduce drastically when we work with the indecomposable version of the Casimir complex. Below we collect these maps for later convenience.

4.3.2 Chain maps ξ^- and $\widehat{\xi}^-$ in indecomposable case

When $n \leq 0$ the maps in Definition 4.2 simplify to

$$\xi_1^- = \xi_3^- = \begin{array}{c} \uparrow \\ \text{X} \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \text{X} \\ \downarrow \end{array} \quad \xi_2^- = \begin{array}{c} \downarrow \\ \curvearrowright \\ \downarrow \end{array} \quad \xi_4^- = \begin{array}{c} \downarrow \\ \text{X} \\ \downarrow \end{array}$$

$$\begin{aligned}
 \xi_5^- = \xi_6^- &= \begin{array}{c} \uparrow \downarrow \\ \diagdown \diagup \\ \bullet \\ \diagup \diagdown \\ \downarrow \end{array} - \begin{array}{c} \uparrow \downarrow \\ \diagup \diagdown \\ \bullet \\ \diagdown \diagup \\ \downarrow \end{array} & \xi_8^- = - \begin{array}{c} \downarrow \\ \bullet \end{array} & \xi_9^- = - \begin{array}{c} \downarrow \\ \cup \\ \downarrow \end{array} & \xi_7^- = 0 \\
 \hat{\xi}_1^- = \hat{\xi}_5^- &= \begin{array}{c} \downarrow \\ \diagdown \diagup \\ \bullet \\ \diagup \diagdown \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \diagup \diagdown \\ \bullet \\ \diagdown \diagup \\ \downarrow \end{array} & \hat{\xi}_2^- = 0 & \hat{\xi}_3^- = \begin{array}{c} \downarrow \\ \cup \\ \downarrow \\ n \end{array} & \hat{\xi}_4^- = - \begin{array}{c} \downarrow \\ \bullet \end{array} \\
 \hat{\xi}_6^- &= - \begin{array}{c} \downarrow \\ \diagdown \diagup \\ \downarrow \end{array} & \hat{\xi}_7^- = \hat{\xi}_8^- &= \begin{array}{c} \downarrow \\ \diagdown \diagup \\ \bullet \\ \diagup \diagdown \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \diagup \diagdown \\ \bullet \\ \diagdown \diagup \\ \downarrow \end{array} & \hat{\xi}_9^- = - \begin{array}{c} \downarrow \\ \cup \\ \downarrow \end{array}
 \end{aligned}$$

In the diagrams above we have omitted the label n on the far right region of each diagram for simplicity. We will sometimes make use of this convention in the following sections when the labelling of each region is clear from the context.

4.3.3 Chain maps ξ^+ and $\hat{\xi}^+$ in indecomposable case

For $n \leq 0$ the chain maps $\xi^+ : \mathcal{CE}1_n \rightarrow \mathcal{E}1_n$ and $\hat{\xi}^+ : \mathcal{E}1_n \rightarrow \mathcal{CE}1_n$ defined in (4.42) and (4.43) can be written as

$$\begin{array}{ccccc}
 \mathcal{E}\mathcal{E}\mathcal{F}1_n\{2\} & \xrightarrow{\uparrow\uparrow\downarrow} & \mathcal{E}\mathcal{E}\mathcal{F}1_n & \xrightarrow{-\uparrow\uparrow\downarrow} & \mathcal{E}\mathcal{E}\mathcal{F}1_n\{-2\} \\
 \uparrow\uparrow\downarrow & & \uparrow\cup & & \uparrow\cup \\
 \mathcal{E}1_n\{1+n\} & \xrightarrow{\uparrow\cup} & \mathcal{E}\mathcal{E}\mathcal{F}1_n & \xrightarrow{-\uparrow\cup} & \mathcal{E}1_n\{-1-n\} \\
 \uparrow\cup & & \uparrow\cup & & \uparrow\cup \\
 \mathcal{E}\mathcal{F}\mathcal{E}1_n\{2\} & \xrightarrow{\uparrow\downarrow\uparrow} & \mathcal{E}\mathcal{F}\mathcal{E}1_n & \xrightarrow{-\uparrow\downarrow\uparrow} & \mathcal{E}\mathcal{F}\mathcal{E}1_n\{-2\} \\
 \uparrow\downarrow\uparrow & & \uparrow\downarrow\uparrow & & \uparrow\downarrow\uparrow \\
 \mathcal{E}1_n\{n+1\} & \xrightarrow{\cup\uparrow} & \mathcal{E}\mathcal{F}\mathcal{E}1_n & \xrightarrow{-\cup\uparrow} & \mathcal{E}1_n\{n-1\}
 \end{array}$$

(4.52)

$$\begin{aligned}
 \xi_1^+ = \xi_3^+ &= \begin{array}{c} \bullet \\ \diagdown \diagup \\ \downarrow \end{array} - \begin{array}{c} \bullet \\ \diagup \diagdown \\ \downarrow \end{array} & \xi_2^+ &= \begin{array}{c} \uparrow \\ \cup \\ \uparrow \end{array} & \xi_4^+ &= \begin{array}{c} \uparrow \\ \diagdown \diagup \\ \downarrow \end{array} \\
 \xi_5^+ = \xi_6^+ &= \begin{array}{c} \downarrow \\ \diagdown \diagup \\ \bullet \\ \diagup \diagdown \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \diagup \diagdown \\ \bullet \\ \diagdown \diagup \\ \downarrow \end{array} & \xi_7^+ &= - \begin{array}{c} \downarrow \\ \bullet \end{array} & \xi_8^+ &= - \begin{array}{c} \downarrow \\ \cup \\ \downarrow \end{array} \\
 \hat{\xi}_1^+ = \hat{\xi}_4^+ &= \begin{array}{c} \downarrow \\ \diagdown \diagup \\ \bullet \\ \diagup \diagdown \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \diagup \diagdown \\ \bullet \\ \diagdown \diagup \\ \downarrow \end{array} & \hat{\xi}_2^+ &= \begin{array}{c} \cup \\ \uparrow \end{array} & \hat{\xi}_3^+ &= - \begin{array}{c} \downarrow \\ \bullet \\ n \end{array} \\
 \hat{\xi}_5^+ &= - \begin{array}{c} \uparrow \\ \diagdown \diagup \\ \downarrow \end{array} & \hat{\xi}_6^+ = \hat{\xi}_7^+ &= \begin{array}{c} \downarrow \\ \diagdown \diagup \\ \bullet \\ \diagup \diagdown \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \diagup \diagdown \\ \bullet \\ \diagdown \diagup \\ \downarrow \end{array} & \hat{\xi}_8^+ &= - \begin{array}{c} \uparrow \\ \cup \\ \uparrow \end{array}
 \end{aligned}$$

4.3.4 Chain maps $\sigma(\xi^-)$ and $\sigma(\widehat{\xi}^-)$ in indecomposable case

When $n \geq 0$ it is useful to work with the indecomposable in $Kom(\mathcal{U})$ complex $\sigma(\mathcal{C}\mathbf{1}_{-n})$ instead of $\mathcal{C}\mathbf{1}_n$. For $n \geq 0$ the maps

$$\widehat{\xi}_\sigma^- \mathbf{1}_n := \sigma(\widehat{\xi}^- \mathbf{1}_{-n+2}): \mathcal{F}\sigma(\mathcal{C}\mathbf{1}_{-n}) \rightarrow \sigma(\mathcal{C}\mathbf{1}_{-n+2})\mathcal{F}\mathbf{1}_n \quad (4.53)$$

$$\xi_\sigma^- \mathbf{1}_n := \sigma(\xi^- \mathbf{1}_{-n+2}): \sigma(\mathcal{C}\mathbf{1}_{-n+2})\mathcal{F}\mathbf{1}_n \rightarrow \mathcal{F}\sigma(\mathcal{C}\mathbf{1}_{-n}) \quad (4.54)$$

simplify to the form

(4.55)

where

$$\begin{aligned} (\widehat{\xi}_\sigma^-)_1 = (\widehat{\xi}_\sigma^-)_4 &= \begin{array}{c} \text{---} \diagdown \text{---} \\ \text{---} \diagup \text{---} \end{array} - \begin{array}{c} \text{---} \diagup \text{---} \\ \text{---} \diagdown \text{---} \end{array} & (\widehat{\xi}_\sigma^-)_2 &= \begin{array}{c} \cup \\ \downarrow \end{array} & (\widehat{\xi}_\sigma^-)_3 &= \begin{array}{c} \downarrow \\ \bullet \end{array} \\ (\widehat{\xi}_\sigma^-)_5 &= \begin{array}{c} \downarrow \\ \diagdown \end{array} & (\widehat{\xi}_\sigma^-)_6 = (\widehat{\xi}_\sigma^-)_8 &= \begin{array}{c} \text{---} \diagdown \text{---} \\ \text{---} \diagup \text{---} \end{array} - \begin{array}{c} \text{---} \diagup \text{---} \\ \text{---} \diagdown \text{---} \end{array} & (\widehat{\xi}_\sigma^-)_7 &= \begin{array}{c} \downarrow \\ \cup \end{array} \\ (\xi_\sigma^-)_1 = (\xi_\sigma^-)_3 &= \begin{array}{c} \bullet \\ \diagdown \end{array} - \begin{array}{c} \bullet \\ \diagup \end{array} & (\xi_\sigma^-)_2 &= \begin{array}{c} \downarrow \\ \cup \end{array} & (\xi_\sigma^-)_4 &= \begin{array}{c} \downarrow \\ \diagdown \end{array} \\ (\xi_\sigma^-)_5 = (\xi_\sigma^-)_8 &= \begin{array}{c} \text{---} \diagdown \text{---} \\ \text{---} \diagup \text{---} \end{array} - \begin{array}{c} \text{---} \diagup \text{---} \\ \text{---} \diagdown \text{---} \end{array} & (\xi_\sigma^-)_6 &= \begin{array}{c} \downarrow \\ \bullet \end{array} & (\xi_\sigma^-)_7 &= \begin{array}{c} \downarrow \\ \cup \end{array} \end{aligned}$$

4.3.5 Chain maps $\sigma(\xi^+)$ and $\sigma(\widehat{\xi}^+)$ in indecomposable case

 For $n \geq 0$ the chain maps

$$\widehat{\xi}_\sigma^+ \mathbf{1}_n := \sigma(\widehat{\xi}^+ \mathbf{1}_{-n-2}) : \sigma(\mathcal{C}\mathbf{1}_{-n-2})\mathcal{E}\mathbf{1}_n \rightarrow \mathcal{E}\sigma(\mathcal{C}\mathbf{1}_{-n}) \quad (4.56)$$

$$\xi_\sigma^+ \mathbf{1}_n := \sigma(\xi^+ \mathbf{1}_{-n-2}) : \mathcal{E}\sigma(\mathcal{C}\mathbf{1}_{-n}) \rightarrow \sigma(\mathcal{C}\mathbf{1}_{-n-2})\mathcal{E}\mathbf{1}_n \quad (4.57)$$

simplify to the form

where

$$(\widehat{\xi}_\sigma^+)_1 = (\widehat{\xi}_\sigma^+)_4 = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \quad (\widehat{\xi}_\sigma^+)_2 = \begin{array}{c} \uparrow \\ \downarrow \end{array} \cup \quad (\widehat{\xi}_\sigma^+)_3 = - \begin{array}{c} \uparrow \\ \downarrow \end{array} \bullet$$

$$(\widehat{\xi}_\sigma^+)_5 = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \quad (\widehat{\xi}_\sigma^+)_6 = (\widehat{\xi}_\sigma^+)_8 = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \quad (\widehat{\xi}_\sigma^+)_7 = - \begin{array}{c} \uparrow \\ \downarrow \end{array} \cup \begin{array}{c} \uparrow \\ \downarrow \end{array}$$

$$(\xi_\sigma^+)_1 = (\xi_\sigma^+)_3 = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \quad (\xi_\sigma^+)_2 = \begin{array}{c} \uparrow \\ \downarrow \end{array} \cup \begin{array}{c} \uparrow \\ \downarrow \end{array} \quad (\xi_\sigma^+)_4 = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array}$$

$$(\xi_\sigma^+)_5 = (\xi_\sigma^+)_8 = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} - \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \quad (\xi_\sigma^+)_6 = - \begin{array}{c} \uparrow \\ \downarrow \end{array} \bullet \quad (\xi_\sigma^+)_7 = - \begin{array}{c} \uparrow \\ \downarrow \end{array} \cup \begin{array}{c} \uparrow \\ \downarrow \end{array}$$

4.4 Symmetries of the commutativity chain maps

Given a 2-morphism α and a 1-morphism x in \mathcal{U} we write αx in place of the composite $\alpha 1_x$ whenever this composite makes sense, likewise for $x\alpha$.

The following Propositions gives several alternative characterizations of the various commutativity chain maps.

Proposition 4.9. For all $n \in \mathbb{Z}$ the equalities in $Kom(\mathcal{U})$

$$\widehat{\xi}^- \mathbf{1}_n = \mathcal{F} \widehat{\varrho}^\psi \circ \psi(\xi^- \mathbf{1}_n) \circ \varrho^\psi \mathcal{F} \quad (4.59)$$

$$\xi^- \mathbf{1}_n = \widehat{\varrho}^\psi \mathcal{F} \circ \psi(\widehat{\xi}^- \mathbf{1}_n) \circ \mathcal{F} \varrho^\psi \quad (4.60)$$

$$\xi^+ \mathbf{1}_n = \mathcal{E} \widehat{\varrho}^\psi \circ \psi(\widehat{\xi}^+ \mathbf{1}_n) \circ \varrho^\psi \mathcal{E} \quad (4.61)$$

$$\widehat{\xi}^+ \mathbf{1}_n = \widehat{\varrho}^\psi \mathcal{E} \circ \psi(\xi^+ \mathbf{1}_n) \circ \mathcal{E} \varrho^\psi \quad (4.62)$$

hold.

Proof. Note that (4.59) is how the map $\widehat{\xi}^-$ was defined. Equation (4.60) follows from the strictly commutative square

$$\begin{array}{ccccc}
 \mathcal{F}\mathcal{C}\mathbf{1}_n & \xrightarrow{\xi^-} & \mathcal{C}\mathcal{F}\mathbf{1}_n & & \\
 \mathcal{F}\varrho^\psi \downarrow & \searrow -\text{Id} & & \nearrow -\text{Id} & \\
 & \mathcal{F}\mathcal{C}\mathbf{1}_n & \xrightarrow{\xi^-} & \mathcal{C}\mathcal{F}\mathbf{1}_n & \\
 \mathcal{F}\psi(\widehat{\varrho}^\psi) \nearrow & & & & \psi(\varrho^\psi)\mathcal{F} \searrow \\
 \mathcal{F}\psi(\mathcal{C})\mathbf{1}_n & \xrightarrow{\psi(\widehat{\xi}^-)} & \psi(\mathcal{C})\mathcal{F}\mathbf{1}_n & & \\
 & & & & \widehat{\varrho}^\psi \mathcal{F} \uparrow
 \end{array} \quad (4.63)$$

where the triangles commute since $\psi(\varrho^\psi) = -\widehat{\varrho}^\psi$, $\psi(\widehat{\varrho}^\psi) = -\varrho^\psi$, and ϱ^ψ has inverse $\widehat{\varrho}^\psi$ by Proposition 4.1. The bottom square is ψ applied to (4.59).

To prove (4.61) consider the diagram:

$$\begin{array}{ccccc}
 \mathcal{C}\mathcal{E}\mathbf{1}_n & \xrightarrow{\xi^+} & \mathcal{E}\mathcal{C}\mathbf{1}_n & & \\
 \varrho^\psi \mathcal{E} \downarrow & \searrow \varrho^{\sigma\omega} \mathcal{E} & & \nearrow \mathcal{E} \widehat{\varrho}^{\sigma\omega} & \\
 & \sigma\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\sigma\omega(\xi^-)} & \mathcal{E}\sigma\omega(\mathcal{C})\mathbf{1}_n & \\
 \sigma\omega(\mathcal{F}\varrho^\psi) \downarrow & & & & \sigma\omega(\widehat{\varrho}^\psi \mathcal{F}) \uparrow \\
 \psi\sigma\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\sigma\omega\psi(\widehat{\xi}^-)} & \mathcal{E}\psi\sigma\omega(\mathcal{C})\mathbf{1}_n & & \\
 -\psi(\widehat{\varrho}^{\sigma\omega})\mathcal{E} \nearrow & & & & -\mathcal{E}\psi(\varrho^{\sigma\omega}) \searrow \\
 \psi(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\psi(\widehat{\xi}^+)} & \mathcal{E}\psi(\mathcal{C})\mathbf{1}_n & & \\
 & & & & \mathcal{E}\widehat{\varrho}^\psi \uparrow
 \end{array} \quad (4.64)$$

The bottom square commutes on the nose since it is ψ applied to the definition of $\widehat{\xi}^+$ in (4.43) with two minus signs distributed through the map. The top square is the definition of ξ^+ in (4.42). The center square is $\sigma\omega$ applied to (4.59). Noting that

$$\sigma\omega(\mathcal{F}\varrho^\psi) = \sigma\omega(\varrho^\psi)\mathcal{E} \quad (4.65)$$

$$\sigma\omega(\widehat{\varrho}^\psi\mathcal{F}) = \sigma\omega(\widehat{\varrho}^\psi)\mathcal{E} \quad (4.66)$$

the left and right squares commute since

$$\psi(\widehat{\varrho}^{\sigma\omega}) \circ \varrho^\psi = -\sigma\omega(\varrho^\psi) \circ \varrho^{\sigma\omega}, \quad \psi(\varrho^{\sigma\omega}) \circ \widehat{\varrho}^\psi = -\sigma\omega(\widehat{\varrho}^\psi) \circ \widehat{\varrho}^{\sigma\omega} \quad (4.67)$$

by Proposition 4.1 part (b). Equation (4.62) follows from (4.61) by applying ψ and arguing as in the proof that (4.59) implies (4.60) above. \square

Just as the maps ξ^+ and $\widehat{\xi}^+$ are defined from ξ^- and $\widehat{\xi}^-$ using the symmetry $\sigma\omega$, the following proposition characterizes ξ^- and $\widehat{\xi}^-$ in terms of ξ^+ and $\widehat{\xi}^+$ via the symmetry $\sigma\omega$.

Proposition 4.10. For all $n \in \mathbb{Z}$ the equalities in $Kom(\mathcal{U})$

$$\xi^- \mathbf{1}_n = \widehat{\varrho}^{\sigma\omega} \mathcal{F} \circ \omega \sigma(\xi^+ \mathbf{1}_n) \circ \mathcal{F} \varrho^{\sigma\omega} \quad (4.68)$$

$$\widehat{\xi}^- \mathbf{1}_n = \mathcal{F} \widehat{\varrho}^{\sigma\omega} \circ \omega \sigma(\widehat{\xi}^+ \mathbf{1}_n) \circ \varrho^{\sigma\omega} \mathcal{F} \quad (4.69)$$

hold.

Proof. The first equation follows from the commutative square

$$\begin{array}{ccc} \mathcal{F}\mathcal{C}\mathbf{1}_n & \xrightarrow{\xi^-} & \mathcal{C}\mathcal{F}\mathbf{1}_n \\ \mathcal{F}\varrho^{\sigma\omega} \downarrow & \text{Id} \searrow & \text{Id} \nearrow \\ & \mathcal{F}\mathcal{C}\mathbf{1}_n & \xrightarrow{\xi^-} \mathcal{C}\mathcal{F}\mathbf{1}_n \\ \sigma\omega(\varrho^{\sigma\omega}\mathcal{E}) = \mathcal{F}\sigma\omega(\varrho^{\sigma\omega}) \nearrow & & \searrow \sigma\omega(\mathcal{E}\widehat{\varrho}^{\sigma\omega}) = \sigma\omega(\widehat{\varrho}^{\sigma\omega}\mathcal{F}) \\ \mathcal{F}\sigma\omega(\mathcal{C})\mathbf{1}_n & \xrightarrow{\sigma\omega(\xi^+)} & \sigma\omega(\mathcal{C})\mathcal{F}\mathbf{1}_n \end{array} \quad (4.70)$$

where the bottom square commutes on the nose since it is $\sigma\omega$ applied to the definition of ξ^+ . The left and right triangles commute on the nose since $\sigma\omega(\varrho^{\sigma\omega}) = \widehat{\varrho}^{\sigma\omega}$ and $\sigma\omega(\widehat{\varrho}^{\sigma\omega}) = \varrho^{\sigma\omega}$ and $\varrho^{\sigma\omega}$ has inverse $\widehat{\varrho}^{\sigma\omega}$ by Proposition 4.1. The second claim in the Proposition is proven similarly using the definition of $\widehat{\xi}^+$. \square

Notice that there are four equations in Proposition 4.9 but only two in Proposition 4.10. The missing equalities are definitions (4.42) and (4.43).

Proposition 4.11. For all $n \in \mathbb{Z}$ the equalities (=) and homotopy equivalences (\simeq) hold.

$$\xi^- \mathbf{1}_n = \widehat{\varrho}^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho^\sigma \quad (4.71)$$

$$\widehat{\xi}^- \mathbf{1}_n \simeq \mathcal{F} \widehat{\varrho}^\sigma \circ \sigma(\xi^- \mathbf{1}_{-n+2}) \circ \varrho^\sigma \mathcal{F} \quad (4.72)$$

$$\xi^+ \mathbf{1}_n = \mathcal{E} \widehat{\varrho}^\sigma \circ \sigma(\widehat{\xi}^+ \mathbf{1}_{-n-2}) \circ \varrho^\sigma \mathcal{E} \quad (4.73)$$

$$\widehat{\xi}^+ \mathbf{1}_n \simeq \widehat{\varrho}^\sigma \mathcal{E} \circ \sigma(\xi^+ \mathbf{1}_{-n-2}) \circ \mathcal{E} \varrho^\sigma \quad (4.74)$$

Proof. We need to show the equality of two chain maps in equation (4.71). Each is given by three two-by-two matrices and we check the equality of coefficients one by one. That is twelve equations to check. Here we prove just one of the more complicated equalities of matrix entries. For example, the equality for the upper left term of the first matrix is

$$\begin{aligned} \xi_1^- \mathbf{1}_n &= \widehat{\varrho}_1^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}_1^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho_1^\sigma + \widehat{\varrho}_3^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}_2^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho_1^\sigma \\ &\quad + \widehat{\varrho}_1^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}_3^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho_2^\sigma + \widehat{\varrho}_3^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}_4^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho_2^\sigma. \end{aligned} \quad (4.75)$$

One can check that

$$\widehat{\varrho}_1^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}_1^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho_1^\sigma = \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =-n+1 \\ 1 \leq f_1}} \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \quad (4.76)$$

$$= \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} - \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \quad (4.77)$$

where the summation term is zero by the dotted curl relation (3.17) since

$$\sum_{\substack{f_1+f_2+f_3 \\ =-n+1 \\ 1 \leq f_1}} \begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array} n = \sum_{\substack{f_1+f_2+f_3 \\ =-n+1 \\ 1 \leq f_1}} \sum_{\substack{g_1+g_2 \\ =n-2+f_3}} \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \end{array} n \quad (4.78)$$

and the for both summations to be nonzero we must have $-n+1 \geq 0$ and $f_3+n-2 \geq 0$ which is impossible since $f_3 \leq -n+1$. Similarly,

$$\widehat{\varrho}_3^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}_2^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho_1^\sigma = - \sum_{\substack{f_1+f_2 \\ =-n+1 \\ 1 \leq f_1}} \sum_{\substack{g_1+g_2 \\ =n-3}} \begin{array}{c} \text{Diagram 13} \\ \text{Diagram 14} \end{array} n = 0 \quad (4.79)$$

since the summation indices can never both be non-negative.

We also have

$$\widehat{\varrho}_1^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}_3^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho_2^\sigma = \begin{array}{c} \text{Diagram 15} \\ \text{Diagram 16} \end{array} n = \sum_{\substack{g_1+g_2 \\ =n-2}} \begin{array}{c} \text{Diagram 17} \\ \text{Diagram 18} \end{array} n \quad (4.80)$$

and

$$\widehat{\varrho}_3^\sigma \mathcal{F} \circ \sigma(\widehat{\xi}_4^- \mathbf{1}_{-n+2}) \circ \mathcal{F} \varrho_2^\sigma = - \sum_{\substack{g_1+g_2 \\ =n-3}} \begin{array}{c} \text{Diagram 19} \\ \text{Diagram 20} \end{array} n \quad (4.81)$$

Hence, equation (4.75) amounts to proving the equality

$$\xi_1^- \mathbf{1}_n = \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array}^n - \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array}^n + \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array}^n - \sum_{\substack{g_1+g_2 \\ =n-3}} \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array}^n \quad (4.82)$$

Using the nilHecke relation (3.4) and cancelling terms we can write

$$\begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array}^n - \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \end{array}^n = \begin{array}{c} \text{Diagram 13} \\ \text{Diagram 14} \end{array}^n - \begin{array}{c} \text{Diagram 15} \\ \text{Diagram 16} \end{array}^n \quad (4.83)$$

Now we can apply (3.15) to both terms on the right-hand side, sliding the bottom right crossing in each diagram through the diagonal line. Note that after applying (3.15) the location of the dots above produce some terms that cancel leaving only

$$\begin{array}{c} \text{Diagram 17} \\ \text{Diagram 18} \end{array}^n - \begin{array}{c} \text{Diagram 19} \\ \text{Diagram 20} \end{array}^n + \sum_{\substack{g_1+g_2+g_3 \\ +g_4=-n}} \begin{array}{c} \text{Diagram 21} \\ \text{Diagram 22} \end{array}^{g_1+1} \downarrow^{g_4} \begin{array}{c} \text{Diagram 23} \\ \text{Diagram 24} \end{array}^n - \sum_{\substack{g_1+g_2+g_3 \\ +g_4=-n}} \begin{array}{c} \text{Diagram 25} \\ \text{Diagram 26} \end{array}^{g_1} \downarrow^{g_4+1} \begin{array}{c} \text{Diagram 27} \\ \text{Diagram 28} \end{array}^n \quad (4.84)$$

where the last two terms can be shown to be zero by simplifying the dotted curl and arguing as above. Therefore,

$$\begin{array}{c} \text{Diagram 29} \\ \text{Diagram 30} \end{array}^n - \begin{array}{c} \text{Diagram 31} \\ \text{Diagram 32} \end{array}^n = \begin{array}{c} \text{Diagram 33} \\ \text{Diagram 34} \end{array}^n - \begin{array}{c} \text{Diagram 35} \\ \text{Diagram 36} \end{array}^n \quad (4.85)$$

$$\stackrel{(3.4)}{=} \begin{array}{c} \text{Diagram 37} \\ \text{Diagram 38} \end{array}^n - \begin{array}{c} \text{Diagram 39} \\ \text{Diagram 40} \end{array}^n + \begin{array}{c} \text{Diagram 41} \\ \text{Diagram 42} \end{array}^n \quad (4.86)$$

$$\stackrel{(3.12)}{=} - \begin{array}{c} \text{Diagram 43} \\ \text{Diagram 44} \end{array} + \begin{array}{c} \text{Diagram 45} \\ \text{Diagram 46} \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-3}} \begin{array}{c} \text{Diagram 47} \\ \text{Diagram 48} \end{array}^{f_1} - \sum_{\substack{f_1+f_2+f_3 \\ =n-3}} \begin{array}{c} \text{Diagram 49} \\ \text{Diagram 50} \end{array}^{f_1} + \begin{array}{c} \text{Diagram 51} \\ \text{Diagram 52} \end{array} \quad (4.87)$$

Now plug this into the right-hand side of (4.82), use the identity decomposition equation (3.12) on the last three terms above and note that the additional bubble terms arising from the application of (3.12) vanish by considering the conditions on the summation indices as above. After reordering the non-vanishing terms of the right-hand side of (4.82) we are

left with

$$- \begin{array}{c} \uparrow \\ \text{---} \\ \downarrow \end{array} + \begin{array}{c} \uparrow \\ \text{---} \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-3}} \begin{array}{c} \curvearrowright f_1 \\ \bullet + f_2 \\ \downarrow \\ \curvearrowleft f_{3+1} \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-3}} \begin{array}{c} \curvearrowright f_1 \\ \bullet + f_2 \\ \downarrow \\ \curvearrowleft f_3 \end{array} - \sum_{\substack{g_1+g_2 \\ =n-3}} \begin{array}{c} \curvearrowright g_1 \\ \bullet + g_2 \\ \downarrow \\ \curvearrowleft \end{array} \quad (4.88)$$

By recalling the third term and simplifying the last two terms, the above becomes

$$- \begin{array}{c} \uparrow \\ \text{---} \\ \downarrow \end{array} + \begin{array}{c} \uparrow \\ \text{---} \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f'_3 \\ =n-2 \\ 1 \leq f'_3}} \begin{array}{c} \curvearrowright f_1 \\ \bullet + f_2 \\ \downarrow \\ \curvearrowleft f'_3 \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-3 \\ 1 \leq f_3}} \begin{array}{c} \curvearrowright f_1 \\ \bullet + f_2 \\ \downarrow \\ \curvearrowleft f_3 \end{array} \quad (4.89)$$

But by sliding the bubbles using a rotated version of (3.21) we can simplify the last two terms

$$\begin{aligned} & \sum_{\substack{f_1+f_2+f_3 \\ =n-3 \\ 1 \leq f_3}} \begin{array}{c} \curvearrowright f_1 \\ \bullet + f_2 \\ \downarrow \\ \curvearrowleft f_3 \end{array} - \sum_{\substack{f_1+f_2+f'_3 \\ =n-2 \\ 1 \leq f'_3}} \begin{array}{c} \curvearrowright f_1 \\ \bullet + f_2 \\ \downarrow \\ \curvearrowleft f'_3 \end{array} \stackrel{(3.21)}{=} \sum_{\substack{f_1+f_2+f_3 \\ =n-3 \\ 1 \leq f_3}} \sum_{\substack{g_1+g_2 \\ =f_2}} (f_2 + 1 - g_2) \begin{array}{c} \curvearrowright f_1 \\ \bullet + g_2 \\ \downarrow \\ \curvearrowleft f_3 \end{array} \\ & - \sum_{\substack{f_1+f_2+f'_3 \\ =n-2 \\ 1 \leq f'_3}} \sum_{\substack{g_1+g_2 \\ =f_2}} (f_2 + 1 - g_2) \begin{array}{c} \curvearrowright f_1 \\ \bullet + g_2 \\ \downarrow \\ \curvearrowleft f'_3 \end{array} \\ & = - \sum_{\substack{f_1+f_2+f_3+f_4 \\ =n-2 \\ 1 \leq f_3}} \begin{array}{c} \curvearrowright f_1 \\ \bullet + f_4 \\ \downarrow \\ \curvearrowleft f_3 \end{array} \quad (4.90) \end{aligned}$$

proving equation (4.75). The rest of the proof of (4.71) follows by many more computations analogous to the one above.

Equations (4.72)–(4.74) follow by applying various symmetries to (4.71). The left and right hand sides of (4.72) constitute the perimeter of the following diagram

$$\begin{array}{ccc} \mathcal{CF}1_n & \xrightarrow{\widehat{\xi}^{-1}_n} & \mathcal{FC}1_n \\ \downarrow \varrho^\sigma \mathcal{F} & \searrow \text{Id} & \nearrow \text{Id} \\ & \mathcal{CF}1_n & \xrightarrow{\widehat{\xi}^{-1}_n} & \mathcal{FC}1_n \\ & \nearrow \sigma(\mathcal{F}\varrho^\sigma) = \sigma(\varrho^\sigma)\mathcal{F} & \searrow \sigma(\widehat{\varrho}^\sigma \mathcal{F}) = \mathcal{F}\sigma(\widehat{\varrho}^\sigma) & \\ \sigma(\mathcal{C})\mathcal{F}1_n & \xrightarrow{\sigma(\xi^{-1}_{-n+2})} & \mathcal{F}\sigma(\mathcal{C})1_n & \end{array} \quad (4.91)$$

Since $\sigma(\varrho^\sigma) = \widehat{\varrho}^\sigma$ and $\sigma(\widehat{\varrho}^\sigma) = \varrho^\sigma$ the left and right triangles commute up to homotopy by Proposition 4.1. The bottom square is σ applied to (4.71) with parameter $-n + 2$.

Equation (4.73) follows from the commutativity of the diagram

$$\begin{array}{ccc}
\mathcal{C}\mathcal{E}\mathbf{1}_n & \xrightarrow{\xi^+} & \mathcal{E}\mathcal{C}\mathbf{1}_n \\
\downarrow \varrho^\sigma \mathcal{E} & \searrow \varrho^{\sigma\omega} \mathcal{E} & \nearrow \mathcal{E} \widehat{\varrho}^{\sigma\omega} \\
& \sigma\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\sigma\omega(\xi^-)} & \mathcal{E}\sigma\omega(\mathcal{C})\mathbf{1}_n & \\
& \uparrow \omega(\varrho^\sigma \mathcal{F}) & & \downarrow \omega(\mathcal{F}\widehat{\varrho}^\sigma) & \\
& \omega(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\omega(\widehat{\xi}^-)} & \mathcal{E}\omega(\mathcal{C})\mathbf{1}_n & \\
\downarrow \sigma(\varrho^{\sigma\omega})\mathcal{E} = \sigma(\varrho^{\sigma\omega})\mathcal{E} & & & \downarrow \mathcal{E}\sigma(\widehat{\varrho}^{\sigma\omega}) = \mathcal{E}\sigma(\widehat{\varrho}^{\sigma\omega}) & \\
\sigma(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\sigma(\widehat{\xi}^+)} & \mathcal{E}\sigma(\mathcal{C})\mathbf{1}_n & \uparrow \mathcal{E}\widehat{\varrho}^\sigma & \\
& & & & \mathcal{E}\mathcal{C}\mathbf{1}_n
\end{array} \tag{4.92}$$

Here $\omega(\varrho^\sigma \mathcal{F}) = \omega(\varrho^\sigma)\mathcal{E}$ and $\omega(\mathcal{F}\widehat{\varrho}^\sigma) = \mathcal{E}\omega(\widehat{\varrho}^\sigma)$, the left and right squares commute on the nose since

$$\varrho^{\omega, \sigma\omega} \circ \sigma(\varrho^{\sigma\omega}) \circ \varrho^\sigma = \varrho^{\sigma\omega}, \quad \widehat{\varrho}^\omega \circ \sigma(\widehat{\varrho}^{\sigma\omega}) \circ \widehat{\varrho}^{\omega, \sigma\omega} = \widehat{\varrho}^{\sigma\omega} \tag{4.93}$$

by Proposition 4.1 part (b). The center square commutes since it is ω applied to (4.71). The bottom square is σ applied to the definition of $\widehat{\xi}^+$, where we used that $\sigma^2 = \text{Id}$. The top square commutes by definition of ξ^+ .

A similar homotopy commutative square to (4.91) shows that (4.74) follows from (4.73). \square

Proposition 4.12. For all $n \in \mathbb{Z}$ the equalities ($=$) and homotopy equivalences (\simeq) hold.

$$\xi^- \mathbf{1}_n = \widehat{\varrho}^\omega \mathcal{F} \circ \omega(\widehat{\xi}^+ \mathbf{1}_{-n}) \circ \mathcal{F} \varrho^\omega \tag{4.94}$$

$$\widehat{\xi}^- \mathbf{1}_n \simeq \mathcal{F} \widehat{\varrho}^\omega \circ \omega(\xi^+ \mathbf{1}_{-n}) \circ \varrho^\omega \mathcal{F} \tag{4.95}$$

$$\xi^+ \mathbf{1}_n = \mathcal{E} \widehat{\varrho}^\omega \circ \omega(\widehat{\xi}^- \mathbf{1}_{-n}) \circ \varrho^\omega \mathcal{E} \tag{4.96}$$

$$\widehat{\xi}^+ \mathbf{1}_n \simeq \widehat{\varrho}^\omega \mathcal{E} \circ \omega(\xi^- \mathbf{1}_{-n}) \circ \mathcal{E} \varrho^\omega \tag{4.97}$$

Proof. The proof follows from Proposition 4.11 and the definitions of ξ^+ and $\widehat{\xi}^+$. For example, the first two equations are proven by the diagrams

$$\begin{array}{ccc}
\mathcal{C}\mathcal{F}\mathbf{1}_n & \xrightarrow{\xi^- \mathbf{1}_n} & \mathcal{F}\mathcal{C}\mathbf{1}_n \\
\downarrow \varrho^\omega \mathcal{F} & \searrow \varrho^\sigma \mathcal{F} & \nearrow \mathcal{F} \widehat{\varrho}^\sigma \\
& \sigma(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\sigma(\widehat{\xi}^- \mathbf{1}_{-n+2})} & \sigma(\mathcal{C})\mathcal{F}\mathbf{1}_n & \\
& \uparrow \omega(\varrho^{\sigma\omega})\mathcal{F} & & \downarrow \mathcal{F} \widehat{\varrho}^\omega & \\
& \omega(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\omega(\widehat{\xi}^+ \mathbf{1}_{-n})} & \mathcal{F}\omega(\mathcal{C})\mathbf{1}_n & \\
& & & & \mathcal{F}\mathcal{C}\mathbf{1}_n
\end{array} \quad
\begin{array}{ccc}
\mathcal{F}\mathcal{C}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^- \mathbf{1}_n} & \mathcal{C}\mathcal{F}\mathbf{1}_n \\
\downarrow \mathcal{F} \varrho^\omega & \searrow \mathcal{F} \varrho^\sigma & \nearrow \widehat{\varrho}^\sigma \mathcal{F} \\
& \mathcal{F}\sigma(\mathcal{C})\mathbf{1}_n & \xrightarrow{\sigma(\xi^- \mathbf{1}_{-n+2})} & \sigma(\mathcal{C})\mathcal{F}\mathbf{1}_n & \\
& \uparrow \mathcal{F}\omega(\varrho^{\sigma\omega}) & & \downarrow \omega(\widehat{\varrho}^{\sigma\omega})\mathcal{F} & \\
& \mathcal{F}\omega(\mathcal{C})\mathbf{1}_n & \xrightarrow{\omega(\xi^+ \mathbf{1}_{-n})} & \omega(\mathcal{C})\mathcal{F}\mathbf{1}_n & \\
& & & & \mathcal{C}\mathcal{F}\mathbf{1}_n
\end{array} \tag{4.98}$$

The top squares commute by Proposition 4.11. The bottom squares are ω applied to the definitions of $\widehat{\xi}^+$ and ξ^+ , respectively. Note that $\omega(\varrho^{\sigma\omega}) = \sigma(\widehat{\varrho}^{\sigma\omega})$ and $\omega(\widehat{\varrho}^{\sigma\omega}) = \sigma(\varrho^{\sigma\omega})$. The left and right triangles in both squares commute on the nose since

$$\sigma(\widehat{\varrho}^{\sigma\omega}) \circ \varrho^\omega = \varrho^\sigma, \quad \widehat{\varrho}^\omega \circ \sigma(\varrho^{\sigma\omega}) = \widehat{\varrho}^\sigma. \quad (4.99)$$

Using Proposition 4.10, similar arguments as the above prove the last two equations in the Proposition. \square

5 Naturality of the Casimir complex

In the previous section we have shown that the Casimir complex commutes with generating 1-morphisms $\mathcal{E}\mathbf{1}_n, \mathcal{F}\mathbf{1}_n$ in $Com(\mathcal{U})$. In this section we show that this commutativity is natural with respect to 2-morphisms.

5.1 Natural transformations κ and $\widehat{\kappa}$

Throughout this section we will find it convenient to view $Kom(\mathcal{U})$ and $Com(\mathcal{U})$ as idempotent additive monoidal categories as explained in the introduction. Consider the complex

$$\mathcal{C} := \bigoplus_{n \in \mathbb{Z}} \mathcal{C}\mathbf{1}_n. \quad (5.1)$$

In this section we show that the functor

$$- \otimes \mathcal{C}: Com(\mathcal{U}) \rightarrow Com(\mathcal{U}), \quad (5.2)$$

is naturally isomorphic to the functor

$$\mathcal{C} \otimes -: Com(\mathcal{U}) \rightarrow Com(\mathcal{U}) \quad (5.3)$$

via an invertible natural transformation

$$\kappa: - \otimes \mathcal{C} \Rightarrow \mathcal{C} \otimes - \quad (5.4)$$

with inverse

$$\widehat{\kappa}: \mathcal{C} \otimes - \Rightarrow - \otimes \mathcal{C}. \quad (5.5)$$

Recall that the tensor product of complexes and juxtaposition of diagrams gives the composition operation in categories $Com(\mathcal{U})$ and $Com(\dot{\mathcal{U}})$. Here we will use composition notation rather than the tensor notation.

Defining the natural transformation κ and its inverse $\widehat{\kappa}$ amounts to specifying for any complex X in $Com(\mathcal{U})$ a chain map

$$\kappa_X: XC \rightarrow \mathcal{C}X, \quad \widehat{\kappa}_X: \mathcal{C}X \rightarrow XC, \quad (5.6)$$

such that for any chain map $f: X \rightarrow Y$ the squares

$$\begin{array}{ccc}
 Y\mathcal{C} & \xrightarrow{\kappa_Y} & \mathcal{C}Y \\
 \uparrow f\mathcal{C} & & \uparrow cf \\
 X\mathcal{C} & \xrightarrow{\kappa_X} & \mathcal{C}X
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{C}Y & \xrightarrow{\widehat{\kappa}_Y} & Y\mathcal{C} \\
 \uparrow cf & & \uparrow f\mathcal{C} \\
 \mathcal{C}X & \xrightarrow{\widehat{\kappa}_X} & X\mathcal{C}
 \end{array}
 \tag{5.7}$$

commute (up to chain homotopy).

On generating 1-morphisms we define

$$\kappa_{\mathcal{F}\mathbf{1}_n} := \xi^- : \mathcal{F}\mathcal{C}\mathbf{1}_n \rightarrow \mathcal{C}\mathcal{F}\mathbf{1}_n \quad \widehat{\kappa}_{\mathcal{F}\mathbf{1}_n} := \widehat{\xi}^- : \mathcal{C}\mathcal{F}\mathbf{1}_n \rightarrow \mathcal{F}\mathcal{C}\mathbf{1}_n \tag{5.8}$$

$$\kappa_{\mathcal{E}\mathbf{1}_n} := \widehat{\xi}^+ : \mathcal{E}\mathcal{C}\mathbf{1}_n \rightarrow \mathcal{C}\mathcal{E}\mathbf{1}_n \quad \widehat{\kappa}_{\mathcal{E}\mathbf{1}_n} := \xi^+ : \mathcal{C}\mathcal{E}\mathbf{1}_n \rightarrow \mathcal{E}\mathcal{C}\mathbf{1}_n. \tag{5.9}$$

For an arbitrary complex X in $Com(\mathcal{U})$ the chain maps κ_X and $\widehat{\kappa}_X$ are determined from the assignments above. For example, if $X = \mathcal{E}\mathcal{F}\mathcal{F}\mathbf{1}_n\{t\}$ then

$$\kappa_X = \mathcal{E}\mathcal{F}\mathcal{F}\mathcal{C}\mathbf{1}_n\{t\} \xrightarrow{\mathcal{E}\mathcal{F}\xi^-\{t\}} \mathcal{E}\mathcal{F}\mathcal{C}\mathcal{F}\mathbf{1}_n\{t\} \xrightarrow{\mathcal{E}\xi^-\mathcal{F}\{t\}} \mathcal{E}\mathcal{C}\mathcal{F}\mathcal{F}\mathbf{1}_n\{t\} \xrightarrow{\widehat{\xi}^+\mathcal{F}\mathcal{F}\{t\}} \mathcal{C}\mathcal{E}\mathcal{F}\mathcal{F}\mathbf{1}_n\{t\}. \tag{5.10}$$

This definition of κ_X produces a commutative diagram

$$\begin{array}{ccc}
 \mathcal{C}XY\mathbf{1}_n & \xrightarrow{\kappa_{XY}} & XY\mathcal{C}\mathbf{1}_n \\
 \searrow \kappa_{XY} & & \nearrow X\kappa_Y \\
 & XCY\mathbf{1}_n &
 \end{array}
 \tag{5.11}$$

in $Com(\mathcal{U})$ for complexes $Y = \mathbf{1}_{n'}Y\mathbf{1}_n$ and $X = \mathbf{1}_{n''}X\mathbf{1}_{n'}$, with $\mathcal{C}XY\mathbf{1}_n = \mathbf{1}_{n''}\mathcal{C}\mathbf{1}_{n''}X\mathbf{1}_{n'}Y\mathbf{1}_n$.

Proposition 5.1. Equations (5.7) hold for all 2-morphisms in $Com(\mathcal{U})$.

Proof. It is enough to check naturality squares (5.7) on generating 2-morphisms (dots, crossing, cups, and caps) in \mathcal{U} . This will be done in Section 5.2. \square

It is clear from the definitions in (5.8) and (5.9) of κ and $\widehat{\kappa}$ and the results in section 4.2 that

$$\widehat{\kappa}_X\kappa_X = \text{Id}_{\mathcal{C}X}, \quad \kappa_X\widehat{\kappa}_X = \text{Id}_{X\mathcal{C}} \tag{5.12}$$

in $Com(\mathcal{U})$, so that κ and $\widehat{\kappa}$ are inverse.

Naturality of κ and $\widehat{\kappa}$ and the universality of the Karoubian envelope allow us to extend κ and $\widehat{\kappa}$ to isomorphisms between functors

$$- \otimes \mathcal{C} : Kar(Com(\mathcal{U})) \rightarrow Kar(Com(\mathcal{U})) \tag{5.13}$$

and

$$\mathcal{C} \otimes - : Kar(Com(\mathcal{U})) \rightarrow Kar(Com(\mathcal{U})). \tag{5.14}$$

The equivalence $Kar(Com(\mathcal{U})) \cong Com(\dot{\mathcal{U}})$ allows us to treat κ and $\widehat{\kappa}$ as isomorphisms

$$\kappa : - \otimes \mathcal{C} \Rightarrow \mathcal{C} \otimes - \quad \text{and} \quad \widehat{\kappa} : \mathcal{C} \otimes - \Rightarrow - \otimes \mathcal{C}$$

between endofunctors on $Com(\dot{\mathcal{U}})$ concluding the proof of Theorem 1.1.

5.2 Naturality with respect to 2-morphisms

It is immediate from the axioms of a 2-category and will be used throughout this section that for any $g \in \mathcal{G}$ the chain maps $\varrho^g: \mathcal{C} \rightarrow g(\mathcal{C})$ induce natural transformations between functors

$$\varrho^g \otimes -: \mathcal{C} \otimes - \Rightarrow g(\mathcal{C}) \otimes -, \quad - \otimes \varrho^g: - \otimes \mathcal{C} \Rightarrow - \otimes g(\mathcal{C}). \quad (5.15)$$

5.2.1 Naturality of κ for dot 2-morphism

We will show that the diagram

$$\begin{array}{ccc} \mathcal{FC}1_n \xrightarrow{\kappa_{\mathcal{F}1_n}} \mathcal{CF}1_n & & \mathcal{FC}1_n \xrightarrow{\xi^-} \mathcal{CF}1_n \\ \downarrow \mathcal{C} \uparrow & & \downarrow \mathcal{C} \uparrow \\ \mathcal{FC}1_n \xrightarrow{\kappa_{\mathcal{F}1_n}} \mathcal{CF}1_n & := & \mathcal{FC}1_n \xrightarrow{\xi^-} \mathcal{CF}1_n \\ & & \uparrow \mathcal{C} \downarrow \end{array} \quad (5.16)$$

commutes up to homotopy (commutes in $Com(\mathcal{U})$). Let

$$(\xi^-)' = \xi^- \circ (\downarrow \mathcal{C}) - (\mathcal{C} \downarrow) \circ \xi^-. \quad (5.17)$$

We construct a chain homotopy $(\xi^-)' \simeq 0$.

$$\begin{array}{ccccc} \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathcal{F}1_n\{2\} \\ \mathcal{F}1_n\{1+n\} \end{array} \right) & \xrightarrow{\left(\begin{array}{cc} \uparrow \downarrow \downarrow & \downarrow \\ \uparrow \downarrow \downarrow & \downarrow \end{array} \right)} & \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathcal{F}1_n \\ \mathcal{E}\mathcal{F}\mathcal{F}1_n \end{array} \right) & \xrightarrow{\left(\begin{array}{cc} - \uparrow \downarrow \downarrow & \uparrow \downarrow \downarrow \\ \downarrow & - \downarrow \end{array} \right)} & \left(\begin{array}{c} \mathcal{E}\mathcal{F}\mathcal{F}1_n\{-2\} \\ \mathcal{F}1_n\{n-3\} \end{array} \right) \\ \uparrow (\xi'^-)^{-1} & \swarrow h^0 & \uparrow (\xi'^-)^0 & \swarrow h^1 & \uparrow (\xi'^-)^1 \\ \left(\begin{array}{c} \mathcal{F}\mathcal{E}\mathcal{F}1_n\{2\} \\ \mathcal{F}1_n\{1-n\} \end{array} \right) & \xrightarrow{\left(\begin{array}{cc} \downarrow \uparrow \downarrow & \downarrow \\ \downarrow \uparrow \downarrow & \downarrow \end{array} \right)} & \left(\begin{array}{c} \mathcal{F}\mathcal{E}\mathcal{F}1_n \\ \mathcal{F}\mathcal{E}\mathcal{F}1_n \end{array} \right) & \xrightarrow{\left(\begin{array}{cc} - \downarrow \uparrow \downarrow & \downarrow \uparrow \downarrow \\ \downarrow & - \downarrow \end{array} \right)} & \left(\begin{array}{c} \mathcal{F}\mathcal{E}\mathcal{F}1_n\{-2\} \\ \mathcal{F}1_n\{n-1\} \end{array} \right) \end{array} \quad (5.18)$$

where, after simplifying the map $(\xi^-)'$, we have

$$(\xi'^-)^{-1}_{11} = \begin{array}{c} \downarrow \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (\xi'^-)^{-1}_{12} = 0 \quad (5.19)$$

$$(\xi'^-)^{-1}_{21} = \begin{array}{c} \downarrow \\ \downarrow \end{array} - \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (\xi'^-)^{-1}_{22} = 0 \quad (5.20)$$

$$(\xi'^{-})_{11}^0 = \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} - \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \quad (5.21)$$

$$(\xi'^{-})_{12}^0 = 0 \quad (5.22)$$

$$(\xi'^{-})_{21}^0 = \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} - \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (5.23)$$

$$(\xi'^{-})_{22}^0 = \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} - \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (5.24)$$

$$(\xi'^{-})_{11}^1 = \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} - \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} + \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (5.25)$$

$$(\xi'^{-})_{12}^1 = \sum_{\substack{f_1+f_2=n \\ 1 \leq f_1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} - \sum_{\substack{f_1+f_2=n \\ 1 \leq f_1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (5.26)$$

$$(\xi'^{-})_{21}^1 = (\xi'^{-})_{22}^1 = 0 \quad (5.27)$$

and the chain homotopy is given by

$$(h^0)_{11} = -(h^0)_{12} = \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (5.28)$$

$$(h^0)_{21} = -(h^0)_{22} = - \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (5.29)$$

$$(h^1)_{11} = \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (h^1)_{12} = \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} - \sum_{\substack{f_1+f_2+f_3 \\ =n-1 \\ 1 \leq f_1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (5.30)$$

$$(h^1)_{21} = \sum_{\substack{f_1+f_2=n \\ 1 \leq f_1}} \begin{array}{c} \diagdown \quad \downarrow \\ \diagup \quad \downarrow \end{array} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array} \quad (h^1)_{22} = 0 \quad (5.31)$$

The naturality square for the map $\kappa_{\mathcal{E}1_n} = \widehat{\xi}^+$ follows from the naturality square in

(5.16):

$$\begin{array}{ccc}
 \mathcal{E}\mathcal{C}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^+} & \mathcal{C}\mathcal{E}\mathbf{1}_n \\
 \mathcal{E}\varrho^\omega \searrow & & \nearrow \widehat{\varrho}^\omega \mathcal{E} \\
 \mathcal{E}\omega(\mathcal{C})\mathbf{1}_n & \xrightarrow{\omega(\widehat{\xi}^-)} & \omega(\mathcal{C})\mathcal{E}\mathbf{1}_n \\
 \uparrow \bullet c & & \uparrow \bullet c \\
 \mathcal{E}\omega(\mathcal{C})\mathbf{1}_n & \xrightarrow{\omega(\widehat{\xi}^-)} & \omega(\mathcal{C})\mathcal{E}\mathbf{1}_n \\
 \mathcal{E}\varrho^\omega \nearrow & & \searrow \widehat{\varrho}^\omega \mathcal{E} \\
 \mathcal{E}\mathcal{C}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^+} & \mathcal{C}\mathcal{E}\mathbf{1}_n
 \end{array} \quad (5.32)$$

where the middle square is the image of (5.16) under the 2-functor ω . The left and right squares commute by the naturality of ϱ^ω and $\widehat{\varrho}^\omega$, and the top and bottom squares commutes by Proposition 4.12.

5.2.2 Naturality of $\widehat{\kappa}$ for dot 2-morphism

We will show that the diagram

$$\begin{array}{ccc}
 \mathcal{C}\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\kappa}_{\mathcal{F}\mathbf{1}_n}} & \mathcal{F}\mathcal{C}\mathbf{1}_n \\
 \uparrow \bullet c & & \uparrow \bullet c \\
 \mathcal{C}\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\kappa}_{\mathcal{F}\mathbf{1}_n}} & \mathcal{F}\mathcal{C}\mathbf{1}_n
 \end{array} := \begin{array}{ccc}
 \mathcal{C}\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^-} & \mathcal{F}\mathcal{C}\mathbf{1}_n \\
 \uparrow \bullet c & & \uparrow \bullet c \\
 \mathcal{C}\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^-} & \mathcal{F}\mathcal{C}\mathbf{1}_n
 \end{array} \quad (5.33)$$

commutes up to homotopy. To see this apply $\sigma\omega$ to the diagram (5.32), and use the naturality of $\varrho^{\sigma\omega}$ and $\widehat{\varrho}^{\sigma\omega}$, and Proposition 4.10 to see that each of the five small squares in (5.34) commutes.

$$\begin{array}{ccc}
 \mathcal{C}\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^-} & \mathcal{F}\mathcal{C}\mathbf{1}_n \\
 \varrho^{\sigma\omega} \mathcal{F} \searrow & & \nearrow \mathcal{F} \widehat{\varrho}^{\sigma\omega} \\
 \sigma\omega(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\sigma\omega(\widehat{\xi}^+)} & \mathcal{F}\sigma\omega(\mathcal{C})\mathbf{1}_n \\
 \uparrow \bullet \sigma\omega(\mathcal{C}) & & \uparrow \bullet \sigma\omega(\mathcal{C}) \\
 \sigma\omega(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\sigma\omega(\widehat{\xi}^+)} & \mathcal{F}\sigma\omega(\mathcal{C})\mathbf{1}_n \\
 \varrho^{\sigma\omega} \mathcal{F} \nearrow & & \searrow \mathcal{F} \widehat{\varrho}^{\sigma\omega} \\
 \mathcal{C}\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^-} & \mathcal{F}\mathcal{C}\mathbf{1}_n
 \end{array} \quad (5.34)$$

The naturality square

$$\begin{array}{ccc}
 \mathcal{CE}1_n & \xrightarrow{\widehat{\kappa}_{\mathcal{E}1_n}} & \mathcal{E}C1_n \\
 \uparrow \bullet \uparrow c & & \uparrow \bullet \uparrow c \\
 \mathcal{CE}1_n & \xrightarrow{\widehat{\kappa}_{\mathcal{E}1_n}} & \mathcal{E}C1_n
 \end{array}
 :=
 \begin{array}{ccc}
 \mathcal{CE}1_n & \xrightarrow{\xi^+} & \mathcal{E}C1_n \\
 \uparrow \bullet \downarrow c & & \uparrow \bullet \uparrow c \\
 \mathcal{CE}1_n & \xrightarrow{\xi^+} & \mathcal{E}C1_n
 \end{array}
 \quad (5.35)$$

can similarly be shown to commute by applying $\sigma\omega$ to the square (5.16) and appealing to Proposition 4.10.

5.2.3 Naturality of κ and $\widehat{\kappa}$ for crossing 2-morphisms

We will show that the diagram

$$\begin{array}{ccc}
 \mathcal{F}F\mathcal{C}1_n & \xrightarrow{\kappa_{\mathcal{F}F1_n}} & \mathcal{C}\mathcal{F}F1_n \\
 \uparrow \text{crossing } c & & \uparrow c \text{ crossing} \\
 \mathcal{F}F\mathcal{C}1_n & \xrightarrow{\kappa_{\mathcal{F}F1_n}} & \mathcal{C}\mathcal{F}F1_n
 \end{array}
 :=
 \begin{array}{ccc}
 \mathcal{F}F\mathcal{C}1_n & \xrightarrow{\mathcal{F}\xi^-} & \mathcal{F}C\mathcal{F}1_n & \xrightarrow{\xi^- \mathcal{F}} & \mathcal{C}\mathcal{F}F1_n \\
 \uparrow \text{crossing } c & & \uparrow c \text{ crossing} & & \uparrow c \text{ crossing} \\
 \mathcal{F}F\mathcal{C}1_n & \xrightarrow{\mathcal{F}\xi^-} & \mathcal{F}C\mathcal{F}1_n & \xrightarrow{\xi^- \mathcal{F}} & \mathcal{C}\mathcal{F}F1_n
 \end{array}
 \quad (5.36)$$

commutes for all $n \in \mathbb{Z}$ by treating the cases $n \leq 0$ and $n > 0$ separately. For $n \leq 0$ consider the map

$$Z := (\mathcal{C} \text{ crossing } c) \circ (\xi^- \mathcal{F}) \circ (\mathcal{F}\xi^-) - (\xi^- \mathcal{F}) \circ (\mathcal{F}\xi^-) \circ (\text{crossing } c). \quad (5.37)$$

One can check that the map Z is identically zero, so that the diagram (5.36) commutes on the nose for $n \leq 0$ (i.e. commutes in $Kom(\mathcal{U})$).

For $n > 0$ we deduce commutativity from the commutativity of the diagram

$$\begin{array}{ccccccc}
 \mathcal{F}F\mathcal{C}1_n & \xrightarrow{\mathcal{F}\xi^-} & \mathcal{F}F\mathcal{C}1_n & \xrightarrow{\xi^- \mathcal{F}} & \mathcal{C}\mathcal{F}F1_n & & \\
 \uparrow \text{crossing } c & \searrow \mathcal{F}\mathcal{F}\varrho^\sigma & \mathcal{F}\widehat{\varrho}^\sigma \mathcal{F} & \searrow \mathcal{F}\varrho^\sigma \mathcal{F} & \widehat{\varrho}^\sigma \mathcal{F}\mathcal{F} & \uparrow c \text{ crossing} & \\
 \mathcal{F}\mathcal{F}\sigma(\mathcal{C})1_n & \xrightarrow{\mathcal{F}\widehat{\xi}_\sigma^-} & \mathcal{F}\sigma(\mathcal{C})\mathcal{F}1_n & \xrightarrow{\text{Id}} & \mathcal{F}\sigma(\mathcal{C})\mathcal{F}1_n & \xrightarrow{\widehat{\xi}_\sigma^- \mathcal{F}} & \sigma(\mathcal{C})\mathcal{F}F1_n \\
 \uparrow \text{crossing } \sigma(c) & & \uparrow \text{crossing } \sigma(c) & & \uparrow \text{crossing } \sigma(c) & & \\
 \mathcal{F}\mathcal{F}\sigma(\mathcal{C})1_n & \xrightarrow{\mathcal{F}\widehat{\xi}_\sigma^-} & \mathcal{F}\sigma(\mathcal{C})\mathcal{F}1_n & \xrightarrow{\text{Id}} & \mathcal{F}\sigma(\mathcal{C})\mathcal{F}1_n & \xrightarrow{\widehat{\xi}_\sigma^- \mathcal{F}} & \sigma(\mathcal{C})\mathcal{F}F1_n \\
 \uparrow \mathcal{F}\mathcal{F}\varrho^\sigma & & \uparrow \mathcal{F}\widehat{\varrho}^\sigma \mathcal{F} & & \uparrow \mathcal{F}\varrho^\sigma \mathcal{F} & & \uparrow \widehat{\varrho}^\sigma \mathcal{F}\mathcal{F} \\
 \mathcal{F}F\mathcal{C}1_n & \xrightarrow{\mathcal{F}\xi^-} & \mathcal{F}C\mathcal{F}1_n & \xrightarrow{\xi^- \mathcal{F}} & \mathcal{C}\mathcal{F}F1_n & &
 \end{array}
 \quad (5.38)$$

where the center rectangle

$$\begin{array}{ccccc}
 \mathcal{F}\mathcal{F}\sigma(\mathcal{C})\mathbf{1}_n & \xrightarrow{\mathcal{F}\widehat{\xi}_\sigma^-} & \mathcal{F}\sigma(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\xi}_\sigma^- \mathcal{F}} & \sigma(\mathcal{C})\mathcal{F}\mathcal{F}\mathbf{1}_n \\
 \swarrow \times \sigma(\mathcal{C}) \uparrow & & & & \uparrow \sigma(\mathcal{C}) \searrow \times \\
 \mathcal{F}\mathcal{F}\sigma(\mathcal{C})\mathbf{1}_n & \xrightarrow{\mathcal{F}\widehat{\xi}_\sigma^-} & \mathcal{F}\sigma(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\xi}_\sigma^- \mathcal{F}} & \sigma(\mathcal{C})\mathcal{F}\mathcal{F}\mathbf{1}_n
 \end{array} \quad (5.39)$$

commutes on the nose since the map (of complexes)

$$Z' := (\sigma(\mathcal{C}) \searrow \times) \circ (\widehat{\xi}_\sigma^- \mathcal{F}) \circ (\mathcal{F}\widehat{\xi}_\sigma^-) - (\widehat{\xi}_\sigma^- \mathcal{F}) \circ (\mathcal{F}\widehat{\xi}_\sigma^-) \circ (\searrow \times \sigma(\mathcal{C})). \quad (5.40)$$

is already zero in $Kom(\mathcal{U})$ for $n > 0$. The triangles commute up to homotopy since ϱ^σ is the homotopy inverse of $\widehat{\varrho}^\sigma$ by Proposition 4.1. The left and right squares follow from the naturality of ϱ^σ , and the commutativity up to homotopy of the remaining squares is implied by (4.71).

For all $n \in \mathbb{Z}$ the naturality square for $\kappa_{\mathcal{E}\mathcal{E}\mathbf{1}_n}$ and the crossing is proven as follows

$$\begin{array}{ccccc}
 \mathcal{E}\mathcal{E}\mathcal{C}\mathbf{1}_n & \xrightarrow{\mathcal{E}\widehat{\xi}^+} & \mathcal{E}\mathcal{C}\mathcal{E}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^+ \mathcal{E}} & \mathcal{C}\mathcal{E}\mathcal{E}\mathbf{1}_n \\
 \mathcal{E}\mathcal{E}\varrho^\omega \searrow & & \mathcal{E}\widehat{\varrho}^\omega \mathcal{E} \nearrow & & \widehat{\varrho}^\omega \mathcal{E}\mathcal{E} \nearrow \\
 \mathcal{E}\mathcal{E}\omega(\mathcal{C})\mathbf{1}_n & \xrightarrow{\mathcal{E}\omega(\xi^-)} & \mathcal{E}\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\text{Id}} & \mathcal{E}\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\omega(\xi^-)\mathcal{E}} & \omega(\mathcal{C})\mathcal{E}\mathcal{E}\mathbf{1}_n \\
 \swarrow \times c \uparrow & & \swarrow \times \omega(\mathcal{C}) \uparrow & & \omega(\mathcal{C}) \swarrow \times \uparrow & & \swarrow \times c \uparrow \\
 \mathcal{E}\mathcal{E}\omega(\mathcal{C})\mathbf{1}_n & \xrightarrow{\mathcal{E}\omega(\xi^-)} & \mathcal{E}\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\text{Id}} & \mathcal{E}\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\omega(\xi^-)\mathcal{E}} & \omega(\mathcal{C})\mathcal{E}\mathcal{E}\mathbf{1}_n \\
 \mathcal{E}\mathcal{E}\varrho^\omega \nearrow & & \mathcal{E}\widehat{\varrho}^\omega \mathcal{E} \searrow & & \mathcal{E}\varrho^\omega \mathcal{E} \nearrow & & \widehat{\varrho}^\omega \mathcal{E}\mathcal{E} \searrow \\
 \mathcal{E}\mathcal{E}\mathcal{C}\mathbf{1}_n & \xrightarrow{\mathcal{E}\widehat{\xi}^+} & \mathcal{E}\mathcal{C}\mathcal{E}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^+ \mathcal{E}} & \mathcal{C}\mathcal{E}\mathcal{E}\mathbf{1}_n
 \end{array} \quad (5.41)$$

where the center rectangle is the image of (5.36) under ω . Each of the two triangles commutes up to homotopy since ϱ^ω is the homotopy inverse of $\widehat{\varrho}^\omega$. The squares on the left and right commute by the naturality of ϱ^ω and $\widehat{\varrho}^\omega$. The remaining squares commute by equation (4.97) in Proposition 4.12.

The naturality of $\widehat{\kappa}$ and the crossing is established by applying the symmetry 2-functor ψ to the above arguments.

5.2.4 Naturality of κ for cap 2-morphisms

We show that the naturality square

$$\begin{array}{ccc}
 \mathcal{C}\mathbf{1}_n & \xrightarrow{\kappa_{\mathbf{1}_n}} & \mathcal{C}\mathbf{1}_n \\
 \uparrow \wr^c & & \uparrow c \\
 \mathcal{E}\mathcal{F}\mathcal{C}\mathbf{1}_n & \xrightarrow{\kappa_{\mathcal{E}\mathcal{F}\mathbf{1}_n}} & \mathcal{C}\mathcal{E}\mathcal{F}\mathbf{1}_n
 \end{array}
 \quad := \quad
 \begin{array}{ccc}
 \mathcal{C}\mathbf{1}_n & \xrightarrow{\mathcal{C}} & \mathcal{C}\mathbf{1}_n \\
 \uparrow \wr^c & & \uparrow c \\
 \mathcal{E}\mathcal{F}\mathcal{C}\mathbf{1}_n & \xrightarrow{\mathcal{E}\xi^-} \mathcal{E}\mathcal{C}\mathcal{F}\mathbf{1}_n \xrightarrow{\widehat{\xi}^+\mathcal{F}} & \mathcal{C}\mathcal{E}\mathcal{F}\mathbf{1}_n
 \end{array}
 \quad (5.42)$$

commutes up to homotopy by considering the cases $n > 0$ and $n \leq 0$ separately.

For $n \leq 0$ the complex $\mathcal{C}\mathbf{1}_n$ is indecomposable. In this case, let

$$Z = \left(\mathcal{C} \wr \right) \circ \left(\widehat{\xi}^+\mathcal{F} \right) \circ \left(\mathcal{E}\xi^- \right) - \left(\wr \mathcal{C} \right). \quad (5.43)$$

We will specify a homotopy $Z \simeq 0$.

$$\begin{array}{ccccc}
 \begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} \\ \mathbf{1}_n\{1-n\} \end{pmatrix} & \xrightarrow{\begin{pmatrix} \uparrow \downarrow \wr \\ \uparrow \downarrow \wr \end{pmatrix}} & \begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n \\ \mathcal{E}\mathcal{F}\mathbf{1}_n \end{pmatrix} & \xrightarrow{\begin{pmatrix} -\uparrow \downarrow \wr & \uparrow \downarrow \wr \\ \wr & -\wr \end{pmatrix}} & \begin{pmatrix} \mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \\ \mathbf{1}_n\{n-1\} \end{pmatrix} \\
 \uparrow (Z)^{-1} & & \uparrow (Z)^0 & & \uparrow (Z)^1 \\
 & \swarrow h^0 & & \searrow h^1 & \\
 \begin{pmatrix} \mathcal{E}\mathcal{F}\mathcal{E}\mathcal{F}\mathbf{1}_n\{2\} \\ \mathcal{E}\mathcal{F}\mathbf{1}_n\{1-n\} \end{pmatrix} & \xrightarrow{\begin{pmatrix} \uparrow \downarrow \uparrow \downarrow \wr & \uparrow \downarrow \wr \\ \uparrow \downarrow \uparrow \downarrow \wr & \uparrow \downarrow \wr \end{pmatrix}} & \begin{pmatrix} \mathcal{E}\mathcal{F}\mathcal{E}\mathcal{F}\mathbf{1}_n \\ \mathcal{E}\mathcal{F}\mathcal{E}\mathcal{F}\mathbf{1}_n \end{pmatrix} & \xrightarrow{\begin{pmatrix} -\uparrow \downarrow \uparrow \downarrow \wr & \uparrow \downarrow \uparrow \downarrow \wr \\ \uparrow \downarrow \uparrow \downarrow \wr & -\uparrow \downarrow \uparrow \downarrow \wr \end{pmatrix}} & \begin{pmatrix} \mathcal{E}\mathcal{F}\mathcal{E}\mathcal{F}\mathbf{1}_n\{-2\} \\ \mathcal{E}\mathcal{F}\mathbf{1}_n\{n-1\} \end{pmatrix}
 \end{array}
 \quad (5.44)$$

where, after simplifying, Z is given by terms

$$(Z)_{11}^{-1} = \begin{array}{c} \uparrow \downarrow \\ \uparrow \downarrow \end{array} \wr^n - \begin{array}{c} \uparrow \downarrow \\ \uparrow \downarrow \end{array} \wr^n - \begin{array}{c} \uparrow \downarrow \\ \uparrow \downarrow \end{array} \wr^n + \begin{array}{c} \uparrow \downarrow \\ \uparrow \downarrow \end{array} \wr^n \quad (5.45)$$

$$(Z)_{12}^{-1} = 0 \quad (5.46)$$

$$(Z)_{21}^{-1} = - \begin{array}{c} \uparrow \downarrow \\ \uparrow \downarrow \end{array} \wr^n \quad (5.47)$$

$$(Z)_{22}^{-1} = - \begin{array}{c} \uparrow \downarrow \\ \uparrow \downarrow \end{array} \wr^n \quad (5.48)$$

$$(Z)_{11}^0 = \text{diagram} - \text{diagram} - \text{diagram} - \text{diagram} \quad (5.49)$$

$$(Z)_{12}^0 = \text{diagram} - \text{diagram} \quad (5.50)$$

$$(Z)_{21}^0 = 0 \quad (5.51)$$

$$(Z)_{22}^0 = \text{diagram} - \text{diagram} - \text{diagram} \quad (5.52)$$

$$(Z)_{11}^1 = \text{diagram} - \text{diagram} - \text{diagram} - \text{diagram} \quad (5.53)$$

$$(Z)_{12}^1 = \text{diagram} \quad (5.54)$$

$$(Z)_{21}^1 = 0 \quad (5.55)$$

$$(Z)_{22}^1 = - \text{diagram} \quad (5.56)$$

and the chain homotopy is given by

$$(h^0)_{11} = - \text{diagram} - \sum_{\substack{g_1+g_2+g_3 \\ =-n}} g_1 \text{diagram} \quad (5.57)$$

$$(h^0)_{12} = 0 \quad (5.58)$$

$$(h^0)_{21} = \text{diagram} + \sum_{\substack{g_1+g_2+g_3 \\ =-n}} g_1 \text{diagram} \quad (5.59)$$

$$(h^0)_{22} = - \text{diagram} \quad (5.60)$$

We will show that Z' is homotopic to zero

$$\begin{array}{ccccc}
 \begin{pmatrix} \mathcal{F}\mathcal{E}1_n\{2\} \\ 1_n\{1+n\} \end{pmatrix} & \xrightarrow{\begin{pmatrix} \downarrow \uparrow \curvearrowright \\ \downarrow \uparrow \curvearrowright \end{pmatrix}} & \begin{pmatrix} \mathcal{F}\mathcal{E}1_n \\ \mathcal{F}\mathcal{E}1_n \end{pmatrix} & \xrightarrow{\begin{pmatrix} -\downarrow \uparrow & \downarrow \uparrow \\ \curvearrowright & -\curvearrowright \end{pmatrix}} & \begin{pmatrix} \mathcal{F}\mathcal{E}1_n\{-2\} \\ 1_n\{-n-1\} \end{pmatrix} \\
 \uparrow (Z')^{-1} & \swarrow h^0 & \uparrow (Z')^0 & \swarrow h^1 & \uparrow (Z')^1 \\
 \begin{pmatrix} \mathcal{E}\mathcal{F}\mathcal{F}\mathcal{E}1_n\{2\} \\ \mathcal{E}\mathcal{F}1_n\{1+n\} \end{pmatrix} & \xrightarrow{\begin{pmatrix} \uparrow \downarrow \downarrow \uparrow & \uparrow \downarrow \curvearrowright \\ \uparrow \downarrow \downarrow \uparrow & \uparrow \downarrow \curvearrowright \end{pmatrix}} & \begin{pmatrix} \mathcal{E}\mathcal{F}\mathcal{F}\mathcal{E}1_n \\ \mathcal{E}\mathcal{F}\mathcal{F}\mathcal{E}1_n \end{pmatrix} & \xrightarrow{\begin{pmatrix} -\uparrow \downarrow \downarrow \uparrow & \uparrow \downarrow \downarrow \uparrow \\ \uparrow \downarrow \curvearrowright & -\uparrow \downarrow \curvearrowright \end{pmatrix}} & \begin{pmatrix} \mathcal{E}\mathcal{F}\mathcal{F}\mathcal{E}1_n\{-2\} \\ \mathcal{E}\mathcal{F}1_n\{-n-1\} \end{pmatrix}
 \end{array} \tag{5.68}$$

where the chain map Z' is given by

$$(Z')_{11}^{-1} = \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} - \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} \tag{5.69}$$

$$(Z')_{12}^{-1} = (Z')_{21}^{-1} = (Z')_{22}^{-1} = 0. \tag{5.70}$$

$$(Z')_{11}^0 = \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} - \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} \tag{5.71}$$

$$(Z')_{12}^0 = 0 \tag{5.72}$$

$$(Z')_{22}^0 = -(Z')_{21}^0 = \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} - \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} \tag{5.73}$$

$$(Z')_{11}^1 = \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} - \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} \tag{5.74}$$

$$(Z')_{12}^1 = (Z')_{22}^1 = (Z')_{11}^0 = 0 \tag{5.75}$$

and the chain homotopy is given by

$$(h^0)_{11} = -(h^0)_{12} = \begin{array}{c} \uparrow n \\ \curvearrowright \end{array} \tag{5.76}$$

$$(h^0)_{21} = (h^0)_{22} = 0 \tag{5.77}$$

and

$$(h^1)_{11} = (h^1)_{21} = \begin{array}{c} \curvearrowright \\ \uparrow n \end{array} \quad (5.78)$$

$$(h^1)_{12} = (h^1)_{22} = 0. \quad (5.79)$$

Naturality of the other cup follows from the homotopy commutative diagram below:

$$\begin{array}{ccccc}
 & & \mathcal{C}\mathbf{1}_n & \xrightarrow{c} & \mathcal{C}\mathbf{1}_n \\
 & \nearrow \varrho^\omega & \omega(\mathcal{C})\mathbf{1}_n & \xrightarrow{\omega(\mathcal{C})} & \omega(\mathcal{C})\mathbf{1}_n \\
 & & \uparrow \omega(\mathcal{C}) & & \uparrow \omega(\mathcal{C}) \\
 & & \mathcal{F}\mathcal{E}\omega(\mathcal{C})\mathbf{1}_n & \xrightarrow{\mathcal{F}\omega(\xi^{-1-n})} & \mathcal{F}\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n \xrightarrow{\text{Id}} \mathcal{F}\omega(\mathcal{C})\mathcal{E}\mathbf{1}_n \xrightarrow{\omega(\widehat{\xi}^+ \mathbf{1}_{-n})\mathcal{E}} & \omega(\mathcal{C})\mathcal{F}\mathcal{E}\mathbf{1}_n \\
 & \nearrow \mathcal{F}\mathcal{E}\varrho^\omega & \uparrow \mathcal{F}\varrho^\omega \mathcal{E} & & \uparrow \mathcal{F}\widehat{\varrho}^\omega \mathcal{E} \\
 \mathcal{F}\mathcal{E}\mathcal{C}\mathbf{1}_n & \xrightarrow{\mathcal{F}\widehat{\xi}^+} & \mathcal{F}\mathcal{C}\mathcal{E}\mathbf{1}_n & \xrightarrow{\xi^{-\mathcal{E}}} & \mathcal{C}\mathcal{F}\mathcal{E}\mathbf{1}_n \\
 & & \downarrow \widehat{\varrho}^\omega \mathcal{F}\mathcal{E} & & \downarrow \widehat{\varrho}^\omega \mathcal{F}\mathcal{E} \\
 & & \mathcal{C}\mathbf{1}_n & \xrightarrow{c} & \mathcal{C}\mathbf{1}_n
 \end{array} \quad (5.80)$$

The center rectangle follows by applying the symmetry ω to the homotopy commutative square (5.42) and replacing n by $-n$. The two squares on the left and right commute by naturality of ϱ^ω , $\widehat{\varrho}^\omega$. The top square and the bottom middle triangle are commutative since ϱ^ω is the homotopy inverse of $\widehat{\varrho}^\omega$ as shown in Proposition 4.1. The bottom two squares follow from equations (4.94) and (4.97) in Proposition 4.12.

5.2.5 Naturality of $\widehat{\kappa}$ for cap 2-morphisms

We show that the squares

$$\begin{array}{ccc}
 \mathcal{C}\mathbf{1}_n & \xrightarrow{\widehat{\kappa}_{\mathbf{1}_n}} & \mathcal{C}\mathbf{1}_n \\
 \uparrow c & & \uparrow c \\
 \mathcal{C}\mathcal{E}\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\kappa}_{\mathcal{E}\mathcal{F}\mathbf{1}_n}} & \mathcal{E}\mathcal{F}\mathcal{C}\mathbf{1}_n
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{C}\mathbf{1}_n & \xrightarrow{\widehat{\kappa}_{\mathbf{1}_n}} & \mathcal{C}\mathbf{1}_n \\
 \uparrow c & & \downarrow c \\
 \mathcal{C}\mathcal{F}\mathcal{E}\mathbf{1}_n & \xrightarrow{\widehat{\kappa}_{\mathcal{F}\mathcal{E}\mathbf{1}_n}} & \mathcal{F}\mathcal{E}\mathcal{C}\mathbf{1}_n
 \end{array} \quad (5.81)$$

commute up to homotopy. To show that the first square is homotopy commutative consider the diagram below:

$$\begin{array}{ccccc}
 \mathcal{C}1_n & \xrightarrow{\quad \mathcal{C} \quad} & \mathcal{C}1_n & & \\
 \downarrow \varrho^{\sigma\omega} & & \downarrow \widehat{\varrho}^{\sigma\omega} & & \\
 \sigma\omega(\mathcal{C})1_n & \xrightarrow{\quad \sigma\omega(\mathcal{C}) \quad} & \sigma\omega(\mathcal{C})1_n & & \\
 \uparrow c & & \uparrow c & & \\
 \sigma\omega(\mathcal{C})\mathcal{E}\mathcal{F}1_n & \xrightarrow{\quad \sigma\omega(\xi^-)\mathcal{F} \quad} & \mathcal{E}\sigma\omega(\mathcal{C})\mathcal{F}1_n & \xrightarrow{\quad \text{Id} \quad} & \mathcal{E}\sigma\omega(\mathcal{C})\mathcal{F}1_n & \xrightarrow{\quad \mathcal{E}\sigma\omega(\xi^+) \quad} & \mathcal{E}\mathcal{F}\sigma\omega(\mathcal{C})1_n & \\
 \uparrow \varrho^{\sigma\omega}\mathcal{E}\mathcal{F} & & \downarrow \mathcal{E}\widehat{\varrho}^{\sigma\omega}\mathcal{F} & & \uparrow \mathcal{E}\varrho^{\sigma\omega}\mathcal{F} & & \downarrow \mathcal{E}\mathcal{F}\widehat{\varrho}^{\sigma\omega} & \\
 \mathcal{C}\mathcal{E}\mathcal{F}1_n & \xrightarrow{\quad \xi^+\mathcal{F} \quad} & \mathcal{E}\mathcal{C}\mathcal{F}1_n & \xrightarrow{\quad \widehat{\xi}^- \quad} & \mathcal{E}\mathcal{F}\mathcal{C}1_n & & & \\
 & & & & & & &
 \end{array} \quad (5.82)$$

The center rectangle is obtained by applying $\sigma\omega$ to the homotopy commutative square (5.42). The triangle commutes since $\varrho^{\sigma\omega}$ has inverse $\widehat{\varrho}^{\sigma\omega}$ by Proposition 4.1. The bottom two squares were shown to commute in Proposition 4.10. The remaining squares commute up to homotopy by the naturality of $\varrho^{\sigma\omega}$ and Propositions 4.1.

The second naturality square in (5.81) can be shown to commute by applying $\sigma\omega$ to (5.80) and arguing as above.

5.2.6 Naturality of κ for the cup 2-morphisms

We must show that the squares

$$\begin{array}{ccc}
 \mathcal{E}\mathcal{F}\mathcal{C}1_n & \xrightarrow{\widehat{\kappa}_{\mathcal{E}\mathcal{F}1_n}} & \mathcal{C}\mathcal{E}\mathcal{F}1_n \\
 \uparrow c & & \uparrow c \\
 \mathcal{C}1_n & \xrightarrow{\widehat{\kappa}_{1_n}} & \mathcal{C}1_n
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{F}\mathcal{E}\mathcal{C}1_n & \xrightarrow{\widehat{\kappa}_{\mathcal{F}\mathcal{E}1_n}} & \mathcal{C}\mathcal{F}\mathcal{E}1_n \\
 \uparrow c & & \uparrow c \\
 \mathcal{C}1_n & \xrightarrow{\widehat{\kappa}_{1_n}} & \mathcal{C}1_n
 \end{array} \quad (5.83)$$

commute up to homotopy. The proof is given by the following diagrams:

$$\begin{array}{ccccc}
 \mathcal{EFC1}_n & \xrightarrow{\mathcal{E}\xi^-} & \mathcal{ECF1}_n & \xrightarrow{\widehat{\xi}^+\mathcal{F}} & \mathcal{CEF1}_n \\
 \downarrow \mathcal{EF}\varrho^\psi & & \downarrow \mathcal{E}\widehat{\varrho}^\psi\mathcal{F} & & \downarrow \mathcal{E}\varrho^\psi\mathcal{F} \\
 \mathcal{EF}\psi(\mathcal{C})\mathbf{1}_n & \xrightarrow{\mathcal{E}\psi(\widehat{\xi}^-)} & \mathcal{E}\psi(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\text{Id}} & \mathcal{E}\psi(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\psi(\widehat{\xi}^+)\mathcal{F}} & \psi(\mathcal{C})\mathcal{EF}\mathbf{1}_n \\
 \uparrow \psi(\mathcal{C}) & & \uparrow \psi(\mathcal{C}) & & \uparrow \psi(\mathcal{C}) & & \uparrow \widehat{\varrho}^\psi\mathcal{EF} \\
 \psi(\mathcal{C})\mathbf{1}_n & \xrightarrow{\psi(\mathcal{C})} & \psi(\mathcal{C})\mathbf{1}_n & & \psi(\mathcal{C})\mathbf{1}_n & & \psi(\mathcal{C})\mathbf{1}_n \\
 \uparrow \varrho^\psi & & & & & & \downarrow \widehat{\varrho}^\psi \\
 \mathcal{C1}_n & \xrightarrow{\psi(\mathcal{C})} & \mathcal{C1}_n & & \mathcal{C1}_n & & \mathcal{C1}_n
 \end{array} \quad (5.84)$$

c

$$\begin{array}{ccccc}
 \mathcal{FEC1}_n & \xrightarrow{\mathcal{F}\widehat{\xi}^+} & \mathcal{FCE1}_n & \xrightarrow{\xi^-\mathcal{F}} & \mathcal{CFE1}_n \\
 \downarrow \mathcal{FE}\varrho^\psi & & \downarrow \mathcal{F}\widehat{\varrho}^\psi\mathcal{E} & & \downarrow \mathcal{F}\varrho^\psi\mathcal{E} \\
 \mathcal{FE}\psi(\mathcal{C})\mathbf{1}_n & \xrightarrow{\mathcal{F}\psi(\widehat{\xi}^+)} & \mathcal{F}\psi(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\text{Id}} & \mathcal{F}\psi(\mathcal{C})\mathcal{E}\mathbf{1}_n & \xrightarrow{\psi(\widehat{\xi}^-)\mathcal{E}} & \psi(\mathcal{C})\mathcal{FE}\mathbf{1}_n \\
 \uparrow \psi(\mathcal{C}) & & \uparrow \psi(\mathcal{C}) & & \uparrow \psi(\mathcal{C}) & & \uparrow \widehat{\varrho}^\psi\mathcal{FE} \\
 \psi(\mathcal{C})\mathbf{1}_n & \xrightarrow{\psi(\mathcal{C})} & \psi(\mathcal{C})\mathbf{1}_n & & \psi(\mathcal{C})\mathbf{1}_n & & \psi(\mathcal{C})\mathbf{1}_n \\
 \uparrow \varrho^\psi & & & & & & \downarrow \widehat{\varrho}^\psi \\
 \mathcal{C1}_n & \xrightarrow{\psi(\mathcal{C})} & \mathcal{C1}_n & & \mathcal{C1}_n & & \mathcal{C1}_n
 \end{array} \quad (5.85)$$

c

The center rectangles commute up to homotopy since they are obtained from the homotopy commutative squares in (5.81). The triangles and the left, bottom and right squares in both diagrams above commute since ϱ^ψ is inverse $\widehat{\varrho}^\psi$ by Proposition 4.1. The remaining squares commute by Proposition 4.9.

5.2.7 Naturality of $\widehat{\kappa}$ for the cup 2-morphisms

The naturality for one cap is given by the commutative diagram in $Com(\mathcal{U})$

$$\begin{array}{ccccc}
 \mathcal{CE}\mathcal{F}\mathbf{1}_n & \xrightarrow{\varepsilon\xi^+} & \mathcal{EC}\mathcal{F}\mathbf{1}_n & \xrightarrow{\widehat{\xi}^-\mathcal{F}} & \mathcal{E}\mathcal{F}\mathcal{C}\mathbf{1}_n \\
 \uparrow \varrho^\psi \mathcal{E}\mathcal{F} & & \varepsilon\widehat{\varrho}^\psi \mathcal{F} \uparrow & & \mathcal{E}\mathcal{F}\widehat{\varrho}^\psi \uparrow \\
 \psi(\mathcal{C})\mathcal{E}\mathcal{F}\mathbf{1}_n & \xrightarrow{\psi(\widehat{\xi}^+)\mathcal{F}} & \mathcal{E}\psi(\mathcal{C})\mathcal{F}\mathbf{1}_n & \xrightarrow{\varepsilon\psi(\xi^-)} & \mathcal{E}\mathcal{F}\psi(\mathcal{C})\mathbf{1}_n \\
 \uparrow \psi(\mathcal{C}) & & \uparrow \psi(\mathcal{C}) & & \uparrow \psi(\mathcal{C}) \\
 \psi(\mathcal{C})\mathbf{1}_n & \xrightarrow{\psi(\mathcal{C})} & \psi(\mathcal{C})\mathbf{1}_n & & \\
 \uparrow \varrho^\psi & & & & \downarrow \widehat{\varrho}^\psi \\
 \mathcal{C}\mathbf{1}_n & \xrightarrow{c} & \mathcal{C}\mathbf{1}_n & &
 \end{array} \tag{5.86}$$

where the middle square commutes up to homotopy by applying ψ to the homotopy commutative diagram (5.42). The left and right squares commute by the naturality of ϱ^ψ and $\widehat{\varrho}^\psi$. The top square commutes on the nose since ϱ^ψ has inverse $\widehat{\varrho}^\psi$. The bottom two squares commute by Proposition 4.9.

Naturality for $\widehat{\kappa}$ with respect to the other cap is proven similarly.

References

- [1] P. Balmer and M. Schlichting. Idempotent completion of triangulated categories. *J. Algebra*, 236(2):819–834, 2001.
- [2] A. Beilinson, G. Lusztig, and R. MacPherson. A geometric setting for the quantum deformation of GL_n . *Duke Math. J.*, 61(2):655–677, 1990.
- [3] A. Beliakova and S. Wehrli. Categorification of the colored Jones polynomial and Rasmussen invariant of links. *Canad. J. Math.*, 60(6):1240–1266, 2008. arXiv:math/0510382.
- [4] D. Benson. *Representations and cohomology. I: Basic representation theory of finite groups and associative algebras*, volume 30 of *Cambridge Studies in Advanced Mathematics*. Cambridge U. Press, second edition, 1998.
- [5] Kassel C and V. Turaev. Double construction for monoidal categories. *Acta Math.*, 175(1):1–48, 1995.
- [6] L. Crane and I. B. Frenkel. Four-dimensional topological quantum field theory, Hopf categories, and the canonical bases. *J. Math. Phys.*, 35(10):5136–5154, 1994, hep-th/9405183.

- [7] I.B. Frenkel. Unpublished notes, circa 1993.
- [8] K. Habiro. A unified Witten-Reshetikhin-Turaev invariant for integral homology spheres. *Invent. Math.*, 171(1):1–81, 2008. arXiv:math/0605314.
- [9] A. Joyal and R. Street. Tortile Yang-Baxter operators in tensor categories. *J. Pure Appl. Algebra*, 71(1):43–51, 1991.
- [10] C. Kassel. *Quantum groups*, volume 155 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [11] M. Khovanov. A categorification of the Jones polynomial. *Duke Math. J.*, 101(3):359–426, 2000. math.QA/9908171.
- [12] M. Khovanov and A. Lauda. A diagrammatic approach to categorification of quantum groups III. *Quantum Topology*, 1:1–92, 2010, math.QA/0807.3250.
- [13] M. Khovanov, A. Lauda, M. Mackaay, and M. Stošić. Extended graphical calculus for categorified quantum $\mathfrak{sl}(2)$. 2010, arXiv:1006.2866.
- [14] A. Lauda. Categorified quantum $\mathfrak{sl}(2)$ and equivariant cohomology of iterated flag varieties. math.QA/0803.3848, 2008. To appear in *Algebras and Representation Theory*.
- [15] A. D. Lauda. A categorification of quantum $\mathfrak{sl}(2)$. *Adv. Math.*, 225:3327–3424, 2010. math.QA/0803.3652.
- [16] R.J. Lawrence. A universal link invariant using quantum groups. In *Differential geometric methods in theoretical physics (Chester, 1988)*, pages 55–63. World Sci. Publ., Teaneck, NJ, 1989.
- [17] R.J. Lawrence. A universal link invariant. In *The interface of mathematics and particle physics (Oxford, 1988)*, volume 24 of *Inst. Math. Appl. Conf. Ser. New Ser.*, pages 151–156. Oxford Univ. Press, New York, 1990.
- [18] G. Lusztig. *Introduction to quantum groups*, volume 110 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1993.
- [19] S. Majid. Representations, duals and quantum doubles of monoidal categories. In *Proceedings of the Winter School on Geometry and Physics (Srńí, 1990)*, number 26, 1991.
- [20] N. Reshetikhin and V. Turaev. Invariants of 3-manifolds via link polynomials and quantum groups. *Invent. Math.*, 103(3):547–597, 1991.
- [21] E. Witten. Quantum field theory and the Jones polynomial. *Comm. Math. Phys.*, 121(3):351–399, 1989.

A.B.: *Institut für Mathematik, Universität Zürich, Winterthurerstr. 190 CH-8057 Zürich*
email: `anna@math.uzh.ch`

M.K.: *Department of Mathematics, Columbia University, New York, NY 10027, USA*
email: `khovanov@math.columbia.edu`

A.L.: *Department of Mathematics, Columbia University, New York, NY 10027, USA*
email: `lauda@math.columbia.edu`