

ON THE INTEGRALITY OF THE WITTEN–RESHETIKHIN–TURAEV 3–MANIFOLD INVARIANTS

ANNA BELIAKOVA, QI CHEN, AND THANG LE

ABSTRACT. We prove that the $SU(2)$ and $SO(3)$ Witten–Reshetikhin–Turaev invariants of any 3–manifold with any colored link inside at any root of unity are algebraic integers.

0. INTRODUCTION

In the late 80s, Witten [Wi] and Reshetikhin–Turaev [RT] associated with any closed oriented 3-manifold M (possibly with a colored link inside), any root of unity ξ and any compact Lie group G a complex number $\tau_M^G(\xi)$, called the quantum or WRT invariant of M .

For more than 20 years, the problem of integrality of the WRT invariants has been intensively studied. The interest to this problem was drawn by the theory of perturbative 3-manifold invariants generalizing those of Casson and Walker [O], by the construction of Integral Topological Quantum Field Theories [G, GM] and their topological applications and more recently, by attempts to categorify the WRT invariants [Kho].

In the case $G = SU(2)$, there is a projective version $\tau_M^{SO(3)}(\xi)$, introduced by Kirby and Melvin [KM] and defined at roots of unity of *odd* order. This projective version, when defined, determines the $SU(2)$ version.

In this paper we completely solve the integrality problem for both $SO(3)$ and $SU(2)$ versions of the WRT invariant for all 3-manifolds with arbitrary links inside. Before stating our results, let us give a brief introduction into the history of this subject.

In 1995 Murakami [Mu] established the integrality of the WRT $SO(3)$ -invariant for rational homology 3-spheres at roots of unity of *prime orders*. This result was extended to all 3-manifolds by Masbaum and Roberts [MR]. Masbaum and Wenzl [MW], and independently Takata and Yokota [TY], proved the integrality of the projective WRT $SU(n)$ -invariant for all 3-manifolds, always under the assumption that the orders of the roots of unity are *prime*. Finally the third author [Le2] established the integrality of the projective WRT invariant associated with any compact simple Lie group, again at roots of unity of prime orders.

The case for the roots of unity of *non-prime* orders is more complicated. The first integrality result for *all roots of unity* was obtained by Habiro [Ha2] in the case of $SU(2)$ and *integral homology 3-spheres*. Habiro’s proof relies on the existence of the unified invariant for integral homology 3-spheres as an element of Habiro’s ring, a certain cyclotomic completion of the polynomial ring $\mathbb{Z}[q]$. This unified invariant is a kind of generating function for the set of WRT $SU(2)$ invariants at all roots of unity. The integrality in this approach follows directly from the general properties of Habiro’s ring.

Habiro and the third author [HL] subsequently defined the unified WRT invariant for all *simple Lie groups* and *integral homology 3-spheres*, thus proving that the WRT invariant

A.B. and T.L. are supported by the Swiss and American National Science Foundations.

Mathematics Classification (2010): Primary 57M27; Secondary 57M25.

of any *integral homology 3-sphere* associated to any simple Lie group and any root of unity is always an algebraic integer. However, the case of manifolds other than homology spheres was unknown, even with $G = SU(2)$.

In this paper we give a complete solution for the integrality problem for *all 3-manifolds* with *arbitrary* link inside at *all roots* of unity for the case of the group $SU(2)$. Our invariants are normalized as in [KM] and we show that integrality in that case implies integrality for all other normalizations used in the literature.

Theorem 1. *The WRT $SU(2)$ -invariant of any 3-manifold M with any colored link inside at any root of unity is an algebraic integer.*

Theorem 2. *The WRT $SO(3)$ -invariant of any 3-manifold M with any colored link inside at any root of unity of odd order is an algebraic integer.*

Theorem 2 is a generalization of a result in [BL] to manifolds which contain a link inside. However, we give here a new independent proof along the same lines as in the $SU(2)$ case. Theorem 1 is the main result of the paper. The key new ideas used in the proofs are the following.

One of the main tools is a significant generalization of some divisibility result (Theorem 2.2) which was originally obtained in [Le3] using a number-theoretical identity of Andrews' [A], whose special cases are the classical Rogers-Ramanujan identities.

Further, to include the case of even colored links in 3-manifolds, we had to introduce a new basis for the Grothendieck ring of the quantum $sl(2)$, which is orthogonal to the odd part of the center with respect to the Rosso form. This led to an important new result (Theorem 1.1) generalizing that of Habiro, which states that the colored Jones polynomial can be presented as a sum of integral "blocks". This result is proved in the Appendix, and it is of independent interest in the quantum link invariant theory.

For manifolds obtained by surgery along links with diagonal linking matrix we show that the contribution of each integral block to the WRT invariant is integral by using our main tool (Theorem 2.2). The general case can be reduced to the diagonal one by using some classification results for linking pairings. However, it is more demanding in the $SU(2)$ case than in the $SO(3)$ one, since the linking pairings on abelian groups of even order are more complicated [KK].

As a byproduct, we generalize the relationship between $SU(2)$ and $SO(3)$ invariants at odd roots of unity to the case when a 3-manifold contains an arbitrary colored link inside. For empty links and links colored by the fundamental representation, this relationship was established in [KM] and [MR], respectively.

At the moment of this writing, our proof cannot be generalized to higher-ranked Lie groups because we do not have an analog of Theorem 1.1 (splitting into integral blocks) in those cases. The paper is as self-contained as possible. The only two results used without proofs here are [Le3, Theorem 7] and [BBIL, Theorem 2].

We organize this paper as follows. In Section 1 we fix notations, recall the definition of the WRT invariant and state a generalization of Habiro's result. The main strategy of our proofs is explained in Section 1.6. In Section 2 we prove some divisibility results for generic values of the quantum parameter. Formulas related to roots of unity are proved in Section 3. Section 4 deals with the symmetry principle and the splitting of the $SU(2)$ invariant at odd roots of unity into the product of the $SO(3)$ and Deloup's invariants. Section 5 discusses how to construct 3-manifolds that can be obtained by surgery along links with diagonal linking matrices. The last two sections are devoted to the proofs of Theorem 2 and Theorem 1, respectively.

1. THE COLORED JONES POLYNOMIAL AND THE WRT INVARIANT

1.1. **Notations.** Let $q^{1/4}$ be a formal parameter. Set

$$\{n\} := q^{n/2} - q^{-n/2}, \quad \{n\}! := \prod_{i=1}^n \{i\}, \quad [n] := \frac{\{n\}}{\{1\}}, \quad \left[\begin{matrix} n \\ k \end{matrix} \right] := \frac{\{n\}!}{\{k\}!\{n-k\}!},$$

and

$$(z; q)_m = \prod_{i=0}^{m-1} (1 - q^i z), \quad \binom{m}{n}_q := \frac{(q^{m-n+1}; q)_n}{(q; q)_n} = q^{(m-n)n/2} \left[\begin{matrix} m \\ n \end{matrix} \right].$$

Throughout this paper, let ξ be a primitive root of unity of order r and $\xi^{1/4}$ be a complex number such that $(\xi^{1/4})^4 = \xi$. There are 4 possible choices for $\xi^{1/4}$, and we will make some restrictions later.

When working in the $SO(3)$ case, we will always assume that $r \geq 3$ is *odd*. In the $SU(2)$ case, $r \geq 2$ will be an arbitrary positive integer.

For $f \in \mathbb{Q}[q^{\pm 1/4}]$, we define the following evaluation map

$$\text{ev}_\xi(f) := f|_{q^{1/4}=\xi^{1/4}}.$$

It should be noted that although we write $\text{ev}_\xi(f)$, this quantity depends on the choice of a 4-th root $\xi^{1/4}$ of ξ .

If f is a function on positive integers n_1, \dots, n_k with values in $\mathbb{Q}[q^{\pm 1/4}]$, we define

$$\sum_{n_1, \dots, n_k} \xi, SO(3) f := \frac{1}{4^k} \sum_{\substack{n_j=0 \\ n_j \text{ odd}}}^{4r-1} \text{ev}_\xi(f), \quad \sum_{n_1, \dots, n_k} \xi, SU(2) f := \frac{1}{4^k} \sum_{n_j=0}^{4r-1} \text{ev}_\xi(f).$$

All 3-manifolds in this paper are supposed to be closed and oriented. Every link in a 3-manifold is framed, oriented and has ordered components.

1.2. **The colored Jones polynomial.** Suppose L is a framed oriented link in S^3 with m ordered components. For an m -tuple of positive integers $\mathbf{n} = (n_1, \dots, n_m)$, one has the colored Jones polynomial $J_L(\mathbf{n}) \in \mathbb{Z}[q^{\pm 1/4}]$, see e.g. [Tu, MM]. The number n_i is usually called the color of the i -th component, and stands for the n_i -dimensional irreducible sl_2 -representation in the theory of quantum link invariants. We use the normalization so that $J_U(n) = [n]$ where U is the unknot with 0 framing. It is well known that if \tilde{L} is obtained from L by increasing the framing on the i -th component by 1 then

$$J_{\tilde{L}}(\mathbf{n}) = q^{\frac{n_i^2-1}{4}} J_L(\mathbf{n}). \quad (1)$$

Although there are fractional powers $q^{\pm 1/4}$, there exists an integer $a = a(L, \mathbf{n})$ such that $J_L(\mathbf{n}) \in q^{a/4} \mathbb{Z}[q^{\pm 1}]$. For a precise formula of a see [Le1]. This formula implies that if all the colors n_j 's are odd, then $J_L(\mathbf{n}) \in \mathbb{Z}[q^{\pm 1}]$.

1.3. **Habiro's expansion and its generalization.** Assume that $L \sqcup L'$ is a framed link in S^3 with disjoint sublinks L and L' . Suppose L has m ordered components and L' has l ordered components. Fix an l -tuple of positive integers $\mathbf{s} = (s_1, \dots, s_l)$, and let's consider $J_{L \sqcup L'}(\mathbf{n}, \mathbf{s})$ as a function on m -tuples $\mathbf{n} = (n_1, \dots, n_m)$. Since \mathbf{s} is fixed, we will remove it from the notation for simplicity. The function $J_{L \sqcup L'}(\mathbf{n})$ can be rearranged into another function $c_{L \sqcup L'}(\mathbf{k})$ generalizing an important result of Habiro [Ha2, Theorem 8.2].

To state the result we need to introduce a few notations. Let $\tilde{\ell}_{ij}$ be the linking number between the i -th component of L and the j -th component of L' . For any $i = 1, \dots, m$, we define

$$\varepsilon_i \in \{0, 1\} \quad \text{by} \quad \varepsilon_i := \sum_{j=1}^l \tilde{\ell}_{ij}(s_j - 1) \pmod{2}. \quad (2)$$

Theorem 1.1. *Assume that $L \sqcup L' \subset S^3$ is as described above. Suppose that L has 0 linking matrix. Then for every m -tuple $\mathbf{k} = (k_1, \dots, k_m)$ of non-negative integers with $k = \max(k_1, \dots, k_m)$ there exists*

$$c_{L \sqcup L'}(\mathbf{k}) \in \frac{(q^{k+1}; q)_{k+1}}{1-q} \mathbb{Z}[q^{\pm 1/4}] \quad (3)$$

such that for every m -tuple $\mathbf{n} = (n_1, \dots, n_m)$ of non-negative integers,

$$J_{L \sqcup L'}(\mathbf{n}) = \sum_{k_i \geq 0} c_{L \sqcup L'}(\mathbf{k}) \prod_{i=1}^m \begin{bmatrix} n_i + k_i \\ 2k_i + 1 \end{bmatrix} \{k_i\}! \frac{\lambda_{n_i}^{\varepsilon_i}}{\lambda_{k_i+1}^{\varepsilon_i}} \quad (4)$$

where $\lambda_n = q^{n/2} + q^{-n/2}$.

For the case when all $\varepsilon_i = 0$, or, in particular, when all s_i 's are odd, the statement is equivalent to [BBuL, Theorem 3]. A proof of Theorem 1.1 is given in Appendix A. Note that for a fixed \mathbf{n} the right hand side of (4) is a finite sum because $\begin{bmatrix} n+k \\ 2k+1 \end{bmatrix} = 0$ if $n \leq k$.

This is the presentation of the colored Jones polynomial as a sum of integral blocks mentioned in Introduction. The existence of $c_{L \sqcup L'}(\mathbf{k}) \in \mathbb{Q}(q^{1/4})$ that satisfies (4) is easy to prove. The real content of Theorem 1.1 is the integrality (3).

1.4. The WRT invariant. We review here the definition of the WRT $SU(2)$ invariant of a 3-manifold M with a colored link L' inside [RT] and its $SO(3)$ version [KM].

We use the convention that the pair (M, L') is obtained from (S^3, L') by surgery along L . Here L' is an \mathbf{s} -colored framed link. For $G = SU(2)$ or $G = SO(3)$ set

$$F_{L \sqcup L'}^G(\xi) := \sum_{n_1, \dots, n_m} \xi^{G} \left\{ J_{L \sqcup L'}(\mathbf{n}) \prod_{i=1}^m [n_i] \right\}. \quad (5)$$

For simplicity, we assume here that all entries of \mathbf{s} are odd if $G = SO(3)$. In general for $G = SO(3)$, we have to multiply (5) by a power of ξ , depending on the linking matrix of L' and the parity of colors. This is done in Section 4.2. Since the additional factor is a unit, it does not affect integrality.

We want to emphasize that although it is not explicit from the notation, (5) depends on a choice of a 4-th root $\xi^{1/4}$ of ξ .

It is known that $F_{L \sqcup L'}^G(\xi)$ is invariant under the handle slide move and if normalized appropriately, is an invariant of the pair (M, L') .

Let U^\pm be the unknot with ± 1 framing. It is easy to see that $F_{U^-}^G(\xi)$ is the complex conjugate of $F_{U^+}^G(\xi)$. Let

$$\mathcal{D}^G := |F_{U^+}^G(\xi)| = \sqrt{F_{U^+}^G(\xi) F_{U^-}^G(\xi)}. \quad .$$

This number is called the rank of a TQFT in [Tu]. We normalize by dividing (5) by certain powers of $F_{U^\pm}^G(\xi) \neq 0$. Hence, we want to know when $F_{U^\pm}^G(\xi) \neq 0$. The following is probably known. For completeness we include a proof in Section 3.3.

Lemma 1.2. *One has $F_{U^\pm}^G(\xi) = 0$ if and only if*

$$G = SU(2) \text{ and } \xi^{1/4} \text{ has order } 2 \text{ ord}(\xi) = 2r. \quad (\star)$$

In [KM, RT] and [Lic] it is assumed that $\text{ord}(\xi^{1/4}) = 4 \text{ ord}(\xi)$. However, there are other cases when $F_{U^\pm}^G(\xi) \neq 0$. Here we consider all of them.

In the entire paper we will assume that condition (\star) is not satisfied, so that $F_{U^\pm}^G(\xi) \neq 0$. Then the WRT invariant of the pair (M, L') is defined by

$$\tau_{M,L'}^G(\xi) = \frac{F_{L \sqcup L'}^G(\xi)}{(F_{U^+}^G(\xi))^{\beta_+} (F_{U^-}^G(\xi))^{\beta_-} (\mathcal{D}^G)^\beta}, \quad (6)$$

where β_+, β_- and β are respectively the number of positive, negative, and 0 eigenvalues of the linking matrix of L .

The invariant $\tau_{M,L'}^G(\xi)$ is multiplicative with respect to connected sum. If $-M$ is M with the reverse orientation, then $\tau_{-M}^G(\xi)$ is the complex conjugate of $\tau_M^G(\xi)$, and $\tau_{S^3}^G(\xi) = 1$.

Remark 1.3. We will prove later that $\mathcal{D}^G \in \mathbb{Z}[\xi^{1/4}, e_8]$, where $e_8 = \exp(\pi\sqrt{-1}/4)$. Note that $\mathbb{Z}[\xi^{1/4}, e_8] = \mathbb{Z}[\exp(2\pi\sqrt{-1}/t)]$, where $t = 8r$ if r is odd and $t = 4r$ if r is even. In the last case, $e_8 \in \mathbb{Z}[\xi^{1/4}]$.

Hence a priori, $\tau_{M,L'}^G(\xi) \in \mathbb{Q}(\xi^{1/4}, e_8)$. Since the ring of integers of $\mathbb{Q}(\xi^{1/4}, e_8)$ is $\mathbb{Z}[\xi^{1/4}, e_8]$, our invariant is algebraically integral if it belongs to $\mathbb{Z}[\xi^{1/4}, e_8]$.

Further, if $G = SO(3)$, M is a rational homology 3-sphere, and all the s_i 's are odd, then $\tau_{M,L'}^{SO(3)}(\xi) \in \mathbb{Q}(\xi)$ by definition. So, in that case integrality means that $\tau_{M,L'}^{SO(3)}(\xi) \in \mathbb{Z}[\xi]$ for any root of unity ξ of odd order.

Relations with other invariants. If we put $\xi^{1/4} := \exp(\pi\sqrt{-1}/2r)$, then our invariant $\tau_M^{SU(2)}(\xi)$ and $\tau_M^{SO(3)}(\xi)$ are respectively $\tau_r(M)$ and $\tau_r'(M)$ in [KM]. In that case, our $\mathcal{D}^{SU(2)}$ equals to b^{-1} in the notation of [KM].

Again, if $\xi^{1/4} = \exp(\pi\sqrt{-1}/2r)$, the original Reshetikhin-Turaev invariant [RT] differs from $\tau_r(M)$ by a multiplication with a certain root of unity, so this does not affect integrality.

The set of invariants considered in [MR] coincides with ours assuming r is an odd prime. More precisely, the invariants $I_{2r}(M)$ and $I_r(M)$, defined in [MR] as functions of a variable A , coincide with ours $\tau_M^{SU(2)}(\xi)$ and $\tau_M^{SO(3)}(\xi)$ after setting $A = -\xi^{1/4}$ and $A = -\xi^{(r+1)^2/4}$, respectively. At these roots of unity, the $SO(3)$ invariants determines those for $SU(2)$.

In [Lic], Lickorish chose a different normalization and worked with $\tau_M^{SU(2)}(\xi)(\mathcal{D}^{SU(2)})^\beta$ in our notation. Clearly, integrality of this invariant will follow from the integrality of $\tau_{M,L'}^G(\xi)$.

1.5. Diagonal case. Of particular importance is the case when the linking matrix of L is a diagonal matrix $\text{diag}(b_1, \dots, b_m)$, $b_i \in \mathbb{Z}$ for any i . Let L^0 be the framed link obtained from L by switching all the framings to 0. Recall from (2) that for $1 \leq i \leq m$, $\varepsilon_i := \sum_{k=1}^i \tilde{\ell}_{ik}(s_k - 1) \pmod{2}$. Using (1) and (4), we can rewrite $F_{L \sqcup L'}^G(\xi)$ as follows:

$$F_{L \sqcup L'}^G(\xi) = \sum_{k_i \geq 0} \text{ev}_\xi(c_{L^0 \sqcup L'}(\mathbf{k})/\{1\}^m) \prod_{i=1}^m H^G(k_i, b_i, \varepsilon_i) \quad (7)$$

where

$$H^G(k, b, \varepsilon) := \sum_n^{\xi, G} q^{b \frac{n^2-1}{4}} \begin{bmatrix} n+k \\ 2k+1 \end{bmatrix} \frac{\lambda_n^\varepsilon}{\lambda_{k+1}^\varepsilon} \{k\}! \{n\}. \quad (8)$$

By (5) and (8) we also have

$$F_{U^\pm}^G(\xi) = \frac{H^G(0, \pm 1, 0)}{\text{ev}_\xi(\{1\})}. \quad (9)$$

From (7) and (9) we get the following.

Proposition 1.4. *Suppose the linking matrix of L is a diagonal matrix $\text{diag}(b_1, \dots, b_m)$ with exactly t non-zero elements b_1, \dots, b_t . Assume the entries of \mathbf{s} are odd when $G = SO(3)$. Then*

$$\tau_{M,L'}^G(\xi) = \sum_{k_i=0}^{\lfloor \frac{r-2}{2} \rfloor} \text{ev}_\xi(c_{L^0 \sqcup L'}(\mathbf{k})) \prod_{i=1}^t \frac{H^G(k_i, b_i, \varepsilon_i)}{H^G(0, \text{sn}(b_i), 0)} \prod_{i=t+1}^m \frac{H^G(k_i, 0, \varepsilon_i)}{\text{ev}_\xi(\{1\}) \mathcal{D}^G}, \quad (10)$$

where $\text{sn}(b_i)$ is the sign of b_i .

Note that in the above sum the index k_i is from 0 to $\lfloor \frac{r-2}{2} \rfloor$. This is because $(\xi^{k+1}; \xi)_{k+1} = 0$ when $k > (r-2)/2$, so $\text{ev}_\xi(c_{L^0 \sqcup L'}(\mathbf{k})) = 0$ when $k = \max\{k_i\} > (r-2)/2$ according to Proposition 1.1.

To allow an arbitrary coloring \mathbf{s} of L' for $G = SO(3)$, we have to multiply the right hand side of (10) by a unit, defined in Section 4.2.

We say that M is *diagonal of prime type*, when M can be obtained by surgery along a link with diagonal linking matrix whose entries are (up to a sign) 0, 1 or prime powers.

1.6. Strategy for the proof of Theorems 1 and 2. We first prove the integrality of $\tau_{M,L'}^G(\xi)$ for the case when M is diagonal of prime type. By (10), in this case it suffices to show that

$$\frac{H^G(k, b, \varepsilon)}{H^G(0, \text{sn}(b), 0)} \quad \text{and} \quad \frac{H^G(k, 0, \varepsilon)}{(1 - \xi) \mathcal{D}^G}$$

are algebraic integers when $0 \leq k \leq \lfloor \frac{r-2}{2} \rfloor$. This is proved in Proposition 6.1 for $G = SO(3)$ and in Proposition 7.1 for $G = SU(2)$, under assumptions r is odd and even, respectively.

The general case can be reduced to the diagonal one of prime type by applying some standard results on diagonalization, presented in Section 5. Roughly speaking, $M \# M$ becomes diagonal of prime type after adding a diagonalizing manifold N , which is a connected sum of some simple lens spaces. In the $SO(3)$ case, this already solves the problem, since the WRT invariant of N is invertible. In the $SU(2)$ case, the WRT invariants of N might be 0. We show that there is an odd colored link $L \subset N$ such that $\tau_{N,L}^{SU(2)}$ is integral and non-zero. However, another difficulty arises since $\tau_{N,L}^{SU(2)}$ is not invertible. To overcome this difficulty we will look at the connected sum of many copies of $M \# M$ with (N, L) , which we will show to be diagonal of prime type. Further, we make substantial use of the fact that in any Dedekind domain, every ideal has a unique prime factorization.

The case $G = SU(2)$ and r odd is solved in Section 4. There we generalize the splitting formula of Kirby and Melvin [KM] by showing that the $SU(2)$ invariant of any 3-manifold with a colored link inside at a root of unity of odd order is a product of the $SO(3)$ invariant and another integer invariant, previously defined by Deloup.

2. BASIC DIVISIBILITY: THE CASE OF GENERIC q

In this section we establish a divisibility result for generic q which will help us to prove that each factor of (10) is integral.

2.1. **The ideal I_k .** Let I_k be the ideal of $\mathbb{Z}[z^{\pm 1}, q^{\pm 1}]$ generated by $(q^a z; q)_k$ for all $a \in \mathbb{Z}$. This ideal plays an important role in the theory of quantum invariants, see [Ha2, Le3, HL].

We will use the following characterization of I_k , which is Proposition 4.3 of [Le3].

Proposition 2.1. *The ideal I_k is the set of all $f \in \mathbb{Z}[z^{\pm 1}, q^{\pm 1}]$ such that $f(q^b, q)$ is divisible by $(q; q)_k$ for every $b \in \mathbb{Z}$.*

We will often use the following q -binomial formula

$$(q^a z; q)_k = \sum_{j=0}^k (-1)^j \binom{k}{j}_q q^{\binom{j}{2} + aj} z^j. \quad (11)$$

2.2. **Divisibility for generic q .** For a positive integer k let

$$X_k := \frac{(q; q)_k}{(q; q)_{\lfloor k/2 \rfloor}} = \prod_{j=\lfloor k/2 \rfloor + 1}^k (1 - q^j). \quad (12)$$

A map $Q : \mathbb{Z} \rightarrow \mathbb{Z}$ is said to be an *integral quadratic form* if

$$Q(n) = a_2 n^2 + a_1 n + a_0$$

for some integers a_2, a_1, a_0 .

For a quadratic form Q let $\mathcal{L}_Q : \mathbb{Z}[z^{\pm 1}, q^{\pm 1}] \rightarrow \mathbb{Z}[q^{\pm 1}]$ be the $\mathbb{Z}[q^{\pm 1}]$ -linear map defined by

$$\mathcal{L}_Q(z^j) = q^{Q(j)}.$$

Note that this map is not an algebra homomorphism if a_2 or $a_0 \neq 0$.

Let σ be the $\mathbb{Z}[q^{\pm 1}]$ -algebra automorphism of $\mathbb{Z}[z^{\pm 1}, q^{\pm 1}]$ defined by $\sigma(z) = z^{-1}$. An important observation is that if $a_1 = 0$, then $\mathcal{L}_Q \sigma = \mathcal{L}_Q$.

Theorem 2.2. *For an arbitrary integral quadratic form Q and any $f(z, q) \in I_k$, the element $\mathcal{L}_Q(f)$ is divisible by X_k , i.e.*

$$\mathcal{L}_Q(f) \in X_k \mathbb{Z}[q^{\pm 1}].$$

Remark 2.3. This theorem will be used substantially. It is a generalization of [Le3, Theorem 7], which was proved with the help of Andrews' identity. The case $Q(n) = n^2$ of Theorem 2.2 appeared in [HL] for the construction of the unified WRT invariant.

Proof. By the definition of I_k , it is enough to consider the case when $f = z^m (q^a z; q)_k$. Suppose

$$Q(n) = a_2 n^2 + a_1 n + a_0.$$

Let $Q_0(n) = a_2 n^2$. Rewriting f as a sum with help of (11), one can show

$$\mathcal{L}_Q(q^{-a_1 m} z^m (q^{a-a_1} z; q)_k) = q^{a_0} \mathcal{L}_{Q_0}(z^m (q^a z; q)_k).$$

Hence, the substitution of $q^{-a_1} z$ for z transforms \mathcal{L}_Q into \mathcal{L}_{Q_0} . Without loss of generality, we can further assume $Q = Q_0$.

The rest of the proof is by induction on k .

The case $k = 1$ is trivial. Suppose that the statement holds true for $k - 1$.

Since

$$z^m (q^{a+1} z; q)_k - z^m (q^a z; q)_k = q^a (1 - q^k) z^{m+1} (q^{a+1} z; q)_{k-1},$$

we see that, by the induction hypothesis, $\mathcal{L}_{Q_0}(z^m (q^{a+1} z; q)_k)$ is divisible by X_k if and only if $\mathcal{L}_{Q_0}(z^m (q^a z; q)_k)$ is. Therefore we only need to show the statement for a single value of a . We will take $a = -\lfloor k/2 \rfloor$. The cases of odd and even k will be considered separately.

Suppose $k = 2l + 1$. Then $a = -l$ and $\mathcal{L}_{Q_0}(z^m(q^{-l}z; q)_{2l+1})$ is divisible by $X_k = (q^{l+1}; q)_{l+1}$ by Lemma 2.4 (b) below.

Now suppose $k = 2l$. Then $a = -l$ and we need to show that for every integer m , X_k divides $\mathcal{L}_{Q_0}(B(m, l))$ where

$$B(m, l) := z^m(q^{-l}z; q)_{2l} .$$

Since

$$B(m, l) - q^l B(m+1, l) = z^m(q^{-l}z; q)_{2l+1}$$

and $X_k = (q^{l+1}; q)_l$ divides $\mathcal{L}_{Q_0}(z^m(q^{-l}z; q)_{2l+1})$ by Lemma 2.4 (b) below, it is enough to show that X_k divides $\mathcal{L}_{Q_0}(B(m, l))$ for only a single value of m . We choose $m = -l$, and we will show that $X_k = (q^{l+1}; q)_l$ divides $\mathcal{L}_{Q_0}(B(-l, l))$.

Using that σ is the algebra automorphism of $\mathbb{Z}[z^{\pm 1}, q^{\pm 1}]$ sending z to z^{-1} , we get

$$\begin{aligned} (\text{id} + \sigma)B(-l, l) &= z^{-l}(q^{-l}z; q)_{2l} + z^l(q^{-l}z^{-1}; q)_{2l} \\ &= z^{-l}(q^{-l}z; q)_{2l} + z^{-l}q^{-l}(q^{1-l}z; q)_{2l} \\ &= z^{-l}(q^{1-l}z; q)_{2l-1}(1 - q^{-l}z + q^{-l}(1 - q^l z)) \\ &= z^{-l}(q^{1-l}z; q)_{2l-1}(1 - z)(1 + q^{-l}) \\ &= -q^{-l}(1 + q^l) y_{l-1} , \end{aligned}$$

where

$$y_l := z^{-l}(1 - z^{-1})(q^{-l}z; q)_{2l+1} = (-1)^l q^{-\frac{l(l+1)}{2}} \prod_{j=0}^l (z - q^j)(z^{-1} - q^j) .$$

From $\mathcal{L}_{Q_0}\sigma = \mathcal{L}_{Q_0}$ it follows that

$$2\mathcal{L}_{Q_0}(B(-l, l)) = \mathcal{L}_{Q_0}((\text{id} + \sigma)B(-l, l)) = -q^{-l}(1 + q^l)\mathcal{L}_{Q_0}(y_{l-1}) ,$$

which is divisible by $2(1 + q^l)(q^l; q)_l = 2(q^{l+1}; q)_l$ thanks to Lemma 2.4 (a). This completes the induction, whence the proof. \square

Lemma 2.4. *With the same notations as above we have*

- (a) *if $f \in \mathbb{Z}[z^{\pm 1}, q^{\pm 1}]$ is invariant under σ , then $2(q^{l+1}; q)_{l+1}$ divides $\mathcal{L}_{Q_0}(fy_l)$;*
- (b) *for any $f \in \mathbb{Z}[z^{\pm 1}, q^{\pm 1}]$, $\mathcal{L}_{Q_0}((q^{-l}z; q)_{2l+1} f)$ is divisible by $(q^{l+1}; q)_{l+1}$.*

Proof. (a) First we prove the case $f = 1$. We will show that this case follows from [Le3, Theorem 7], which was proved by using the Andrews identity. In fact we have

$$\begin{aligned} y_l &= z^{-l}(1 - z^{-1})(q^{-l}z; q)_{2l+1} \\ &= z^{-l}(q^{-l}z; q)_{2l+1} - z^{-l-1}(q^{-l}z; q)_{2l+1} . \end{aligned} \tag{13}$$

It is easy to see that the two terms of the right hand side of (13) are related by

$$\sigma(z^{-l}(q^{-l}z; q)_{2l+1}) = -z^{-l-1}(q^{-l}z; q)_{2l+1} . \tag{14}$$

Hence

$$\begin{aligned} \mathcal{L}_{Q_0}(y_l) &= 2\mathcal{L}_{Q_0}(z^{-l}(q^{-l}z; q)_{2l+1}) \\ &= 2 \sum_{j=0}^{2l+1} (-1)^j \begin{bmatrix} 2l+1 \\ j \end{bmatrix} q^{a_2(j-l)^2} , \end{aligned}$$

which, according to [Le3, Theorem 7], is divisible by

$$2 \frac{\{2l+1\}!}{\{l\}!} = 2(-1)^{l+1} q^{-(l+1)(3l+2)/4} (q^{l+1}; q)_{l+1}$$

in $\mathbb{Z}[q^{\pm 1/2}]$. Since $\mathcal{L}_{Q_0}(y_l)$ and $2(q^{l+1}; q)_{l+1}$ are both in $\mathbb{Z}[q^{\pm 1}]$, (a) is true when $f = 1$.

Consider the general case. Since $\sigma(f) = f$, f is a polynomial in $(z + z^{-1})$. It is enough to prove (a) for $f = (z + z^{-1})^m$. Since

$$(z + z^{-1})y_l = y_{l+1} + (q^{l+1} + q^{-l-1})y_l,$$

$(z + z^{-1})^m y_l$ is a $\mathbb{Z}[q^{\pm 1}]$ -linear combination of $y_l, y_{l+1}, \dots, y_{l+m}$. Because $(q^{l+i}; q)_{l+i}$ is divisible by $(q^{l+1}; q)_{l+1}$ for every positive integer i , the case $f = (z + z^{-1})^m$ follows from the case $f = 1$.

(b) For non-negative integer m , using

$$\sigma(z^m(q^{-l}z; q)_{2l+1}) = -z^{-m-2l-1}(q^{-l}z; q)_{2l+1},$$

we get

$$2\mathcal{L}_{Q_0}(z^m(q^{-l}z; q)_{2l+1}) = \mathcal{L}_{Q_0}((\text{id} + \sigma)z^m(q^{-l}z; q)_{2l+1}) = \mathcal{L}_{Q_0}(y_l \sum_{j=-m-l}^{m+l} z^j),$$

which is divisible by $2(q^{l+1}; q)_{l+1}$ according to (a). Similar argument works for negative m . \square

The following corollary is sometimes more convenient than Theorem 2.2.

Corollary 2.5. *For every positive integer k and every integral quadratic form Q , X_k divides*

$$\sum_{j=0}^k (-1)^j \binom{k}{j}_q q^{Q(j) + \binom{j}{2}}.$$

Proof. From (11) we have

$$\mathcal{L}_Q((z; q)_k) = \sum_{j=0}^k (-1)^j \binom{k}{j}_q q^{Q(j) + \binom{j}{2}}.$$

By Theorem 2.2, the left hand side is divisible by X_k , and so is the right hand side. \square

2.3. Polynomials with q -integer values. We also need a generalization of the following classical result in the theory of polynomials with integer values: If $f(z_1, \dots, z_n) \in \mathbb{Q}[z_1, \dots, z_n]$ takes integer values whenever z_1, \dots, z_n are integers, then f is a \mathbb{Z} -linear combination of $\prod_{i=1}^n \binom{z_i}{k_i}$, $k_i \in \mathbb{Z}_{\geq 0}$.

Let us formulate a q -analog of this fact.

Proposition 2.6. *If $f(z_1, \dots, z_n) \in \mathbb{Q}(q)[z_1, \dots, z_n]$ satisfies $f(q^{m_1}, \dots, q^{m_n}) \in \mathbb{Z}[q^{\pm 1}]$ for every $m_1, \dots, m_n \in \mathbb{Z}$, then f is a $\mathbb{Z}[q^{\pm 1}]$ -linear combination of*

$$\prod_{i=1}^n \frac{(z_i; q)_{k_i}}{(q; q)_{k_i}} \quad \text{with } k_i \in \mathbb{Z}_{\geq 0}.$$

Proof. The elements $z_{\mathbf{k}} := \prod_{i=1}^n (z_i; q)_{k_i} / (q; q)_{k_i}$, with $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}_{\geq 0}^n$, form a $\mathbb{Q}(q)$ -basis of $\mathbb{Q}(q)[x_1, \dots, x_n]$. Hence there are $c_{\mathbf{k}} \in \mathbb{Q}(q)$ such that $f = \sum_{\mathbf{k} \in \mathbb{Z}_{\geq 0}^n} c_{\mathbf{k}} z_{\mathbf{k}}$. Only a finite number of $c_{\mathbf{k}}$'s are non-zero. We will show that $c_{\mathbf{k}} \in \mathbb{Z}[q^{\pm 1}]$ by induction on $|\mathbf{k}| := k_1 + \dots + k_n$.

Suppose $\mathbf{k} = 0$. Let $z_1 = z_2 = \dots = z_n = 1$, then $z_{\mathbf{k}} = 0$ unless $\mathbf{k} = 0$. Hence $c_0 = f(1, 1, \dots, 1) \in \mathbb{Z}[q^{\pm 1}]$.

Suppose $c_{\mathbf{k}} \in \mathbb{Z}[q^{\pm 1}]$ for $|\mathbf{k}| < l$. The $z_{\mathbf{k}}$'s with $|\mathbf{k}| < l$ will be called terms of lower orders. Consider a $\mathbf{k} = (k_1, \dots, k_n)$ with $|\mathbf{k}| = l$. Note that when $z_i = q^{-k_i}$, $z_{(a_1, \dots, a_n)} = 0$ if for some i one has $a_i > k_i$, and $z_{\mathbf{k}} = \pm 1$. Hence

$$f(q^{-k_1}, \dots, q^{-k_n}) = \pm c_{\mathbf{k}} + \text{terms of lower orders.}$$

By induction, the terms of lower orders are in $\mathbb{Z}[q^{\pm 1}]$. Since the left hand side is in $\mathbb{Z}[q^{\pm 1}]$, we conclude that $c_{\mathbf{k}} \in \mathbb{Z}[q^{\pm 1}]$. \square

Corollary 2.7. *For any integer a , the element $(q^a z_1 z_2; q)_k$ is a $\mathbb{Z}[q^{\pm 1}]$ -linear combination of terms*

$$\frac{(q; q)_k}{(q; q)_{k_1} (q; q)_{k_2}} (z_1; q)_{k_1} (z_2; q)_{k_2}$$

with $k_1, k_2 \leq k$.

Proof. The evaluation of

$$\frac{(q^a z_1 z_2; q)_k}{(q; q)_k} \in \mathbb{Q}(q)[z_1, z_2]$$

at $z_i = q^{m_i}$ belongs to $\mathbb{Z}[q^{\pm 1}]$ for any m_i . Applying Proposition 2.6 we get the result. Note that k_1 and k_2 should be less than or equal to k by degree reason. \square

3. BASIC RESULTS: THE CASE OF ROOTS OF UNITY

In this section we prove a basic divisibility result for the case when q is a root of unity ξ of order r . In Subsection 3.4 we reduce the integrality of $H^G(k, b, \varepsilon)/H^G(0, \pm 1, 0)$ to that of a simpler quotient.

For $x, y \in \mathbb{Q}(\xi^{1/4}, e_8)$ we write $x \sim y$ if x/y is an invertible element in $\mathbb{Z}[\xi^{1/4}, e_8]$.

We use the notation $\bar{k} = r - 1 - k$, and

$$\binom{m}{n}_{\xi} := \text{ev}_{\xi} \binom{m}{n}_q, \quad O_{\xi} := (\xi; \xi)_{\lfloor \frac{r-1}{2} \rfloor}, \quad x_k := \prod_{j=\lfloor k/2 \rfloor + 1}^k (1 - \xi^j) = \text{ev}_{\xi}(X_k),$$

where X_k is defined in (12).

3.1. Divisibility. The main divisibility result at roots of unity is formulated below.

Proposition 3.1. *For every integral quadratic form Q and $f(z, q) \in I_k$ with $0 \leq k < r$ we have*

$$\sum_{n=0}^{r-1} \xi^{Q(n)} f(\xi^n, \xi) \in x_k O_{\xi} \cdot \mathbb{Z}[\xi].$$

We need the following lemma for the proof of Proposition 3.1.

Lemma 3.2. *For any integers a, k , with $0 \leq k < r$, and integral quadratic form Q , the element*

$$y = \sum_{n=0}^{r-1} \xi^{Q(n)} \binom{n+a}{k}_{\xi}$$

is divisible by $x_{\bar{k}}$.

Proof. Using $1 - \xi^m = -\xi^m(1 - \xi^{r-m})$ we have

$$\begin{aligned} \binom{n+k}{n}_\xi &= \frac{(\xi^{k+1}; \xi)_n}{(\xi; \xi)_n} = (-1)^n \xi^{kn+n(n+1)/2} \frac{(\xi^{r-k-n}; \xi)_n}{(\xi; \xi)_n} \\ &= (-1)^n \xi^{nk+n+\binom{n}{2}} \binom{\bar{k}}{n}_\xi. \end{aligned} \quad (15)$$

Set $n' = n + a - k$. One has

$$\begin{aligned} y &= \sum_{n=0}^{r-1} \xi^{Q(n)} \binom{n'+k}{k}_\xi \\ &= \sum_{n'=0}^{r-1} (-1)^{n'} \xi^{Q'(n')+\binom{n'}{2}} \binom{\bar{k}}{n'}_\xi \end{aligned} \quad (16)$$

by (15), where $Q'(n') = Q(n' - a + k) + n'k + n'$. In the right hand side of (16), the index n' actually runs from 0 to $\bar{k} - 1$, since $\binom{\bar{k}}{n'}_\xi = 0$ if $n' \geq \bar{k}$. The right hand side of (16) is divisible by $x_{\bar{k}}$ by Corollary 2.5. \square

Proof of Proposition 3.1. Since the set $\{z^d(zq^a; q)_k : d, a \in \mathbb{Z}\}$ spans I_k over $\mathbb{Z}[q^{\pm 1}]$, we can assume that $f = z^d(zq^a; q)_k$. Then

$$\begin{aligned} \frac{\sum_{n=0}^{r-1} \xi^{Q(n)} f(\xi^n, \xi)}{x_k O_\xi} &= \frac{(\xi; \xi)_{\lfloor k/2 \rfloor}}{O_\xi} \sum_{n=0}^{r-1} \xi^{Q(n)+dn} \binom{n+a-1}{k}_\xi \\ &\in \frac{(\xi; \xi)_{\lfloor k/2 \rfloor}}{O_\xi} x_{\bar{k}} \mathbb{Z}[\xi] \quad (\text{by Lemma 3.2}), \end{aligned}$$

which is in $\mathbb{Z}[\xi]$ by Lemma 3.3 (f) below. \square

3.2. The ring $\mathbb{Z}[\xi]$. It is known that $\mathbb{Z}[\xi]$ is a Dedekind domain with field of fractions $\mathbb{Q}[\xi]$.

Lemma 3.3. *a) If $(a, r) = (b, r)$ then $(1 - \xi^a) \sim (1 - \xi^b)$ in $\mathbb{Z}[\xi]$.*

b) One has $(\xi; \xi)_{r-1} = r$.

c) Suppose $y \in \mathbb{Q}[\xi]$ and $y^s \in \mathbb{Z}[\xi]$ for some positive integer s . Then $y \in \mathbb{Z}[\xi]$.

d) Suppose $y, z \in \mathbb{Q}[\xi]$ with $z \neq 0$. If $a_s := y^s z \in \mathbb{Z}[\xi]$ for infinitely many positive s , then $y \in \mathbb{Z}[\xi]$.

e) One has

$$O_\xi^2 \sim \begin{cases} r & \text{if } r \text{ is odd,} \\ r/2 & \text{if } r \text{ is even.} \end{cases}$$

f) For every integer $0 \leq k < r$, O_ξ divides $(\xi; \xi)_{\lfloor k/2 \rfloor} x_{\bar{k}}$.

Proof. a) Let $c = (a, r) = (b, r)$. Since $1 - \xi^c$ divides $1 - \xi^a$, and also $1 - \xi^a$ divides $1 - \xi^c = 1 - \xi^{aa^*}$ where $aa^* \equiv c \pmod{r}$, we have $1 - \xi^a \sim 1 - \xi^c$. Similarly $1 - \xi^b \sim 1 - \xi^c$.

Part b) is obtained by substituting $X = 1$ into the following identity.

$$1 + X + \cdots + X^{r-1} = \frac{1 - X^r}{1 - X} = \prod_{i=1}^{r-1} (X - \xi^i).$$

Part c) follows from the fact that every Dedekind domain is integrally closed.

d) Let $y = y_1/y_2$ and $z = z_1/z_2$ with $y_1, y_2, z_1, z_2 \in \mathbb{Z}[\xi]$ and $z_i \neq 0$. Then for infinitely many $s > 0$, $z_1 y_1^s = a_s z_2 y_2^s$, and hence

$$(z_1)(y_1)^s = (a_s)(z_2)(y_2)^s, \quad (17)$$

where (x) denotes the principal ideal in $\mathbb{Z}[\xi]$ generated by x . In any Dedekind domain, every ideal decomposes uniquely into a product of prime ideals:

$$(x) = \prod_i \mathfrak{p}_i^{e_i}$$

and this decomposition respects the multiplication. From the uniqueness of prime ideal decomposition and (17), we see easily that $y_2 \mid y_1$, or $y = y_1/y_2 \in \mathbb{Z}[\xi]$.

e) First suppose r is odd. Then $O_\xi = (\xi; \xi)_{\frac{r-1}{2}}$. Since $(1 - \xi^k) \sim (1 - \xi^{r-k})$ by part (a), we have

$$O_\xi^2 \sim (\xi; \xi)_{r-1} = r.$$

Now suppose r is even. Then $O_\xi = (\xi; \xi)_{\frac{r-2}{2}}$. Using $(1 - \xi^k) \sim (1 - \xi^{r-k})$, we have

$$O_\xi^2 \sim (\xi; \xi)_{r-1}/(1 - \xi^{r/2}) = r/2$$

since $\xi^{r/2} = -1$.

f) First suppose r is odd. Note that for odd r , $(1 - \xi^j) \sim (1 - \xi^{2j})$ by part (a). One has

$$\begin{aligned} x_{\bar{k}} = x_{r-k-1} &= \frac{\prod_{j=1}^{r-k-1} (1 - \xi^j)}{\prod_{j=1}^{\lfloor \frac{r-k-1}{2} \rfloor} (1 - \xi^j)} \\ &\sim \frac{\prod_{j=1}^{r-k-1} (1 - \xi^j)}{\prod_{j=1}^{\lfloor \frac{r-k-1}{2} \rfloor} (1 - \xi^{2j})} \quad \text{since } (1 - \xi^j) \sim (1 - \xi^{2j}) \\ &\sim (1 - \xi)(1 - \xi^3) \cdots (1 - \xi^{r-2-2\lfloor \frac{k}{2} \rfloor}). \end{aligned} \quad (18)$$

Using $(1 - \xi^j) \sim (1 - \xi^{2j}) \sim (1 - \xi^{r-2j})$, we have

$$(\xi; \xi)_{\lfloor \frac{k}{2} \rfloor} = \prod_{j=1}^{\lfloor \frac{k}{2} \rfloor} (1 - \xi^j) \sim \prod_{j=1}^{\lfloor \frac{k}{2} \rfloor} (1 - \xi^{r-2j}). \quad (19)$$

Multiplying (18) and (19), we get

$$(\xi; \xi)_{\lfloor \frac{k}{2} \rfloor} x_{\bar{k}} \sim \prod_{j=1}^{(r-1)/2} (1 - \xi^{2j-1}) \sim O_\xi, \quad (20)$$

where the second \sim follows from the fact that $1 - \xi^{r-a} \sim 1 - \xi^a$ for any integer a .

Now suppose r is even.

$$\begin{aligned} \frac{O_\xi}{(\xi; \xi)_{\lfloor \frac{k}{2} \rfloor}} &= (1 - \xi^{r/2-1})(1 - \xi^{r/2-2}) \cdots (1 - \xi^{\lfloor \frac{k}{2} \rfloor+1}) \\ &= (1 + \xi)(1 + \xi^2) \cdots (1 + \xi^{\frac{r}{2}-1-\lfloor \frac{k}{2} \rfloor}). \end{aligned} \quad (21)$$

On the other hand there exists $f \in \mathbb{Z}[\xi]$ such that

$$\begin{aligned} x_{\bar{k}} = x_{r-k-1} &= \frac{\prod_{j=1}^{r-k-1} (1 - \xi^j)}{\prod_{j=1}^{\lfloor \frac{r-k-1}{2} \rfloor} (1 - \xi^j)} \\ &= f \frac{\prod_{j=1}^{\lfloor \frac{r-k-1}{2} \rfloor} (1 - \xi^{2j})}{\prod_{j=1}^{\lfloor \frac{r-k-1}{2} \rfloor} (1 - \xi^j)} \\ &= f (1 + \xi)(1 + \xi^2) \cdots (1 + \xi^{\lfloor \frac{r-k-1}{2} \rfloor}). \end{aligned} \quad (22)$$

Note that $\lfloor \frac{r-k-1}{2} \rfloor = \frac{r}{2} - 1 - \lfloor \frac{k}{2} \rfloor$ for even r . Compare (21) and (22), we see that O_ξ divides $(\xi; \xi)_{\lfloor \frac{k}{2} \rfloor} x_{\bar{k}}$. \square

3.3. Quadratic Gauss sums. For arbitrary integers b and d , the quadratic Gauss sum is defined as

$$G(b, d, \xi) := \sum_{n=0}^{\text{ord}(\xi)-1} \xi^{bn^2+dn}.$$

The following is well-known.

Proposition 3.4. *a) Let $r = \text{ord}(\xi)$ and $c = \text{gcd}(b, r)$. Then*

$$G(b, d, \xi) = \begin{cases} c G(b/c, d/c, \xi^c) & \text{if } c \mid d; \\ 0 & \text{otherwise.} \end{cases}$$

b) Suppose b and r are co-prime. Then

$$G(b, 0, \xi)^2 \sim \begin{cases} r & \text{if } r \text{ is odd;} \\ 0 & \text{if } r \equiv 2 \pmod{4}; \\ 2r & \text{if } r \equiv 0 \pmod{4}. \end{cases}$$

Furthermore $G(b, b, \xi)^2 = 2r$ if $r \equiv 2 \pmod{4}$.

c) Suppose d is odd and $r \equiv 0 \pmod{4}$. Then $G(b, d, \xi) = 0$.

d) Suppose r_1 and r_2 are co-prime and $r = r_1 r_2$. Then

$$G(b, d, \xi) = G(br_1, d, \xi^{r_1}) G(br_2, d, \xi^{r_2}). \quad (23)$$

Proof. Part (a) is clear from the definition when $c \mid d$. Now suppose that $c \nmid d$. We have

$$\begin{aligned} G(b, d, \xi) &= \sum_{t=0}^{r/c-1} \sum_{s=0}^{c-1} \xi^{b(sr/c+t)^2 + d(sr/c+t)} \\ &= \sum_{t=0}^{r/c-1} \xi^{bt^2+dt} \sum_{s=0}^{c-1} \xi^{sdr/c} = 0, \end{aligned}$$

where the last equality follows from the fact that $\xi^{dr/c} \neq 1$ and its order divides c .

b) After a Galois transformation of the form $\xi \rightarrow \xi^a$, with a co-prime to r , one can assume that $b = 1$ and $\xi = \exp(2\pi i/r)$. The result now follows e.g. from [D, Chapter 2].

c) One has

$$\begin{aligned} b \left(n + \frac{r}{2} \right)^2 + d \left(n + \frac{r}{2} \right) &= bn^2 + bnr + br \frac{r}{4} + dn + d \frac{r}{2} \\ &\equiv bn^2 + dn + \frac{r}{2} \pmod{r}. \end{aligned}$$

Hence

$$G(b, d, \xi) = \sum_{n=0}^{r-1} \xi^{bn^2+dn} = \sum_{n=0}^{r-1} \xi^{b(n+r/2)^2+d(n+r/2)} = \xi^{r/2} \sum_{n=0}^{r-1} \xi^{bn^2+dn} = -G(b, d, \xi).$$

It follows that $G(b, d, \xi) = 0$.

d) The proof follows easily from the fact that the map $\mathbb{Z}/r_1 \times \mathbb{Z}/r_2 \rightarrow \mathbb{Z}/(r_1 r_2)$, defined by $(n_1, n_2) \rightarrow r_2 n_1 + r_1 n_2$, is an isomorphism. \square

Proof of Lemma 1.2. Now we are in position to see that $F_{U^+}^G(\xi) = 0$ if and only if $G = SU(2)$ and $\xi^{1/4}$ has order $2r$.

By completing the squares we have

$$(1 - \xi)F_{U^+}^G \sim \frac{2}{1 - \xi} \sum_n^{\xi, G} q^{\frac{n^2-1}{4}} (1 - q^n) \sim \frac{2}{1 - \xi} \left(\sum_n^{\xi, G} q^{\frac{n^2-1}{4}} - \xi^{-1} \sum_n^{\xi, G} q^{\frac{(n+2)^2-1}{4}} \right)$$

$$\sim \begin{cases} G(4^*, 0, \xi) & \text{if } G = SO(3) ; \\ \frac{1}{2}G(1, 0, \xi^{1/4}) & \text{if } G = SU(2) \text{ and } \text{ord}(\xi^{1/4}) = 4r ; \\ G(1, 0, \xi^{1/4}) & \text{if } G = SU(2) \text{ and } \text{ord}(\xi^{1/4}) = 2r ; \\ 2G(1, 0, \xi^{1/4}) & \text{if } G = SU(2) \text{ and } \text{ord}(\xi^{1/4}) = r . \end{cases}$$

Note that for $G = SO(3)$, the sum is over odd n 's, so $n^2 - 1$ is always divisible by 4. Hence, for any choice of $\xi^{1/4}$, we have $\xi^{(n^2-1)/4} = \xi^{4^*(n^2-1)}$ with $4^*4 = 1 \pmod{r}$.

If r is even, then $\text{ord}(\xi^{1/4})$ is always $4r$. Now Proposition 3.4 (b) implies the claim. \square

3.4. Simplification of $H^G(k, b, \varepsilon)$.

Lemma 3.5. a) For integers k, b , and $\varepsilon \in \{0, 1\}$, there is $f_\varepsilon(z, q) \in I_{2k+1+\varepsilon}$ such that

$$H^G(k, b, \varepsilon) \sim \frac{2}{x_{2k+1+\varepsilon}} \sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4} - \frac{3\varepsilon n}{2}} f_\varepsilon(q^n, q) .$$

More precisely, one can choose $f_\varepsilon = z^{-k} (q^{-k} z; q)_{2k+1+\varepsilon}$.

b) One has $\sqrt{2}, \sqrt{r} \in \mathbb{Z}[\xi^{1/4}, e_8]$ and

$$H^G(0, \pm 1, 0) \sim \begin{cases} \sqrt{r} & \text{if } G = SO(3) ; \\ \sqrt{2r} & \text{if } G = SU(2) \text{ and } \text{ord}(\xi^{1/4}) = 4r ; \\ 2\sqrt{r} & \text{if } G = SU(2) \text{ and } \text{ord}(\xi^{1/4}) = r . \end{cases}$$

c) One has $\mathcal{D}^G \in \mathbb{Z}[\xi^{1/4}, e_8]$ and $(1 - \xi)\mathcal{D}^G \sim H^G(0, \pm 1, 0)$.

d) Suppose b and r are even. Then $H^{SU(2)}(k, b, 1) = 0$.

Proof. a) We will use the following simple observation: For $g(z, q) \in \mathbb{Q}[z^{\pm 1/2}, q^{\pm 1/4}]$, we have

$$\sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4}} g(q^{n/2}, q) = \sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4}} g(q^{-n/2}, q). \quad (24)$$

To prove it, one only needs to consider $g(z, q) = z^{a/2}, a \in \mathbb{Z}$. Then

$$\text{LHS} = \sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4} + \frac{an}{2}} = \sum_{n \rightarrow -n}^{\xi, G} q^{\frac{b(n^2-1)}{4} - \frac{an}{2}} = \text{RHS} .$$

One can check that $\{n\} \prod_{j=-k}^k \{n+j\} = (q^{-kn-n} - q^{-kn})(q^{n-k}; q)_{2k+1}$. Then we get

$$\begin{aligned} \text{ev}_\xi \left(\frac{\{2k+1\}!}{\{k\}!} \right) H^G(k, b, 0) &= \sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4}} \{n\} \prod_{j=-k}^k \{n+j\} \\ &= -2 \sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4}} q^{-nk} (q^{n-k}; q)_{2k+1}, \end{aligned}$$

where the last equality follows from (14), (24) and the fact that

$$q^{-kn-n}(q^{n-k}; q)_{2k+1} = -q^{-kn}(q^{n-k}; q)_{2k+1} \Big|_{n \rightarrow -n}.$$

Analogously, we have

$$\begin{aligned} \text{ev}_\xi \left(\frac{\{2k+1\}!}{\{k\}!} \lambda_{k+1} \right) H^G(k, b, 1) &= \sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4}} \{n\} \lambda_n \prod_{j=-k}^k \{n+j\} \\ &= -2 \sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4}} q^{-n(k+3/2)} (q^{n-k}; q)_{2k+2}. \end{aligned}$$

This proves a).

b) Let us first show that $\sqrt{r}, \sqrt{2} \in \mathbb{Z}[\xi^{1/4}, e_8]$. Observe that $\sqrt{2} \in \mathbb{Z}[e_8]$. Further,

$$\prod_{j=1}^{(r-1)/2} |1 - \xi^j| = \begin{cases} \sqrt{r} & \text{if } r \text{ is odd;} \\ \sqrt{r/2} & \text{if } r \text{ is even.} \end{cases}$$

Since $|1 - \xi^j| = \pm \sqrt{-1} (\xi^{j/2} - \xi^{-j/2})$, we have $\sqrt{r} \in \mathbb{Z}[\xi^{1/4}, e_4]$.

Part (b) follows now from (9), Proposition 3.4 (b) and the proof of Lemma 1.2.

c) Since $\mathcal{D}_G := |F_{U^+}^G|$, from the proof of Lemma 1.2, we get

$$|1 - \xi| \mathcal{D}^G = \begin{cases} \sqrt{r} & \text{if } G = SO(3); \\ \sqrt{2r} & \text{if } G = SU(2) \text{ and } \text{ord}(\xi^{1/4}) = 4r; \\ 2\sqrt{r} & \text{if } G = SU(2) \text{ and } \text{ord}(\xi^{1/4}) = r. \end{cases}$$

Clearly, \sqrt{r} is divisible by $|1 - \xi|$, so $\mathcal{D}^G \in \mathbb{Z}[\xi^{1/4}, e_8]$. The second statement follows from (b).

d) By part (a), it is enough to show that

$$\sum_n^{\xi, G} q^{\frac{b(n^2-1)}{4} - \frac{3n}{2}} f(q^n, q) = 0$$

for any $f \in \mathbb{Z}[z^{\pm 1}, q^{\pm 1}]$. We can assume $f = z^a$, $a \in \mathbb{Z}$. Assume $b = 2b'$. Then

$$4 \sum_n^{\xi, SU(2)} q^{\frac{b(n^2-1)}{4} - \frac{3n}{2}} q^{na} = 2\xi^{-b'/2} \sum_{n=0}^{2r-1} (\xi^{1/2})^{b'n^2 - 3n + 2na} = 2\xi^{-b'/2} G(b', 2a - 3, \xi^{1/2}),$$

which is 0 by Proposition 3.4 (c), since $\text{ord}(\xi^{1/2})$ is always $2r$ if r is even. \square

3.5. Lens spaces. Suppose $L \sqcup L'$ is the Hopf link with framing $b \neq 0$ on L and framing 0 on L' . Besides, the color of L' is a fixed number a . By surgery on L from (S^3, L') we get the pair $(\mathbf{L}(b, 1), L')$, where $\mathbf{L}(b, 1)$ is the lens space. It is known that $J_{L \sqcup L'}(n) = q^{b(n^2-1)/4} [na]$. Hence we have

$$\tau_{\mathbf{L}(b,1),L'}^G(\xi) = \frac{\sum_n \xi^{,G} q^{b(n^2-1)/4} [na][n]}{\sum_n \xi^{,G} q^{\text{sn}(b)(n^2-1)/4} [n]^2}. \quad (25)$$

Note that the invariant of $\mathbf{L}(b, -1) = \mathbf{L}(-b, 1)$ is just a complex conjugate of (25).

Lemma 3.6. *a) If b and r are co-prime, then $\tau_{\mathbf{L}(b,1)}^{SO(3)}$ is invertible in $\mathbb{Z}[\xi]$.*

b) Suppose r is even. For $b = 2^k$, there is a knot K in the lens space $M = \mathbf{L}(2^k, -1)$ colored by an odd number such that

$$\tau_{M,K}^{SU(2)}(\xi) \neq 0.$$

Proof. a) The $SO(3)$ invariant of $\mathbf{L}(b, 1)$ can be easily computed. By completing the square we have

$$\tau_{\mathbf{L}(b,1)}^{SO(3)}(\xi) = \xi^{(\text{sn}(b)-b)/4} \frac{(1 - \xi^{-b^*})}{(1 - \xi^{-1})} \frac{G(b, 0, \xi)}{G(1, 0, \xi)},$$

which is a unit in $\mathbb{Z}[\xi^{1/4}]$ by Proposition 3.4 (b). Here $b^*b \equiv 1 \pmod{r}$.

b) Let $L \sqcup L'$ be the Hopf link with framing $-b = -2^k$ on L and framing 0 on L' . Suppose L' is colored by $a = 2s + 1$. Surgery on L gives us a pair $(M, K) = (\mathbf{L}(2^k, -1), K)$.

An easy calculation shows

$$\tau_{M,K}^{SU(2)}(\xi) \sim \frac{G(-b, 4s + 4, \xi^{1/4}) - G(-b, 4s, \xi^{1/4})}{(1 - \xi) G(-1, 0, \xi^{1/4})}. \quad (26)$$

For $b = 2$, then $M = \mathbb{R}P^3$. Choose $s = 0$, or $a = 1$. Then $\tau_{M,K}^{SU(2)}(\xi) = \tau_M^{SU(2)}(\xi) \neq 0$.

For $b = 4$ again choose $s = 0$. Then one and only one term in the numerator of (26) is zero, by Proposition 3.4.

Suppose $b = 2^k > 4$. Then $c := (b, 4r) > 4$. Choose s such that $G(-b, 4s, \xi^{1/4}) \neq 0$ (see Proposition 3.4). Then $c \mid 4s$, and c does not divide $4s + 4$. Hence $G(-b, 4s + 4, \xi^{1/4}) = 0$. We conclude that $\tau_{M,K}^{SU(2)}(\xi) \neq 0$. \square

4. SYMMETRY PRINCIPLE AND SPLITTING OF THE $SU(2)$ INVARIANT

The symmetry principle of the colored Jones polynomial and the splitting of the $SU(2)$ WRT invariant were discovered by Kirby and Melvin in [KM]. In [Le1, Le3], the third author generalized these to all higher ranked Lie groups. Here we extend the symmetry principle and splitting to the case of pairs of a 3-manifold and a colored link inside. We show that the symmetry principle for a link in an arbitrary 3-manifold holds only for $SO(3)$ invariant, but does not hold for the $SU(2)$ invariant.

4.1. Symmetry Principle for links in S^3 . Suppose ξ is a root of unity of order r . Then the colored Jones polynomial at ξ is periodic with period $2r$, i.e.

$$\text{ev}_\xi (J_L(n_1, \dots, n_i + 2r, \dots, n_m)) = \text{ev}_\xi (J_L(n_1, \dots, n_i, \dots, n_m)), \quad (27)$$

and under the reflection $r - n \rightarrow r + n$ it behaves as follows:

$$\text{ev}_\xi (J_L(n_1, \dots, r + n_i, \dots, n_m)) = -\text{ev}_\xi (J_L(n_1, \dots, r - n_i, \dots, n_m)) \quad (28)$$

(see [Le1]). This means that one can restrict the colors to the interval $[0, r]$.

The symmetry principle tells us how J_L behaves under the transformation $n \rightarrow r - n$. More precisely, let $\mathbb{Z}/2 = \{0, 1\}$ act on $\mathbb{Z}/r\mathbb{Z}$ by $0 * n = n$, $1 * n = r - n$. For $\mathbf{a} =$

$(a_1, \dots, a_m) \in \{0, 1\}^m$ and $\mathbf{n} = (n_1, \dots, n_m) \in (\mathbb{Z}/r\mathbb{Z})^m$, let $\mathbf{a} * \mathbf{n} = (a_1 * n_1, \dots, a_m * n_m)$. In addition, we set $\hat{n} := n - 1$ for any integer $n \in \mathbb{Z}$.

Proposition 4.1. *Suppose (ℓ_{ij}) is the linking matrix of L . With the notations as above one has*

$$\text{ev}_\xi(J_L(\mathbf{a} * \mathbf{n})) = (-\xi^{r/2})^{\sum_i a_i} \xi^t \text{ev}_\xi(J_L(\mathbf{n})),$$

where

$$t = \frac{r(r-2)}{4} \sum_{i,j} \ell_{ij} a_i a_j + \frac{r}{2} \sum_{i,j} \ell_{ij} a_i \hat{n}_j, \quad (29)$$

and (ℓ_{ij}) is the linking matrix of L .

Proof. This is the sl_2 case of [Le1, Theorem 2.6]. The factor $(-\xi^{r/2})^{\sum_i a_i}$ comes from the difference between our J_L and Q_L in [Le1], where Q_L is equal to J_L times the quantum dimensions of the colors on L . \square

Remark 4.2. If $\text{ord}(\xi^{1/2}) = 2r$, then $-\xi^{r/2} = 1$, and this case was considered in [KM]. Proposition 4.1 handles also the case when $\text{ord}(\xi^{1/2}) \neq 2r$, i.e. $\text{ord}(\xi^{1/2}) = r$.

A simple but useful observation is that if all entries of \mathbf{n} are odd, then the second term in (29) is an integer multiple of r , hence can be removed.

4.2. WRT $SO(3)$ invariant for an arbitrary colored link in M . In the literature, the WRT $SO(3)$ invariant of the pair (M, L') was defined in the case when all colors of L' are odd or all equal to 2 (compare [MR]). Here we extend this definition to arbitrary colors. Since the colors of L' will play an important role in this section, we will make the dependence on them explicit in the notation.

Note that $SO(3)$ invariants of M with evenly colored links inside are not coming from Topological Quantum Field Theories. The main reason is that fusion preserves odd colors. However, fusion of an odd and an even color produce an even color. This violates the invariance of (6) under sliding in the case when some of the s_i 's are even. We will show that this defect can easily be taken into account by a simple factor depending on the linking matrix and parity of the colors only.

Throughout the remaining of this section let $r = \text{ord}(\xi)$ be odd and $\mathbf{s} = (s_1, \dots, s_l)$ be the color on L' . Let (ℓ_{ij}) and (p_{ij}) be the linking matrices of L and L' respectively. The linking number between the i -th component of L and the j -th component of L' will be denoted by $\hat{\ell}_{ij}$.

Let

$$F_{L \sqcup L'}^{SO(3)}(\xi; \mathbf{s}) := \xi^{\mu(L'; \mathbf{s})} \sum_{n_1, \dots, n_m} \xi^{SO(3)}[\mathbf{n}] J_{L \sqcup L'}(\mathbf{n}, \mathbf{s}), \quad (30)$$

where $[\mathbf{n}] := \prod_{i=1}^m [n_i]$ and

$$\mu(L'; \mathbf{s}) := -\frac{r(r-2)}{4} \sum_{i,j=1}^l p_{ij} \hat{s}_i \hat{s}_j.$$

Observe that, when all s_i are odd, (30) coincides with (5).

Lemma 4.3. $F_{L \sqcup L'}^{SO(3)}(\xi; \mathbf{s})$ is invariant under the handle slide of a component of L or L' over a component of L .

Proof. The invariance under sliding of one component of L over another component of L follows by standard arguments (see e.g. [Lic]).

Let $L \sqcup L''$ be the link obtained from $L \sqcup L'$ by sliding a component of L' over a component of L . It is enough to show that

$$F_{L \sqcup L'}^{SO(3)}(\xi; \mathbf{s}) = F_{L \sqcup L''}^{SO(3)}(\xi; \mathbf{s}).$$

Using the fact that $\hat{s} * \hat{s} * s = s$ for any s with $\hat{s} \equiv s - 1 \pmod{2}$ and Proposition 4.1, we have

$$\xi^{\mu(L', \mathbf{s})} J_{L \sqcup L'}(\mathbf{n}, \mathbf{s}) = (-\xi^{r/2})^{\sum_i \hat{s}_i} J_{L \sqcup L'}(\mathbf{n}, \hat{\mathbf{s}} * \mathbf{s}). \quad (31)$$

Here we used the fact $\hat{s} * s$ is always odd, and hence all summands of $\sum p_{ij} \widehat{\hat{s}_i \hat{s}_j * s_j}$ are even.

By the invariance of $F_{L \sqcup L'}^{SO(3)}(\xi, \mathbf{s})$ in the standard case when all colors are odd, we get

$$\sum_{n_1, \dots, n_m} \xi^{SO(3)}[\mathbf{n}] J_{L \sqcup L'}(\mathbf{n}, \hat{\mathbf{s}} * \mathbf{s}) = \sum_{n_1, \dots, n_m} \xi^{SO(3)}[\mathbf{n}] J_{L \sqcup L''}(\mathbf{n}, \hat{\mathbf{s}} * \mathbf{s}). \quad (32)$$

Further using Proposition 4.1 again, we obtain

$$J_{L \sqcup L''}(\mathbf{n}, \hat{\mathbf{s}} * \mathbf{s}) = (-\xi^{r/2})^{\sum_i \hat{s}_i} \xi^{\mu(L'', \mathbf{s})} J_{L \sqcup L''}(\mathbf{n}, \mathbf{s}). \quad (33)$$

Inserting (31), (33) into (32) we get the result. \square

Lemma 4.3 suggests to define $\tau_{M, L'}^{SO(3)}(\xi; \mathbf{s})$ for arbitrary \mathbf{s} by substituting $F_{L \sqcup L'}^{SO(3)}(\xi)$ given by (30) into (6). When all colors of L' are odd, the only additional factor $\xi^{\mu(L', \mathbf{s})}$ is 1 and we get back our old invariant.

Corollary 4.4. $\tau_{M, L'}^{SO(3)}(\xi; \mathbf{s})$ is an invariant of the pair (M, L') .

Remark 4.5. For a colored link L in the 3-sphere, our invariant equals to

$$\tau_{S^3, L}^{SO(3)}(\xi; \mathbf{s}) = \xi^{-r(r-2)/4} \sum_{i,j} l_{ij} \hat{s}_i \hat{s}_j \operatorname{ev}_\xi(J_L(\mathbf{s})).$$

Hence if some colors of L are even, this invariant might differ from the colored Jones polynomial by some factor depending on the linking matrix (l_{ij}) of L .

4.3. Symmetry Principle for the WRT $SO(3)$ invariant. We use the same notations as in the previous section.

Proposition 4.6. For $\mathbf{a} \in \{0, 1\}^l$ and $\mathbf{s} \in (\mathbb{Z}/r\mathbb{Z})^l$ one has

$$\tau_{M, L'}^{SO(3)}(\xi; \mathbf{a} * \mathbf{s}) = (-\xi^{r/2})^{\sum_i a_i} \tau_{M, L'}^{SO(3)}(\xi; \mathbf{s}).$$

Proof. By 4.1 we have

$$\sum_{n_1, \dots, n_m} \xi^{SO(3)}[\mathbf{n}] J_{L \sqcup L'}(\mathbf{n}, \mathbf{a} * \mathbf{s}) = (-\xi^{r/2})^{\sum_i a_i} \xi^u \sum_{n_1, \dots, n_m} \xi^{SO(3)}[\mathbf{n}] J_{L \sqcup L'}(\mathbf{n}, \mathbf{s}),$$

where $u = \frac{r(r-2)}{4} \sum_{i,j} p_{ij} a_i a_j + \frac{r}{2} \sum_{i,j} p_{ij} a_i \hat{s}_j$. Here we use the fact that \mathbf{n} is odd in the above sum. On the other hand

$$w := \mu(L'; \mathbf{s}) - \mu(L'; \mathbf{a} * \mathbf{s}) = \frac{r(r-2)}{4} \sum_{i,j} p_{ij} (\widehat{\widehat{a}_i * s_i \widehat{a}_j * s_j} - \hat{s}_i \hat{s}_j).$$

Then

$$u - w \equiv \frac{r(r-2)}{4} \sum_i p_{ii}(\hat{s}_i^2 - \widehat{a_i * s_i}^2 + 2a_i \hat{s}_i + a_i^2) \\ + \frac{r}{2} \sum_{i < j} p_{ij}(\hat{s}_i \hat{s}_j - \widehat{a_i * s_i a_j * s_j} + a_i \hat{s}_j + a_j \hat{s}_i + a_i a_j) \equiv 0 \pmod{r},$$

which can be verified directly. \square

Remark 4.7. Proposition 4.6 is not true for the WRT $SU(2)$ invariant. For example, consider the Hopf link with framing 2 on the first component and framing 0 on the second. Surgery on the first component produces a pair $(\mathbb{R}P^3, K)$. If $\text{ord}(\xi) = 3$ and $\text{ord}(\xi^{1/4}) = 12$ then $\tau_{M,K}^{SU(2)}(1; \xi) = 0$ and $\tau_{M,K}^{SU(2)}(1 * 1 = 2; \xi) \neq 0$.

4.4. Splitting. In [KM] it was proved that when both the $SO(3)$ and $SU(2)$ WRT invariants can be defined, i.e. when r is odd, then one has the splitting

$$\tau_M^{SU(2)}(\xi) = \tau_M^{\mathbb{Z}/2}(\xi) \tau_M^{SO(3)}(\xi),$$

where $\tau_M^{\mathbb{Z}/2}(\xi)$ is a simple invariant depending only on the linking pairing of M . Here we generalize this result for invariants of pairs $L' \subset M$. We will follow the approach in [Le2], where the splitting is generalized to all higher ranked simple Lie algebras.

Let s_1, \dots, s_l be the colors on L' and set

$$F_{L \sqcup L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) = \xi^{\frac{r(r-2)}{4} \sum p_{ij} \hat{s}_i \hat{s}_j} \sum_{\alpha_1, \dots, \alpha_m \in \{0,1\}} \xi^{\frac{r(r-2)}{4} \sum \ell_{ij} \alpha_i \alpha_j + \frac{r}{2} \sum \varepsilon_i \alpha_i}, \quad (34)$$

where (ℓ_{ij}) and (p_{ij}) are the linking matrices of L and L' respectively, and ε_i is defined by (2).

For example

$$F_{U^\pm}^{\mathbb{Z}/2}(\xi) = 1 + \xi^{\pm \frac{r(r-2)}{4}}. \quad (35)$$

We will assume that $r = \text{ord}(\xi)$ is odd and $\xi^{1/4}$ is chosen so that $\text{ord}(\xi^{1/4}) \neq 2r$, i.e. $\text{ord}(\xi^{1/4})$ is either r or $4r$. This choice guarantees that $F_{U^\pm}^{\mathbb{Z}/2}(\xi) \neq 0$. Define

$$\tau_{M,L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) = \frac{F_{L \sqcup L'}^{\mathbb{Z}/2}(\xi; \mathbf{s})}{\left(F_{U^+}^{\mathbb{Z}/2}(\xi)\right)^{\beta_+} \left(F_{U^-}^{\mathbb{Z}/2}(\xi)\right)^{\beta_-} \left|F_{U^+}^{\mathbb{Z}/2}(\xi)\right|^\beta}. \quad (36)$$

Then $\tau_{M,L'}^{\mathbb{Z}/2}(\xi; \mathbf{s})$ is an invariant of the pair (M, L') .

Remark 4.8. This type of invariants were studied in [MOO, De1] for 3-manifolds without links inside, and in [De2] for 3-manifolds with links inside. When the abelian group is $\mathbb{Z}/2\mathbb{Z}$, set the parameters c_i in [De2] to be equal to $s_i - 1 \pmod{2}$, and define the quadratic form q on $\mathbb{Z}/2\mathbb{Z}$ as follows: $q(0) = 0$, $q(1) = (r-2)/4$, then the invariant introduced in [De2] is equal to $\tau_{M,L'}^{\mathbb{Z}/2}(\xi; \mathbf{s})$ after setting $\xi^{r/4} = \sqrt{-1}$.

Proposition 4.9. *Suppose $r = \text{ord}(\xi)$ is odd, and $\text{ord}(\xi^{1/4})$ is either r or $4r$.*

(a) *One has the splitting*

$$\tau_{M,L'}^{SU(2)}(\xi; \mathbf{s}) = \tau_{M,L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) \tau_{M,L'}^{SO(3)}(\xi; \mathbf{s}).$$

(b) *If $\text{ord}(\xi^{1/4}) = r$, then $\tau_{M,L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) = 1$ and*

$$\tau_{M,L'}^{SU(2)}(\xi; \mathbf{s}) = \tau_{M,L'}^{SO(3)}(\xi; \mathbf{s}).$$

(c) One has the integrality

$$\tau_{M,L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) \in \mathbb{Z}[\xi^{1/4}, e_8] .$$

Proof. (a) Recall that $\tilde{\ell}_{ij}$ is the linking number between the i -th component of L and the j -th component of L' . Also note that $\forall \mathbf{a} \in (\mathbb{Z}/2\mathbb{Z})^m$, $\mathbf{a} * (\mathbf{a} * \mathbf{n}) = \mathbf{n}$. By Proposition 4.1 we have

$$\text{ev}_\xi([\mathbf{n}]J_{L \sqcup L'}(\mathbf{n}, \mathbf{s})) = \xi^t \text{ev}_\xi([\mathbf{a} * \mathbf{n}]J_{L \sqcup L'}(\mathbf{a} * \mathbf{n}, \mathbf{s})) ,$$

where $t = \frac{r(r-2)}{4} \sum \ell_{ij} a_i a_j + \frac{r}{2} \sum \tilde{\ell}_{ij} a_i \hat{s}_j$. Note that the factor $(-\xi^{r/2})^{\sum a_i}$ is missing because of the quantum integers. Therefore by (5), (30) and (36) we have

$$\begin{aligned} F_{L \sqcup L'}^{SU(2)}(\xi; \mathbf{s}) &= \xi^{\frac{r(r-2)}{4} \sum p_{ij} \hat{s}_i \hat{s}_j} \sum_{\mathbf{a}_i \in \{0,1\}} \xi^{\frac{r(r-2)}{4} \sum \ell_{ij} a_i a_j + \frac{r}{2} \sum \tilde{\ell}_{ij} a_i \hat{s}_j} F_{L \sqcup L'}^{SO(3)}(\xi; \mathbf{s}) \\ &= F_{L \sqcup L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) F_{L \sqcup L'}^{SO(3)}(\xi; \mathbf{s}) , \end{aligned}$$

which implies (a).

(b) If $\text{ord}(\xi^{1/4}) = r$, then by (34), $F_{L \sqcup L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) = 2^m$. In particular, $F_{U^\pm}^{\mathbb{Z}/2}(\xi; \mathbf{s}) = 2$. It follows that $\tau_{M,L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) = 1$.

(c) The case $\text{ord}(\xi^{1/4}) = r$ was covered by (b). Assume that $\text{ord}(\xi^{1/4}) = 4r$. Then from (35) it follows that $F_{U^\pm}^{\mathbb{Z}/2}(\xi) \sim \sqrt{2}$. Hence the denominator of (36) is $\sim (\sqrt{2})^m$.

According to [KM, p. 522], we may assume $\ell_{ij} \equiv 0 \pmod{2}$ if $i \neq j$. Since $\ell_{ij} \alpha_i \alpha_j$ appears twice in the exponent in (34) if $i \neq j$, we can write

$$\begin{aligned} F_{L \sqcup L'}^{\mathbb{Z}/2}(\xi; \mathbf{s}) &\sim \prod_{i=1}^m \left(\sum_{\alpha_i \in \{0,1\}} \xi^{\frac{1}{4}r(r-2)\ell_{ii}\alpha_i^2 + \frac{r}{2}\varepsilon_i \alpha_i} \right) \\ &= \prod_{i=1}^m \left(1 + \xi^{\frac{1}{4}r(r-2)\ell_{ii} + \frac{r}{2}\varepsilon_i} \right) . \end{aligned}$$

Since $\xi^{\frac{1}{4}r(r-2)\ell_{ii} + \frac{r}{2}\varepsilon_i}$ is a 4-th root of unity, each factor in the above product is either 2, 0, or $\sim \sqrt{2}$, and hence is divisible by $\sqrt{2}$. This means $F_{L \sqcup L'}^{\mathbb{Z}/2}(\xi; \mathbf{s})$, the numerator of (36), is divisible by $(\sqrt{2})^m$, and the statement follows. \square

5. DIAGONALIZATION OF 3-MANIFOLDS

We recall and refine some well-known facts about diagonalization of 3-manifolds. The first diagonalization result was obtained in [Oh] and was further developed in [Le3, BL, BBuL].

A 3-manifold is said to be *diagonal of prime type* if it can be obtained by surgery along a framed link $L \subset S^3$ with diagonal linking matrix $\text{diag}(b_1, \dots, b_m)$ such that $b_i = \pm p_i^{\varepsilon_i}$, where each p_i is a prime, 1, or 0. Denote by $\mathbf{L}(b, a)$ the lens space obtained from S^3 by surgery on the unknot with framing b/a . Also $M \# M'$ is the connected sum of M and M' and $M^{\#s}$ is the connected sum of s copies of M .

Proposition 5.1. *For every 3-manifold M , there exists a 3-manifold N of the form*

$$N = \mathbf{L}(2^{k_1}, -1) \# \dots \# \mathbf{L}(2^{k_j}, -1) ,$$

such that for every positive integer s , $M^{\#2s} \# N$ is diagonal of prime type.

To prepare for the proof we recall some well-known facts about linking pairing. A *linking pairing* on a finite abelian group G is a non-singular symmetric bilinear map from $G \times G$ to \mathbb{Q}/\mathbb{Z} . Two linking pairings ν, ν' on respectively G, G' are *isomorphic* if there is an isomorphism between G and G' carrying ν to ν' . With the obvious block sum, the set of equivalence classes of linking pairings is a semigroup.

Any non-singular $n \times n$ symmetric matrix B with integer entries gives rise to a linking pairing ϕ_B on $\mathbb{Z}^n/B\mathbb{Z}^n$ by $\phi_B(x, x') = x^t B^{-1} x' \in \mathbb{Q}/\mathbb{Z}$, where $x, x' \in \mathbb{Z}^n$ and x^t is the transpose of x . A linking pairing is *diagonal of type B* if it is isomorphic to ϕ_B , where B is a non-singular $n \times n$ diagonal matrix with integer entries.

An *enhancement* of an $n \times n$ symmetric matrix B is any matrix of the form $B \oplus D$, where D is a diagonal matrix with entries 0 or ± 1 on the diagonal.

For any closed oriented 3-manifold M , there is a linking pairing $\phi(M)$ on the torsion subgroup of $H_1(M, \mathbb{Z})$ defined by the Poincare duality, see [KK]. For example, if $b \neq 0$ is an integer, then the lens space $\mathbf{L}(b, 1)$ has linking pairing $\phi_{(b)}$, and $\mathbf{L}(b, -1)$ has linking pairing $\phi_{(-b)}$. Here (b) is the 1×1 matrix with entry b .

It is clear that $\phi(M \# M') = \phi(M) \oplus \phi(M')$. The result of [Le3, Section 3.5] shows the following.

Proposition 5.2. *If the linking pairing $\phi(M)$ on the torsion subgroup of $H_1(M, \mathbb{Z})$ is diagonal of type B , then M can be obtained from S^3 by surgery along an oriented framed link whose linking matrix is an enhancement of B .*

Proof of Proposition 5.1. In [BL, Section 2.2] it was noticed that $\phi(M \# M)$ is almost diagonal. More precisely,

$$\phi(M \# M) = \phi_B \oplus \nu, \quad (37)$$

where B is a diagonal matrix whose diagonal entries are prime powers and ν has the form

$$\nu = \bigoplus_{i=1}^j E_0^{k_i}.$$

Here E_0^k is a certain linking form on $\mathbb{Z}/2^k \times \mathbb{Z}/2^k$. We don't need the exact description of E_0^k . For us it is important that (see [KK])

$$E_0^k \oplus \phi_{(-2^k)} = \phi_{(-2^k)} \oplus \phi_{(2^k)} \oplus \phi_{(2^k)}. \quad (38)$$

Note that there is still one $\phi_{(-2^k)}$ in the right hand side of (38). From (38) and (37),

$$\phi(N \# (M \# M)^{\#s}) = s \phi_B \oplus \bigoplus_{i=1}^j (\phi_{(-2^{k_i})} \oplus 2s \phi_{(2^{k_i})}) = \phi_{B'}, \quad (39)$$

where B' is a diagonal matrix with diagonal entries of the form $\pm p^m$ with prime p . By Proposition 5.2, $N \# (M \# M)^{\#s}$ is diagonal of prime type. This completes the proof of Proposition 5.1. \square

6. PROOF OF THE INTEGRALITY IN THE $SO(3)$ CASE

Throughout this section $G = SO(3)$ and ξ is a root of unity of odd order r .

Proposition 6.1. *For integer $0 \leq k \leq (r-3)/2$, arbitrary integer b , and $\varepsilon \in \{0, 1\}$,*

$$\frac{H^{SO(3)}(k, b, \varepsilon)}{H^{SO(3)}(0, \pm 1, 0)} \in \mathbb{Z}[\xi^{1/4}, e_8], \quad (40)$$

and

$$\frac{H^{SO(3)}(k, 0, \varepsilon)}{(1 - \xi) \mathcal{D}^{SO(3)}} \in \mathbb{Z}[\xi^{1/4}, e_8]. \quad (41)$$

Proof. First note that by Lemmas 3.3 (e) and 3.5 (b), $O_\xi \sim H^{SO(3)}(0, \pm 1, 0)$. By Lemma 3.5 (a), there is $f_\varepsilon(z, q) \in I_{2k+1+\varepsilon} \subset \mathbb{Z}[z^{\pm 1}, q^{\pm 1}]$ such that

$$\frac{H^{SO(3)}(k, b, \varepsilon)}{H^{SO(3)}(0, \pm 1, 0)} \sim \frac{2 \sum_n \xi, SO(3) q^{\frac{b(n^2-1)}{4}} q^{\frac{-3\varepsilon n}{2}} f_\varepsilon(q^n, q)}{x_{2k+1+\varepsilon} O_\xi}. \quad (42)$$

Since r is odd, $(n^2 - 1)/4$ and $(1 - n)/2$ are integers, and there are integers $2^*, 4^*$ such that $2^* 2 \equiv 4^* 4 \equiv 1 \pmod{r}$. We then have $\xi^{(n^2-1)/4} = \xi^{4^*(n^2-1)}$, $\xi^{-3n/2} = \xi^{-3/2} \xi^{3(1-n)2^*}$.

The numerator of (42) is

$$\begin{aligned} 2 \sum_n \xi, SO(3) q^{\frac{b(n^2-1)}{4}} q^{\frac{-3\varepsilon n}{2}} f(q^n, q) &= \xi^{-3\varepsilon/2} \sum_{\substack{n=0 \\ n \text{ odd}}}^{2r-1} \xi^{4^*b(n^2-1)+3\varepsilon 2^*(1-n)} f_\varepsilon(\xi^n, \xi) \\ &= \xi^{-3\varepsilon/2} \sum_{n=0}^{r-1} \xi^{4^*b(n^2-1)+3\varepsilon 2^*(1-n)} f_\varepsilon(\xi^n, \xi), \end{aligned} \quad (43)$$

where the second identity follows by replacing odd $n \in [r, 2r - 1]$ with $n - r$, which is even and in $[0, r - 1]$.

By Proposition 3.1, the right hand side of (43) is divisible by the denominator of the right hand side of (42), and (40) follows.

Statement (41) follows from (40) with $b = 0$ and Lemma 3.5(c), which says that $H^{SO(3)}(0, \pm 1, 0) \sim (1 - \xi) \mathcal{D}^{SO(3)}$. \square

Proof of Theorem 2. By Proposition 6.1, each factor in the right hand side of (10) is in $\mathbb{Z}[\xi^{1/4}, e_8]$, hence $\tau_{M, L'}^{SO(3)}(\xi) \in \mathbb{Z}[\xi^{1/4}, e_8]$ if M is diagonal.

Now suppose M is an arbitrary 3-manifold. Let N be the manifold described in Proposition 5.1, for which $M \# M \# N$ is diagonal. Since the WRT invariant is multiplicative with respect to connected sum, we get

$$\left(\tau_{M, L'}^{SO(3)}(\xi) \right)^2 \tau_N^{SO(3)}(\xi) \in \mathbb{Z}[\xi^{1/4}, e_8].$$

Since 2^k is coprime to r , $\tau_{\mathbf{L}(2^k, -1)}^{SO(3)}(\xi)$ is a unit in $\mathbb{Z}[\xi^{1/4}]$ by Lemma 3.6 (a). It follows that $\tau_N^{SO(3)}(\xi)$ is a unit, hence $\left(\tau_{M, L'}^{SO(3)}(\xi) \right)^2 \in \mathbb{Z}[\xi^{1/4}, e_8]$. By Lemma 3.3 (c), $\tau_{M, L'}^{SO(3)}(\xi) \in \mathbb{Z}[\xi^{1/4}, e_8]$. This completes the proof of the theorem. \square

7. PROOF OF THE INTEGRALITY IN THE $SU(2)$ CASE

If the order of ξ is odd, then by the splitting property (Proposition 4.9),

$$\tau_{M, L'}^{SU(2)}(\xi) = \tau_{M, L'}^{\mathbb{Z}/2}(\xi) \tau_{M, L'}^{SO(3)}(\xi).$$

Both factors of the right hand side are algebraic integers by Theorem 2 and Proposition 4.9. Hence $\tau_{M, L'}^{SU(2)}(\xi)$ is also an algebraic integer.

Therefore throughout the remaining part of this section we will assume that $r = \text{ord}(\xi)$ is *even*. Note that in this case the order of $\xi^{1/4}$ is always $4r$ and $e_8 \in \mathbb{Z}[\xi^{1/4}]$.

Proposition 7.1. *Let $r = \text{ord}(\xi)$ be even. Suppose $b = \pm p^s$, where p is 0, 1 or a prime, k an integer, and $\varepsilon \in \{0, 1\}$. Then*

$$\frac{H^{SU(2)}(k, b, \varepsilon)}{H^{SU(2)}(0, \pm 1, 0)} \in \mathbb{Z}[\xi^{1/4}] \quad \text{and} \quad \frac{H^{SU(2)}(k, 0, \varepsilon)}{(1 - \xi) \mathcal{D}^{SU(2)}} \in \mathbb{Z}[\xi^{1/4}] .$$

The following lemma will be used in the proof of the above proposition for odd b .

Lemma 7.2. *Suppose b is odd, r is even, $a \in \mathbb{Z}$, and $f \in I_k$. Then*

$$A := \frac{\sum_n \xi, SU(2) q^{\frac{bn^2}{4} + \frac{an}{2}} f(q^n, q)}{x_k O_\xi}$$

belongs to $\mathbb{Z}_{(2)}[\xi^{1/4}]$, where $\mathbb{Z}_{(2)}$ is the set of all rational numbers with odd denominators.

Proof. Let $r = r_e r_o$, where r_o is odd, and r_e is a power of 2. Then $\text{ord}(\xi^{1/4}) = 4r = (4r_e)r_o$, with $4r_e$ and r_o coprime. By definition,

$$\begin{aligned} \sum_n \xi, SU(2) q^{\frac{bn^2}{4} + \frac{an}{2}} q^{nj} &= \frac{1}{4} G(b, 4j + 2a, \xi^{1/4}) \quad \text{by Proposition 3.4 (d)} \\ &= \frac{1}{4} G(br_o, 4j + 2a, \xi^{r_o/4}) G(4br_e, 4j + 2a, \xi^{r_e}) \\ &= \frac{1}{4} \xi^{da^2/4} G(br_o, 0, \xi^{r_o/4}) \xi^{d(j^2+aj)} G(4br_e, 4j + 2a, \xi^{r_e}) , \end{aligned} \quad (44)$$

where d is any multiple of r_o such that $db \equiv -1 \pmod{4r_e}$.

Let us extend $\Delta(z) = z \otimes z$ to a $\mathbb{Z}[q^{\pm 1}]$ -algebra homomorphism

$$\Delta : \mathbb{Z}[z^{\pm 1}, q^{\pm 1}] \rightarrow \mathbb{Z}[z^{\pm 1}, q^{\pm 1}] \otimes_{\mathbb{Z}[q^{\pm 1}]} \mathbb{Z}[z^{\pm 1}, q^{\pm 1}] .$$

Define $Q(j) = dj^2 + da j$. Also define $\mathbb{Z}[q^{\pm 1/4}]$ -module homomorphism $T : \mathbb{Z}[z^{\pm 1}, q^{\pm 1/4}] \rightarrow \mathbb{Z}[q^{1/4}]$ by:

$$T(z^j) = G(4br_e, 4j + 2a, \xi^{r_e}) .$$

Using (44) we can rewrite

$$\sum_n \xi, SU(2) q^{\frac{bn^2}{4} + \frac{an}{2}} f(q^n, q) = \frac{\xi^{da^2/4}}{4} G(br_o, 0, \xi^{r_o/4}) \text{ev}_\xi \left\{ (\mathcal{L}_Q \otimes T)(\Delta f) \right\} . \quad (45)$$

It is enough to consider the case $f = z^m(q^l z; q)_k$. Applying Corollary 2.7 to $\Delta(z^m(q^l z; q)_k)$, using $z_1 = z \otimes 1$ and $z_2 = 1 \otimes z$, we see that A is a $\mathbb{Z}[\xi^{\pm 1/4}]$ -linear combination of terms of the form

$$B = \left(\frac{G(br_o, 0, \xi^{r_o/4})}{4O_\xi} \right) \left(\frac{(\xi; \xi)_k \text{ev}_\xi \{ \mathcal{L}_Q(z^m(z; q)_{k_1}) \}}{x_k(\xi; \xi)_{k_1}} \right) \left(\frac{T(z^m(z; q)_{k_2})}{(\xi; \xi)_{k_2}} \right) \quad (46)$$

with $k_i \leq k$. There are three factors on the RHS of (46) and we will show that each factor belongs to $\mathbb{Z}_{(2)}[\xi^{1/4}]$. The last factor in B is in $\mathbb{Z}[\xi]$. In fact, if $z^m(z; q)_{k_2} = \sum_j c_j(z) z^j$ then

$$\begin{aligned} T(z^m(z; q)_{k_2}) &= \sum_j c_j(\xi) G(4br_e, 4j + 2a, \xi^{r_e}) = \sum_j c_j(\xi) \sum_{n=0}^{r_o-1} \xi^{2r_e(br_e n^2 + (2j+a)n)} \\ &= \sum_n \xi^{4br_e^2 n^2 + 2r_e a n} \sum_j c_j(\xi) \xi^{4r_e n j} = \sum_n \xi^{4br_e^2 n^2 + 2r_e n(a+2m)} (\xi^{4r_e n}; \xi)_{k_2} , \end{aligned}$$

which is divisible by $(\xi; \xi)_{k_2}$ in $\mathbb{Z}[\xi]$. This shows that the last factor of (46) is in $\mathbb{Z}_{(2)}[\xi^{1/4}]$.

By Theorem 2.2, $\text{ev}_\xi(\mathcal{L}_Q(z^m(z; q)_{k_1}))$ is in $x_{k_1} \mathbb{Z}[\xi]$. Hence the second factor is in

$$\frac{(\xi; \xi)_k x_{k_1}}{x_k (\xi; \xi)_{k_1}} \mathbb{Z}[\xi] = \frac{(\xi; \xi)_{\lfloor k/2 \rfloor}}{(\xi; \xi)_{\lfloor k_1/2 \rfloor}} \mathbb{Z}[\xi] \subset \mathbb{Z}[\xi].$$

By Proposition 3.4 (b), $G^2(br_o, 0, \xi^{r_o/4}) \sim 8r_e$, while $O_\xi^2 \sim r/2$ by Lemma 3.3 (e). It follows that the square of the first factor, and hence the first factor itself, is in $\frac{r_e}{r} \mathbb{Z}[\xi] = \frac{1}{r_o} \mathbb{Z}[\xi] \subset \mathbb{Z}_{(2)}[\xi]$. Here we use the fact that $\mathbb{Z}_{(2)}[\xi]$ is integrally closed.

We can conclude that B , and hence A , is in $\mathbb{Z}_{(2)}[\xi]$. \square

Proof of Proposition 7.1. By Lemma 3.5 there is $f_\varepsilon(z, q) \in I_{2k+1+\varepsilon}$ such that

$$\frac{H^{SU(2)}(k, b, \varepsilon)}{H^{SU(2)}(0, \pm 1, 0)} \sim \frac{\frac{1}{4} \sum_{n=0}^{4r-1} \xi^{\frac{b}{4}(n^2-1)} \xi^{-3\varepsilon n/2} f_\varepsilon(\xi^n, \xi)}{x_{2k+1+\varepsilon} O_\xi}. \quad (47)$$

We split the proof into 3 cases: (1) $b \equiv 0 \pmod{4}$; (2) $b = \pm 2$ and (3) b is odd.

(1) $b = 4b', b' \in \mathbb{Z}$. Since $H^{SU(2)}(k, b, 1) = 0$ by Lemma 3.5 (d), we can assume $\varepsilon = 0$. By (47),

$$\frac{H^{SU(2)}(k, b, 0)}{H^{SU(2)}(0, \pm 1, 0)} \sim \frac{\frac{1}{4} \sum_{n=0}^{4r-1} \xi^{b'(n^2-1)} f_0(\xi^n, \xi)}{x_{2k+1} O_\xi} = \frac{\sum_{n=0}^{r-1} \xi^{b'(n^2-1)} f_0(\xi^n, \xi)}{x_{2k+1} O_\xi},$$

which is in $\mathbb{Z}[\xi]$ by Proposition 3.1.

(2) $b = \pm 2$. Again $H^{SU(2)}(k, b, 1) = 0$ by Lemma 3.5 (d), and we can assume $\varepsilon = 0$. This case was studied in [BBIL], where the exact value of $H^{SU(2)}(k, \pm 2, 0)$ was obtained. By Lemma 5.2 in [BBIL] we have

$$H^{SU(2)}(k, \pm 2, 0) \sim 2\sqrt{r} \prod_{i=0}^k \frac{1 - \xi^{(2i+1)/2}}{1 - \xi^{2i+1}} = 2\sqrt{r} \prod_{i=0}^k \frac{1}{1 + \xi^{(2i+1)/2}}.$$

Hence from Lemma 3.5, with $k \leq r/2 - 1$,

$$\frac{H^{SU(2)}(k, \pm 2, 0)}{H^{SU(2)}(0, \pm 1, 0)} \sim \frac{\sqrt{r}/O_\xi}{\prod_{i=0}^k (1 + \xi^{(2i+1)/2})} \in z \mathbb{Z}[\xi^{1/4}]$$

where

$$z = \frac{\sqrt{r}/O_\xi}{\prod_{i=0}^{r/2-1} (1 + \xi^{(2i+1)/2})}.$$

The square of the numerator of z is $r/O_\xi^2 \sim 2$, by Lemma 3.3.

Let us calculate the square of the denominator. For any integer j one has

$$(1 + \xi^{(2j+1)/2}) \sim (1 - \xi^{(2j+1)/2}).$$

Hence

$$\begin{aligned} \left(\prod_{j=0}^{r/2-1} (1 + \xi^{(2j+1)/2}) \right)^2 &\sim \prod_{j=0}^{r/2-1} (1 + \xi^{(2j+1)/2})(1 - \xi^{(2j+1)/2}) = \prod_{j=0}^{r/2-1} (1 - \xi^{(2j+1)}) \\ &\sim \frac{\prod_{j=1}^{r-1} (1 - \xi^j)}{\prod_{j=1}^{r/2-1} (1 - \xi^{2j})} = \frac{r}{r/2} = 2. \end{aligned}$$

We can conclude that $\frac{H^{SU(2)}(k, \pm 2, 0)}{H^{SU(2)}(0, \pm 1, 0)} \in \mathbb{Z}[\xi^{1/4}]$.

(3) Assume that b is odd. Splitting the sum in the numerator of the right hand side of (47) into even and odd n we get

$$\begin{aligned}
& \frac{1}{4} \sum_{n=0}^{4r-1} \xi^{b(n^2-1)/4} \xi^{-3\epsilon n/2} f_\epsilon(\xi^n, \xi) \\
&= \frac{1}{4} \left\{ \xi^{-b/4} \sum_{n=0}^{2r-1} \xi^{bn^2-3\epsilon n} f_\epsilon(\xi^{2n}, \xi) + \xi^{-3\epsilon/2} \sum_{n=0}^{2r-1} \xi^{b(n^2+n)-3\epsilon n} f_\epsilon(\xi^{2n+1}, \xi) \right\} \\
&= \frac{1}{2} \left\{ \xi^{-b/4} \sum_{n=0}^{r-1} \xi^{bn^2-3\epsilon n} f_\epsilon(\xi^{2n}, \xi) + \xi^{-3\epsilon/2} \sum_{n=0}^{r-1} \xi^{b(n^2+n)-3\epsilon n} f_\epsilon(\xi^{2n+1}, \xi) \right\} \quad (48)
\end{aligned}$$

Since $f_\epsilon(z^2, q)$ and $f_\epsilon(z^2q, q)$ belong to $I_{2k+1+\epsilon}$ (according to Proposition 2.1), each summand in the curly brackets of the right hand side of (48) is divisible by $x_{2k+1+\epsilon}O_\xi$, by Proposition 3.1. It follows from (47) that

$$\frac{H^{SU(2)}(k, b, \epsilon)}{H^{SU(2)}(0, \pm 1, 0)} \in \frac{1}{2} \mathbb{Z}[\xi^{1/4}],$$

which, together with Lemma 7.2, implies

$$\frac{H^{SU(2)}(k, b, \epsilon)}{H^{SU(2)}(0, \pm 1, 0)} \in \frac{1}{2} \mathbb{Z}[\xi^{1/4}] \cap \mathbb{Z}_{(2)}[\xi^{1/4}] = \mathbb{Z}[\xi^{1/4}].$$

Finally

$$\frac{H^{SU(2)}(k, 0, \epsilon)}{(1 - \xi) \mathcal{D}^{SU(2)}} \in \mathbb{Z}[\xi^{1/4}]$$

follows from Lemma 3.5 (c), which says that $H^{SU(2)}(0, \pm 1, 0) \sim (1 - \xi) \mathcal{D}^{SU(2)}$. \square

Proof of Theorem 1. By Proposition 7.1, each factor in the right hand side of (10) is in $\mathbb{Z}[\xi^{1/4}]$, hence $\tau_{M, L'}^{SU(2)}(\xi) \in \mathbb{Z}[\xi^{1/4}]$ if M is diagonal of prime type.

Now suppose M is an arbitrary 3-manifold. According to Proposition 5.1, there exist lens spaces $\mathbf{L}(2^{k_1}, -1), \dots, \mathbf{L}(2^{k_j}, -1)$, such that $M \#^{2s} \# N$ is diagonal of prime type for every positive integer s . Here

$$N := \#_{i=1}^j \mathbf{L}(2^{k_i}, -1).$$

By Lemma 3.6, there is an odd colored knot $K_i \subset \mathbf{L}(2^{k_i}, -1)$ such that $\tau_{\mathbf{L}(2^{k_i}, -1), K_i}^{SU(2)} \neq 0$. The knots K_i together form a link $L'' \subset N$, and

$$\tau_{N, L''}^{SU(2)}(\xi) = \prod_i \tau_{\mathbf{L}(2^{k_i}, -1), K_i}^{SU(2)} \neq 0.$$

Taking the connected sum of (N, L'') with $2s$ copies of (M, L') , we get a diagonal 3-manifold of prime type. Hence,

$$\left(\tau_{M, L'}^{SU(2)}(\xi) \right)^{2s} \tau_{N, L''}^{SU(2)}(\xi) \in \mathbb{Z}[\xi^{1/4}]$$

for every positive integer s . Applying Lemma 3.3 (c), we get $\tau_{M, L'}^{SU(2)}(\xi) \in \mathbb{Z}[\xi^{1/4}]$. \square

APPENDIX A. PROOF OF THEOREM 1.1

A.1. Algebraic preliminaries. We first recall the universal quantized algebra $U_h = U_h(sl_2)$ and some of its properties. For more details see e.g. [Ha2].

The universal quantized algebra $U_h = U_h(sl_2)$ is the h -adically complete $\mathbb{Q}[[h]]$ -algebra, topologically generated by the elements H, E , and F , satisfying the relations

$$HE - EH = 2E, \quad HF - FH = -2F, \quad EF - FE = \frac{K - K^{-1}}{v - v^{-1}},$$

where $K := \exp(hH/2)$, $v := \exp(h/2)$, and $v^2 = q$. The algebra U_h has a structure of Hopf algebra, which makes U_h into a U_h -module via the adjoint representation, and defines a tensor product on the set of U_h -modules. In particular, the completed tensor powers $U_h^{\hat{\otimes} m}$ is a U_h -module via the adjoint representation. For a set $Y \subset U_h^{\hat{\otimes} m}$ its subset of invariant elements is defined by

$$Y^{\text{inv}} := \{y \in Y \mid a \cdot y = \epsilon(y), \quad \forall a \in U_h\},$$

where ϵ is the antipode of U_h and $a \cdot y$ is the adjoint action. It is known that $(U_h)^{\text{inv}}$ is exactly the center of U_h .

For each positive integer n there is a unique n -dimensional irreducible U_h -module, denoted by V_n , we set $V := V_2$. Let

$$R = \text{Span}_{\mathbb{Z}[v^{\pm 1}]} \{V_n, n \geq 1\},$$

which is a $\mathbb{Z}[v^{\pm 1}]$ -algebra whose multiplication is the tensor product. One has

$$V_n V = V_{n+1} + V_{n-1}, \tag{A.1}$$

and as a ring $R = \mathbb{Z}[v^{\pm 1}][V]$, the $\mathbb{Z}[v^{\pm 1}]$ -polynomial algebra in V .

For an U_h -module W and $x \in U_h$ the quantum trace is defined by

$$\text{tr}_q^W(x) = \text{tr}(xK^{-1}, W),$$

which can be linearly extended to the case when W is a $\mathbb{Z}[v^{\pm 1}]$ -linear combination of U_h -modules.

The quantum trace preserves ad-invariance, which means the following. Suppose $W \in R$ and $y \in (U_h^{\hat{\otimes} m})^{\text{inv}}$, then $(\text{id}^{\otimes(m-1)} \otimes \text{tr}_q^W)(y) \in (U_h^{\hat{\otimes}(m-1)})^{\text{inv}}$.

A.2. New bases for R . In R consider the following elements: $P_0^{(0)} = P_0^{(1)} = 1$,

$$P_n^{(0)} = \prod_{j=1}^n (V - \lambda_{2j-1}), \quad P_n^{(1)} = \prod_{j=1}^n (V - \lambda_{2j}),$$

where $\lambda_n = v^n + v^{-n}$. Note that $P_n^{(0)}$ is P_n of [Ha2]. Since $P_n^{(0)}$ is a monic polynomial of degree n in V with coefficients in $\mathbb{Z}[v^{\pm 1}]$, it is clear that the set $\{P_n^{(0)}, n = 0, 1, 2, \dots\}$ forms a $\mathbb{Z}[v^{\pm 1}]$ -basis of R . Similarly, $\{P_n^{(1)}, n = 0, 1, 2, \dots\}$ also forms a $\mathbb{Z}[v^{\pm 1}]$ -basis of R . It is not difficult to express V_n through these bases. In fact, (A.1), together with an easy induction, will give the following identities, the first of which was obtained in [Ha2].

$$V_n = \sum_{k=0}^{n-1} \begin{bmatrix} n+k \\ 2k+1 \end{bmatrix} P_k^{(0)}, \quad V_n = \sum_{k=0}^{n-1} \begin{bmatrix} n+k \\ 2k+1 \end{bmatrix} \frac{\lambda_n}{\lambda_{k+1}} P_k^{(1)}. \tag{A.2}$$

A.3. Integral subalgebras and their completions. Following [Ha2] let $\mathcal{U}_q^{(0)}$ be the $\mathbb{Z}[q^{\pm 1}]$ -subalgebra of U_h generated by $\tilde{F}^{(l)}$, e and $K^{\pm 2}$, where

$$\tilde{F}^{(l)} := q^{l(1-l)/4} F^l K^l / [l]! \quad \text{and} \quad e := \{1\}E .$$

Let $\mathcal{U}_q^{(1)} = K \mathcal{U}_q^{(0)}$ and $\mathcal{U}_q = \mathcal{U}_q^{(0)} \oplus \mathcal{U}_q^{(1)}$.

Let $\mathcal{F}_p(\mathcal{U}_q^{\otimes m}) \subset \mathcal{U}_q^{\otimes m}$ be the $\mathbb{Z}[q^{\pm 1}]$ -span of elements of the form $y_1 \otimes y_2 \otimes \cdots \otimes y_m$, where each y_j belongs to \mathcal{U}_q , and one of them belongs to $\mathcal{U}_q e^p \mathcal{U}_q$. For a set $Y \subset \mathcal{U}_q^{\otimes m}$ define its completion

$$\tilde{Y} := \left\{ \sum_{j=0}^{\infty} z_p \mid z_p \in Y \cap \mathcal{F}_p(\mathcal{U}_q^{\otimes m}) \right\} .$$

In particular, when $m = 1$, one can define $\tilde{\mathcal{U}}_q$ and $\tilde{\mathcal{U}}_q^{(\varepsilon)}$ for $\varepsilon \in \{0, 1\}$. For $\mathbf{e} = (\varepsilon_1, \dots, \varepsilon_m) \in \{0, 1\}^m$, let $\widetilde{\mathcal{U}}_q^{(\mathbf{e})}$ be the completion of $\mathcal{U}_q^{(\varepsilon_1)} \otimes \cdots \otimes \mathcal{U}_q^{(\varepsilon_m)}$ defined as above.

The center $Z(\mathcal{U}_q) = (\mathcal{U}_q)^{\text{inv}}$ is freely generated as a $\mathbb{Z}[q^{\pm 1}]$ -algebra by the quantum Casimir operator

$$C = (1 - q^{-1})\tilde{F}^{(1)}K^{-1}e + K + q^{-1}K^{-1} \in \mathcal{U}_q^{(1)} .$$

Set

$$\sigma_n^{(0)} = \prod_{i=0}^n (qC^2 - (q^i + 2 + q^{-i})) \quad \text{and} \quad \sigma_n^{(1)} = C\sigma_n^{(0)} .$$

Theorem 1.1 in [Ha1] states that

$$(\tilde{\mathcal{U}}_q^{(\varepsilon)})^{\text{inv}} = \left\{ \sum_{p \geq 0} a_p \sigma_p^{(\varepsilon)} \mid a_p \in \mathbb{Z}[q^{\pm 1}] \right\} . \quad (\text{A.3})$$

We will need the following result.

Proposition A.1. *Suppose $x \in \mathcal{U}_q^{(\varepsilon)}$, $\varepsilon \in \{0, 1\}$. Then for every n , $\text{tr}_q^{P_n^{(\varepsilon)}}(x)$ belongs to $(q; q)_n \mathbb{Z}[q^{\pm 1}]$.*

Proof. If $\varepsilon = 0$, this is [Ha2, Lemma 8.5]. The case $\varepsilon = 1$ can be proved similarly. It is enough to set $x = \tilde{F}^{(l)} K^{2j+1} e^{l'}$. It is easy to see that $\text{tr}_q^{P_n^{(1)}}(\tilde{F}^{(l)} K^{2j+1} e^{l'}) = 0$ if $l \neq l'$. Set

$$B(n, l, j) := \text{tr}_q^{P_n^{(1)}}(\tilde{F}^{(l)} K^{2j+1} e^{l'}) .$$

Then it is clear that $B(n, l, j) = 0$ when $l > n$ because e^l vanishes on V_1, V_2, \dots, V_{n+1} . When $l \leq n$, by a similar argument as in the proof of [Ha2, lemma 8.8], we have

$$B(n, l, j) = \{j - n\}\{j + n\}B(n - 1, l, j) + q^j(1 - q^{-l})B(n - 1, l - 1, j + 1) .$$

The above recursive relation and a simple induction will show that

$$B(n, l, j) = q^{-(j+l)n} (q; q)_n (q; q)_{n-l} \binom{j-1}{n-l}_q \binom{j+n}{n-l}_q \in (q; q)_n \mathbb{Z}[q^{\pm 1}] .$$

□

Lemma A.2. *For every non-negative integers k, p and $\varepsilon \in \{0, 1\}$, one has*

$$\text{tr}_q^{P_k^{(\varepsilon)}}(\sigma_p^{(\varepsilon)}) = \delta_{k,p} \frac{\{2k+1\}!}{v^\varepsilon \{1\}} \lambda_{k+1}^\varepsilon . \quad (\text{A.4})$$

Proof. The case $\varepsilon = 0$ is proved in [Ha2, Proposition 6.3]. Hence, we restrict to $\varepsilon = 1$.

As explained in [Ha2, Section 6.3.1], there exists a homomorphism $\varphi : R \rightarrow Z(\mathcal{U}_q \otimes \mathbb{Z}[v^{\pm 1}])$ sending V to vC . In particular, for

$$S_n^{(1)} := V \prod_{j=1}^n (V^2 - (\lambda_j)^2)$$

we have $\varphi(S_n^{(1)}) = v\sigma_n^{(1)}$. Moreover, for any $x, y \in R$,

$$\mathrm{tr}_q^x(\varphi(y)) = J_{\mathcal{H}}(x, y) := \langle x, y \rangle ,$$

where $\langle x, y \rangle$ is the Rosso pairing defined as the colored Jones polynomial of the 0-framed Hopf link \mathcal{H} , whose two components are colored by x and y . Note that this pairing is symmetric. Hence, for $\varepsilon = 1$, the left hand side of (A.4) is equal to $\langle P_k^{(1)}, v^{-1}S_p^{(1)} \rangle$.

Since $\lambda_n := v^n + v^{-n} = \langle V_n, V \rangle / [n]$, for every $f(V) \in R$ we have

$$\langle V_n, f(V) \rangle = [n]f(\lambda_n) . \quad (\text{A.5})$$

Hence if $m < n$, then $\langle V_{2m+2}, P_n^{(1)} \rangle = 0$ and $\langle S_n^{(1)}, V_{m+1} \rangle = 0$.

Using $V_n V = V_{n+1} + V_{n-1}$, we get

$$P_n^{(1)} = V_{n+1} + \text{a } \mathbb{Z}[v^{\pm 1}]\text{-linear combination of } V_1, V_2, \dots, V_n , \quad (\text{A.6})$$

and

$$S_n^{(1)} = V_{2n+2} + \text{a } \mathbb{Z}[v^{\pm 1}]\text{-linear combination of } V_2, V_4, \dots, V_{2n} .$$

Therefore $\langle S_m^{(1)}, P_n^{(1)} \rangle = 0$ if $m \neq n$.

Finally

$$\begin{aligned} v \mathrm{tr}_q^{P_n^{(1)}}(\sigma_n^{(1)}) &= \langle S_n^{(1)}, P_n^{(1)} \rangle = \langle S_n^{(1)}, V_{n+1} \rangle = [n+1]\lambda_{n+1} \prod_{j=1}^n ((\lambda_{n+1})^2 - (\lambda_j)^2) \\ &= [n+1]\lambda_{n+1} \prod_{j=1}^n \{j\} \{2n+2-j\} \end{aligned}$$

□

A.4. Proof of Theorem 1.1. Suppose $L \sqcup L'$ is an oriented framed link with fixed colors $\mathbf{s} = (s_1, \dots, s_l)$ on L' . Here L has m ordered components and ε_i are defined in (2).

According to [Ha2, Theorem 4.1], there is an element $J_T \in (U_h^{\otimes(m+l)})^{\mathrm{inv}}$ such that

$$J_L(\mathbf{n}) = \mathrm{tr}_q^{V_{n_1} \otimes \dots \otimes V_{n_m} \otimes V_{s_1} \otimes \dots \otimes V_{s_l}}(J_T) .$$

(In [Ha2], J_T is the universal invariant of a bottom tangle whose closure is $L \sqcup L'$.)

Using (A.2) to express V_{n_i} as a linear combination of $P_k^{(\varepsilon_i)}$, we have

$$\begin{aligned} J_L(\mathbf{n}) &= \sum_{k_i=0}^{n_i-1} \mathrm{tr}_q^{P_{k_1}^{(\varepsilon_1)} \otimes \dots \otimes P_{k_m}^{(\varepsilon_m)} \otimes V_{s_1} \otimes \dots \otimes V_{s_l}}(J_T) \prod_{i=1}^m \begin{bmatrix} n_i + k_i \\ 2k_i + 1 \end{bmatrix} \frac{\lambda_{n_i}^{\varepsilon_i}}{\lambda_{k_i+1}^{\varepsilon_i}} \\ &= \sum_{k_i=0}^{n_i-1} \mathrm{tr}_q^{P_{k_1}^{(\varepsilon_1)'} \otimes \dots \otimes P_{k_m}^{(\varepsilon_m)'} \otimes V_{s_1} \otimes \dots \otimes V_{s_l}}(J_T) \prod_{i=1}^m \begin{bmatrix} n_i + k_i \\ 2k_i + 1 \end{bmatrix} \{k_i\}! \frac{\lambda_{n_i}^{\varepsilon_i}}{\lambda_{k_i+1}^{\varepsilon_i}} , \end{aligned} \quad (\text{A.7})$$

which is (4) with

$$c_{L \sqcup L'}(\mathbf{k}) = \mathrm{tr}_q^{P_{k_1}^{(\varepsilon_1)'} \otimes \dots \otimes P_{k_m}^{(\varepsilon_m)'} \otimes V_{s_1} \otimes \dots \otimes V_{s_l}}(J_T) .$$

Here $P_k^{(\varepsilon)'} = P_k^{(\varepsilon)} / \{k\}!$.

Without loss of generality, we may assume $k_1 = k = \max(k_1, \dots, k_m)$.

By Theorem A.3 in [BBuL], $(\text{id}^{\otimes m} \otimes \text{tr}_q^{V_{s_1}} \otimes \dots \otimes \text{tr}_q^{V_{s_l}})(J_T) \in q^a (\mathcal{U}_q^{\otimes(\mathbf{e})})^{\text{inv}}$, for some $a \in \frac{1}{4}\mathbb{Z}$. Let

$$y := (\text{id} \otimes \text{tr}_q^{P^{(\varepsilon_2)'}} \otimes \dots \otimes \text{tr}_q^{P^{(\varepsilon_m)'}} \otimes \text{tr}_q^{V_{s_1}} \otimes \dots \otimes \text{tr}_q^{V_{s_l}})(J_T) .$$

Then

$$c_{L \sqcup L'}(\mathbf{k}) = \text{tr}_q^{P^{(\varepsilon_1)'}}(y) .$$

Proposition A.1, as well as the fact that quantum trace preserves ad-invariance, gives us $y \in q^a (\mathcal{U}_q^{(\varepsilon_1)})^{\text{inv}}$. Hence y has a presentation $y = q^a \sum_{p \geq 0} d_p \sigma_p^{(\varepsilon_1)}$ with $d_p \in \mathbb{Z}[q^{\pm 1}]$. We then have

$$\text{tr}_q^{P^{(\varepsilon_1)'}}(y) = q^a \sum_p d_p \text{tr}_q^{P^{(\varepsilon_1)'}}(\sigma_p^{(\varepsilon_1)}) ,$$

which belongs to $\frac{(q^{k+1}; q)_{k+1}}{1-q} \mathbb{Z}[q^{\pm \frac{1}{4}}]$ by Lemma A.2. \square

REFERENCES

- [A] G. Andrews *q-series: their development and applications in analysis, number theory, combinatorics, physics, and computer algebra*, Regional conference series in mathematics, Amer. Math. Soc. **66** (1985)
- [BBuL] A. Beliakova, I. Bühler and T. Le, *A unified quantum $SO(3)$ invariant for rational homology 3-spheres*, Inventiones Math. to appear, arXiv:0801.3893
- [BBIL] A. Beliakova, C. Blanchet and T. Le, *Unified quantum invariants and their refinements for homology 3-spheres with 2-torsion*, Fund. Math. **201(3)** (2008), 217–239
- [BL] A. Beliakova and T. Le, *Integrality of quantum 3-manifold invariants and a rational surgery formula*, Compos. Math. **143(6)** (2007), 1593–1612
- [D] H. Davenport, *Multiplicative Number Theory*, Graduate Texts in Mathematics, **74**, Springer-Verlag, New York, 2000
- [G] P. Gilmer, *Integrality for TQFTs*, Duke Math. J. **125** (2004), 389–413
- [De1] F. Deloup, *Linking forms, reciprocity for Gauss sums and invariants of 3-manifolds*, Trans. Amer. Math. Soc. **351** (1999), 1895–1918
- [De2] F. Deloup, *On abelian quantum invariants of links in 3-manifolds*, Math. Ann. **319** (2001), 759–795
- [GM] P. Gilmer and G. Masbaum, *Integral lattices in TQFT*, Ann. Sci. École Norm. **40** (2007), 815–844
- [Ha1] K. Habiro, *An integral form of the quantized enveloping algebra of sl_2 and its completions*, J. Pure Appl. Algebra **211(1)** (2007), 265–292
- [Ha2] K. Habiro, *A unified Witten-Reshetikhin-Turaev invariant for integral homology spheres*, Invent. Math. **171(1)** (2008), 1–81
- [HL] K. Habiro and T. Le, in preparation
- [KK] A. Kawachi and S. Kojima, *Algebraic classification of linking pairings on 3-manifolds*, Math. Ann. **253** (1980), 29–42
- [Kho] M. Khovanov, *Hopfological algebra and categorification at a root of unity: the first steps*, math.QA/0509083
- [KM] R. Kirby and P. Melvin, *The 3-manifold invariants of Witten and Reshetikhin-Turaev for $sl(2, \mathbb{C})$* , Invent. Math. **105** (1991), 473–545
- [La] S. Lang, *Algebra*, 3rd edition, Addison-Wesley 1997
- [Le1] T. Le, *Integrality and symmetry of quantum link invariants*, Duke Math. J., **102** (2000) 273–306
- [Le2] T. Le, *Quantum invariants of 3-manifolds: integrality, splitting, and perturbative expansion*, Topology Appl. **127** (2003), 125–152
- [Le3] T. Le, *Strong integrality of quantum invariants of 3-manifolds*, Trans. Amer. Math. Soc. **360(6)** (2008), 2941–2963
- [Lic] R. Lickorish, *An Introduction to Knot Theory*, Graduate Texts in Mathematics **175** 1997
- [Lu] G. Lusztig, *Quantum group at roots of 1*, Geom. Dedicata **35** (1990), 89–113.

- [MR] G. Masbaum and J. Roberts, *A simple proof of integrality of quantum invariants at prime roots of unity*, Math. Proc. Camb. Phil. Soc. **121** (1997), 443–454.
- [MW] G. Masbaum and H. Wenzl, *Integral modular categories and integrality of quantum invariants at roots of unity of prime order*, J. Reine Angew. Math. **505** (1998), 209–235.
- [MM] P. Melvin and H. Morton, *The coloured Jones function*, Comm. Math. Phys., **169**, 1995, 501–520.
- [Mu] H. Murakami, *Quantum $SO(3)$ -invariants dominate the $SU(2)$ -invariant of Casson and Walker*, Math. Proc. Camb. Phil. Soc. **117** (1995), 237–249.
- [MOO] H. Murakami, T. Ohtsuki and M. Okada, *Invariants of three-manifolds derived from linking matrices of framed links*, Osaka J. Math. **29** (1992), 545–572.
- [Oh] T. Ohtsuki, *A polynomial invariant of rational homology 3-spheres*, Invent. Math. **123** (1996), 241–257.
- [O] T. Ohtsuki, *Quantum invariants. A study of knots, 3-manifolds, and their sets*, Series on Knots and Everything, **29**, World Scientific Publishing Co., Inc., River Edge, NJ, 2002.
- [RT] N. Reshetikhin and V. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. **103** (1991), 547–597.
- [TY] T. Takata, Y. Yokota, *The $PSU(n)$ invariants of 3-manifolds are polynomials*, J. Knot Theory Ram. **8** (1999), 521–532.
- [Tu] V. Turaev, *Quantum invariants of knots and 3-manifolds*, de Gruyter Studies in Mathematics 18, Walter de Gruyter, Berlin New York 1994.
- [Wa] C. Wall, *Quadratic forms on finite groups, and related topics*, Topology **2** (1963), 281–298.
- [Wi] E. Witten, *Quantum field theory and the Jones polynomial*, Comm. Math. Phys. **121** (1989), 351–399.

I-MATH, UNIVERSITY OF ZURICH
 WINTERTHURERSTRASSE 190
 8057 ZURICH, SWITZERLAND
E-mail address: `anna@math.uzh.ch`

DEPARTMENT OF MATHEMATICS
 WINSTON-SALEM STATE UNIVERSITY
 WINSTON SALEM, NC 27110 USA
E-mail address: `chenqi@wssu.edu`

DEPARTMENT OF MATHEMATICS
 GEORGIA INSTITUTE OF TECHNOLOGY
 ATLANTA, GA 30332-0160, USA
E-mail address: `letu@math.gatech.edu`